SPECTRAL ANTISYMMETRY OF TWISTED GRAPH ADJACENCY

YE LUO AND ARINDAM ROY

ABSTRACT. We address a prime counting problem across the homology classes of a graph, presenting a graph-theoretical Dirichlet-type analogue of the prime number theorem. The main machinery we have developed and employed is a spectral antisymmetry theorem, revealing that the spectra of the twisted graph adjacency matrices have an antisymmetric distribution over the character group of the graph with a special character called the canonical character being an extremum. Additionally, we derive some trace formulas based on the twisted adjacency matrices as part of our analysis.

1. INTRODUCTION

The study of the distribution of prime numbers dates back to Euclid's classical proof demonstrating the infinitude of primes. A more precise understanding of this distribution is captured by the Prime Number Theorem, which states that if $\pi(x)$ denotes the number of primes less than or equal to x, then

(1.1)
$$\pi(x) \sim \frac{x}{\log x}$$

Subsequent research extended this analysis to primes within specific subsets of the positive integers. A foundational result in this direction is Dirichlet's theorem, which asserts that there are infinitely many primes in any arithmetic progression of the form a + qn : n = 0, 1, 2, ..., provided that a and q are coprime positive integers. Furthermore, if $\pi(x;q,a)$ denotes the number of primes up to x that are congruent to $a \mod q$, then the analogue of the Prime Number Theorem in this setting is given by

(1.2)
$$\pi(x;q,a) \sim \frac{1}{\phi(q)} \frac{x}{\log x}$$

where $\phi(q)$ is Euler's totient function. The proofs of both (1.1) and (1.2) rely on explicit formulas that connect the distribution of primes to the zeros of the Riemann zeta function and Dirichlet *L*functions, respectively. In the case of primes in arithmetic progressions, the orthogonality relations of Dirichlet characters also play a crucial role.

The prime number theorem was extended beyond the traditional number system, such as prime ideal theorem in algebraic number theory and prime geodesic theorem in Riemannian geometry. In these generalizations, analogous objects to the Riemann zeta function, like the Dedekind zeta function and the Selberg zeta function, play crucial roles in establishing the corresponding prime number theorems. In this article, we will concentrate on the prime number theorem of graph theory. Our objective is to establish some analogy of (1.2), explicit formula, and orthogonality of characters in graph theory by exploring spectral behaviors over the character group. To define the

⁽A1) SCHOOL OF INFORMATICS, XIAMEN UNIVERSITY, XIAMEN, FUJIAN 361005, CHINA

⁽A2) DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF NORTH CAROLINA AT CHARLOTTE, 9201 UNIVERSITY CITY BLVD., CHARLOTTE, NC 28223, USA

E-mail addresses: (A1) luoye@xmu.edu.cn, (A2) arindam.roy@charlotte.edu.

²⁰²⁰ Mathematics Subject Classification. 05C50, 05C38, 11M41.

Key words and phrases. Homological spectral graph theory, Spectral antisymmetry, Twisted graph adjacency, Canonical character, Complex unit gain graph, Trace formula.

prime number theorem for graph theory and make the paper self-contained, we will first lay the necessary groundwork of graph theory.

Throughout this paper, a graph means a connected finite graph G allowing multiple edges and loops, and we let the vertex set of G be $V(G) = \{v_1, \dots, v_n\}$ and the (unoriented) edge set of G be $E(G) = \{e_1, \dots, e_m\}$. The genus g of G is the first Betti number of G which can be computed by g = m - n + 1. For an orientation O of G, let \mathbf{e}_i and \mathbf{e}_{m+i} be respectively the positively and negatively oriented edges corresponding to e_i with respect to O. Then $\mathbf{E}_O(G) := \{\mathbf{e}_1, \cdots, \mathbf{e}_m\}$ is the set of all positively oriented edges and $\mathbf{E}(G) := {\mathbf{e}_1, \cdots, \mathbf{e}_{2m}}$ is the set of all oriented edges. For each oriented edge $\mathbf{a} \in \mathbf{E}(G)$, denote the *initial* and *terminal* vertices of \mathbf{a} by $\mathbf{a}(0)$ and $\mathbf{a}(1)$ respectively, and the *inverse* \mathbf{a}^{-1} of \mathbf{a} is the oriented edge with $\mathbf{a}^{-1}(0) = \mathbf{a}(1)$ and $\mathbf{a}^{-1}(1) = \mathbf{a}(0)$. A walk Δ on G of length $\tau(\Delta) = N$ is a sequence of oriented edges $\mathbf{a}_1 \cdots \mathbf{a}_N$ such that $\mathbf{a}_{i+1}(0) = \mathbf{a}_i(1)$ for $i = 1, \dots, N-1$, of which the *initial vertex* is $\Delta(0) := \mathbf{a}_1(0)$ and the terminal vertex is $\Delta(1) := \mathbf{a}_N(1)$. If the underlying edges of Δ are all distinct, we also call Δ a path. The inverse of Δ is the walk $\Delta^{-1} := \mathbf{a}_N^{-1} \cdots \mathbf{a}_1^{-1}$. A walk $\Delta' = \mathbf{b}_1 \cdots \mathbf{b}_M$ is a sub-walk of Δ if $\mathbf{b}_j = \mathbf{a}_{i+j}$ for all $j = 1, \dots, M$ and some $i \in \{0, \dots, N-1\}$. We say Δ has a *tail* if $\mathbf{a}_N = \mathbf{a}_1^{-1}$, and has a *backtrack* if $\mathbf{a}_{i+1} = \mathbf{a}_i^{-1}$ for some $1 \leq i \leq N-1$. We say Δ is a *closed walk* with respect to the base vertex v if $v = \mathbf{a}_{N}(1) = \mathbf{a}_{1}(0)$. If in addition Δ has no tail and backtracks, then we call Δ a *circuit*. A *prime circuit* is a circuit C which cannot be written as $C = D^r$ where D is a circuit and r > 1. For any circuit C, there exists a unique prime decomposition $C = P^r$ where P is prime. Two circuits are said to be *equivalent* if the only possible difference is the initial vertex. Denote by [C] the equivalence class of circuits equivalent to C. We call [C] a cycle and [P] a prime cycle if P is a prime circuit. For $C = P^r$ with P prime, denote r([C]) = r(C) := r, which can be considered as the number of automorphisms of C by translation. Denote the set of circuits by \mathcal{C} , the set of prime circuits by \mathcal{P} , the set of cycles by $\overline{\mathcal{C}} = \{[C] \mid C \in \mathcal{C}\}$, and the set of prime cycles by $\overline{\mathcal{P}} = \{ [P] \mid P \in \mathcal{P} \}.$

The *prime number theorem for graphs* evaluates the following counting function (in a fashion of equality instead of inequality for the standard prime number theorem)

(1.3)
$$\pi(l) = \pi_G(l) := \#\{[P] \in \overline{\mathcal{P}} : \tau(P) = l\}$$

where $\tau(P)$ is the length of P. Hashimoto was the first to give an asymptotic formula of $\pi_G(l)$ [Has89, Has92], relying on a zeta function defined by Ihara [Iha66] on p-adic groups. Notably, the Ihara zeta function can be interpreted graph-theoretically, as highlighted by Serre [Ser80]. Overtime, efforts have been made in eliminating the constraint of graph regularity in Ihara's initial formulation and providing satifying determinant expressions of the zeta function, as evidenced by Hashimoto [Has89, Has92] and Bass [Bas92]. For more details readers are referred to Hejhal-Friedman-Gutzwiller-Odlyzko [HFGO99], Kotani-Sonata [KS00], Stark-Terras [ST96, ST00, TS07], Horton-Stark-Terras [HST06], Terras [Ter11], etc.

As an analogue of the Riemann zeta function and the Selberg zeta function, the *Ihara zeta* function of G is defined as

(1.4)
$$z(u) = z_G(u) := \prod_{[P] \in \overline{\mathcal{P}}} \left(1 - u^{\tau(P)}\right)^{-1}.$$

Let R_G be the radius of the largest circle of convergence of $z_G(u)$ and $\nu_G = \gcd\{\tau(P) : [P] \in \overline{\mathcal{P}}\}$. Then an asymptotic formula (see, e.g., [HST06]) of $\pi_G(k)$ is given by

(1.5)
$$\pi_G(l) \sim \frac{\nu_G}{l \cdot R_G^l},$$

which is a graph-theoretic analogue of (1.1). The *edge adjacency matrix* W_1 of G is a $2m \times 2m$ matrix with entries 0 and 1 such that for $\mathbf{a}, \mathbf{b} \in \mathbf{E}(G)$, $(W_1)_{\mathbf{ab}} = 1$ if $\mathbf{a}(1) = \mathbf{b}(0)$ and $\mathbf{b} \neq \mathbf{a}^{-1}$

(meaning that **a** feeds into **b**, denoted by $\mathbf{a} \to \mathbf{b}$), and $(W_1)_{\mathbf{ab}} = 0$ otherwise. To prove (1.5), one can apply the following trace formula:

(1.6)
$$N(l) = \sum_{\lambda \in \operatorname{Spec} W_1} \lambda^l,$$

where $N(l) = \#\{\text{circuits } C \in \mathcal{C} \mid \tau(C) = l\} = \sum_{d|l} d \cdot \pi(d).$

There are a few studies that further explored on the distributions of prime cycles. In [Has92], Hashimoto derived a related result. He considered a finite graph G and a subgroup S of $\operatorname{Aut}(G)$. Then he obtained the distribution of $\pi(S, G; l)$, which is the number of S-equivalence classes of prime paths up to length l. He also obtained an analogue of the Chebotarev's density theorem on the distribution of $\pi(S, G; l, [\sigma])$, which is the number of S-equivalence classes of prime paths up to length l whose Frobenius is a conjugate of σ . In [HS15], Hasegawa and Saito considered an average result of (1.5) and then generalized that result in an arithmetic progression setting. In particular they obtained an asymptotic formula for $\sum_{\substack{l \leq N \\ l \equiv \alpha \mod k}} \pi_G(n)$ for fixed $0 \leq \alpha \leq k$ with $(\alpha, k) = 1$.

They have also obtained a density result. Let S be a subset of \mathcal{P} and define the counting function $\Pi_S(N) := \#\{[P] \in S : \tau(P) \leq N\}$ and the prime zeta-function for S by $P_S(u) := \sum_{[P] \in S} u^{\tau(P)}$. Then their density result can be stated as $\lim_{N\to\infty} \frac{\Pi_S(N)}{\Pi_{\overline{\mathcal{P}}}(N)} = \lim_{u \uparrow R_G} \frac{P_S(u)}{P_{\overline{\mathcal{P}}}(u)}$ provided the limit on the left hand side exits.

In this paper, we investigate the prime distributions of a graph within a distinct framework, akin to a character-based analogy of the prime number theorem in graphs (1.5). Our focus lies on enumerating circuits and primes within homology classes using a machinery we call *homological spectral graph theory*. Specifically, to obtain asymptotic formulas for this prime counting function, we will define the characters for a graph and show that these characters also satisfy an orthogonal relation like Dirichlet characters. Unlike Dirichlet characters, we will show that the character group for a non-bipartite graph contains a non-trivial canonical character which resembles some properties of the trivial character in an opposite manner in sense of the spectra of twisted adjacency matrices. Subsequently, we derive some trace formulas, which serve as another key ingredient in our analysis of the prime counting problem.

1.1. **Related work.** Our results have resemblances on some established work on counting closed and prime geodesics within homology classes in hyperbolic geometry. Let \mathbb{H} be the Poincaré upper half-plane, Γ be a Fuchsian group, i.e. a discrete subgroup of $PSL(2,\mathbb{R}), M = \Gamma \setminus \mathbb{H}$ be the corresponding Riemann surface, and $\pi_M(x)$ be the number of prime geodesics on M of length up to x. If Γ is cocompact, Delsarte [Del42] showed that $\pi_M(x) \sim \operatorname{li}(x)$, and Huber [Hub61] accounted for an error term and demonstrated that $\pi_M(x) = \operatorname{li}(e^x) + \sum_{i=1}^m \operatorname{li}(e^{s_i x}) + \mathcal{O}(x^{-1/2}e^{3x/4})$ where $\operatorname{li}(\cdot)$ is the logarithmic integral function and the s_i 's (with $1/2 < s_i \leq 1$) are the exceptional eigenvalues for the Selberg zeta function [Hej76]. This is known as the prime geodesic theorem. It has similar expressions when Γ is co-finite, as demonstrated by Sarnak [Sar82], Iwaniec [Iwa84], Luo-Sarnak [LS95], Soundararajan-Young[SY13], and others. Notably, Sarnak [Sar82] also presented a Chebotarev density theorem for the co-finite model. Now let M be a compact Riemannian manifold of negative curvature, h be the topological entropy of M, and $\phi: \pi_1(M) \to H_1(M,\mathbb{Z})$ be the projection of the fundamental group to the first integral homology group. Adachi and Sunada [AS87a] delved into the counting problem of $\pi(\alpha, x)$ for $\alpha \in H_1(M, \mathbb{Z})$, which is the number of prime geodesics γ on M of length up to x such that $\phi(\gamma) = \alpha$, and proved that $\lim_{x \to \infty} \frac{\log \pi(\alpha, x)}{x} = h$. This finding was improved shortly thereafter by Philips-Sarnak [PS87], Katsuda-Sunada [KS87], Lalley [Lal89], Pollicott [Pol91] and others. In particular, if M is a hyperbolic surface of genus g, then $\pi(\alpha, x) \sim \frac{(g-1)^g e^x}{x^{g+1}}$ as shown in [PS87, KS87]. As a consequence, prime/closed geodescis are uniformly distributed across the homology classes as the length tends to infinity.

On the other hand, our work is also related to conventional spectral graph theory, which explores the graph properties through the analysis of eigenvalues and eigenvectors of the adjacency matrix, Laplacian matrix, and other matrices derived from the graph structure. For a comprehensive overview of spectral graph theory, readers are referred to classical sources such as [Chu97, CRS10, BH12]. In our work, we will focus on homological spectral graph theory with an aim to apply the spectral analysis of the twisted version of the adjacency matrices to problems of enumerating circuits and cycles in homology classes of the graph. With potential applications (including those related to the cycle counting problem shown in this paper), we also expect that our discoveries will pave a new way for exploration in spectral graph theory.

1.2. Our contributions and paper organization. We summarize our results as follows:

Result (1): Our primary discovery unveils an intriguing antisymmetrical phenomenon in the distribution of the spectra of the twisted vertex/edge adjacency matrices over the character group of a finite graph, as stated in Theorem 3.1. It is noteworthy that there is no requirement to impose the constraint of graph regularity. Specifically, consider a finite graph G of genus g. Its character group $\mathcal{X}(G)$ is defined as the Pontryagin dual of the integral homology group $H_1(G,\mathbb{Z})$. Geometrically, $\mathcal{X}(G)$ is a q-dimensional real torus. We showed that there exists a unique character $\theta \in \mathcal{X}(G)$ known as the canonical character of G such that for any character $\chi \in \mathcal{X}(G)$, the spectra (the multiset of eigenvalues) of the twisted vertex/edge adjacency matrix with respect to χ and with respect to $\theta - \chi$ are negations of each other. Moreover, with the exception of certain special cases, the maximum spectral radii of all twisted vertex/edge adjacency matrices are only achieved at the trivial character (non-twisted) and the canonical character. In this context, the trivial character and the canonical character can be viewed as two extrema on the character group, governing the spectral behavior over the character group in an *antisymmetric* manner. As an illustrative example, we show in Figure 2 the distributions of spectral radii over the character group for specific graphs. A key innovation lies in the proof of the "strict" extremity of the trivial and canonical characters (Lemma 4.10 through Lemma 4.17) which utilizes Gelfand's formula along with combinatorial and matrix analysis techniques. Although our investigation on the spectra of the twisted vertex/edge adjacency matrices was originally motivated by the prime counting problem on graphs, we note that a graph endowed with a specific twisted adjacency matrix is equivalent to a complex unit gain graph [Zas89, Ref12], a subject that has drawn significant interest in recent years. In other words, the twisted adjacency matrices of G are essentially the adjacency matrices of the unit gain graphs with identical underlying graph G. Indeed, we provide a top view from a more geometric aspect by examining complex unit gain graphs as a unified entity, as it can be shown that the character group of a graph G is exactly the space of switching equivalence classes of complex unit gain graphs with the same underlying graph G.

Result (2): We present some trace formulas with respect to twisted edge adjacency matrices in Theorem 3.11. The key observation is that the trace of the *n*-th power of the edge adjacency matrix has a Fourier expansion over $H_1(G, \mathbb{Z})$, where the coefficients correspond to the number of circuits in each homology class (see Theorem 3.12(a)(i)). As an example, some identities based on the trace formulas are presented in Subsection 3.5.

Result (3): Leveraging the spectral antisymmetry theorem and trace formulas, we delved into the counting problem of cycles and prime cycles in homology classes of the graph, providing some asymptotic formulas as outlined in Theorem 3.17. It is noteworthy that, unlike the uniform cycle distribution over the homology classes in hyperbolic geometry, there are instances where cycles of certain lengths within specific homology classes of the graph vanish (refer also to Corollary 3.13), essentially a consequence of spectral antisymmetry. An example of cycle counting is also presented in Subsection 3.5. We organize the paper as follows. In Section 2, we provide an introduction to some primitives such as the notions of orthogonal decomposition of the 1-forms, the character group, twisted adjacency matrices and L-functions within a the graph-theoretical framework. Section 3 presents our main results, including the spectral antisymmetry theorem, trace formulas related to the twisted edge adjacency of graphs, and asymptotic prime distribution formulas within homology classes. Illustrative examples are also provided in dedicated subsections. The proof of the spectral antisymmetry theorem, which constitutes the most intricate aspect of this paper, is detailed in Section 4.

2. Preliminaries: Orthogonal decomposition, characters, twisted adjacency matrices and L-functions

2.1. Orthogonal decomposition. Let $\mathcal{T}(G)$ be the real linear space on $\mathbf{E}(G)$ such that $1 \cdot \mathbf{e}^{-1} = -1 \cdot \mathbf{e}$ for any oriented edge \mathbf{e} . Alternatively, by fixing an orientation on the edges, we may also consider $\mathcal{T}(G)$ as a real linear space on $\mathbf{E}_O(G) = \{\mathbf{e}_1, \cdots, \mathbf{e}_m\}$ where $\mathbf{e}_1, \cdots, \mathbf{e}_m$ are the positively oriented edges. Geometrically, one may consider an element of $\mathcal{T}(G)$ as a "vector field" on G. Then a "1-form" ω is an element of the linear dual $\Omega(G)$ of $\mathcal{T}(G)$. We call $\mathcal{T}(G)$ the tangent space of G, and $\Omega(G)$ the cotangent space of G, sometimes also written as \mathcal{T} and Ω respectively for simplicity when G is provided. For $\alpha \in \mathcal{T}$ and $\omega \in \Omega$, we use $\omega(\alpha)$ to represent the value of the pairing between ω and α . For $i = 1, \cdots, m$, let $d\mathbf{e}_i$ be a 1-form such that $d\mathbf{e}_i(\mathbf{e}_j) = \delta_{ij}$ for $j = 1, \cdots, m$. Then $\{d\mathbf{e}_1, \cdots, d\mathbf{e}_m\}$ is a basis of Ω . In addition, the standard inner product on \mathcal{T} with respect to the basis $\{\mathbf{e}_1, \cdots, \mathbf{e}_m\}$ and the standard inner product on Ω with respect to $\{d\mathbf{e}_1, \cdots, d\mathbf{e}_m\}$ are defined by $\langle \mathbf{e}_i, \mathbf{e}_j \rangle = \langle d\mathbf{e}_i, d\mathbf{e}_j \rangle := d\mathbf{e}_i(\mathbf{e}_j) = \delta_{ij}$ for $i, j = 1, \cdots, m$.

For a walk $\Delta = \mathbf{a}_1 \cdots \mathbf{a}_N$, let the *abelianization* of Δ be $\Delta^{ab} = \mathbf{a}_1 + \cdots + \mathbf{a}_N \in \mathcal{T}$, and the integral of a 1-form ω along Δ is $\int_{\Delta} \omega := \omega(\Delta^{ab}) = \omega(\mathbf{a}_1) + \cdots + \omega(\mathbf{a}_N)$.

Let R be a commutative ring with multiplicative identity. Let $C_0(G, R)$ be the free R-module on V(G) and $C_1(G, R)$ be the free R-module on $\mathbf{E}_O(G)$. Let $C^0(G, R) = \operatorname{Hom}_R(C_0(G, R), R)$ and $C^1(G, R) = \operatorname{Hom}_R(C_1(G, R), R)$. As in convention, an element of $C_0(G, R)$ (resp. an element of $C_1(G, R)$) is called a 0-chain (resp. 1-chain) with coefficients in R, and an element of $C^0(G, R)$ (resp. an element of $C^1(G, R)$) is called a 0-cochain (resp. 1-cochain) with coefficients in R. Note that $C_1(G, \mathbb{Z})$ is a full-rank lattice in $C_1(G, \mathbb{R}) = \mathcal{T}(G)$, and $C^1(G, \mathbb{Z})$ is a full-rank lattice in $C^1(G, \mathbb{R}) = \Omega(G)$.

There is a canonical isomorphism $i_R^0: C_0(G, R) \xrightarrow{\sim} C^0(G, R)$ sending a 0-chain $\sum_{v \in V(G)} c_v \cdot v \in C_0(G, R)$ to a 0-cochain $f \in C^0(G, R)$ given by $f(v) = c_v$, and a canonical isomorphism $i_R^1: C_1(G, R) \xrightarrow{\sim} C^1(G, R)$ sending a 1-chain $\sum_{\mathbf{e} \in \mathbf{E}_O(G)} c_{\mathbf{e}} \cdot \mathbf{e} \in C_1(G, R)$ to a 1-cochain $\omega \in C^1(G, R)$ given by $\omega(\mathbf{e}) = c_{\mathbf{e}}$. Note that $i_{\mathbb{R}}^1(\mathbf{e}_i) = d\mathbf{e}_i$ for $i = 1, \cdots, m$.

Let $\partial_R : C_1(G, R) \to C_0(G, R)$ be the boundary map defined by

$$\partial_R \left(\sum_{\mathbf{e} \in \mathbf{E}_O(G)} c_{\mathbf{e}} \cdot \mathbf{e} \right) = \sum_{\mathbf{e} \in \mathbf{E}_O(G)} \left(c_{\mathbf{e}} \cdot \mathbf{e}(1) - c_{\mathbf{e}} \cdot \mathbf{e}(0) \right),$$

and $d_R: C^0(G, R) \to C^1(G, R)$ be the differential map defined by

$$d_R f = \sum_{\mathbf{e} \in \mathbf{E}_O(G)} \left(f(\mathbf{e}(1)) - f(\mathbf{e}(0)) \right) \cdot d\mathbf{e}$$

for each $f \in C^0(G, R)$. Then the adjoint operator $\partial_R^* : C_1(G, R) \to C_0(G, R)$ of ∂ is defined by $\partial_R^* = (i_R^1)^{-1} \circ d \circ i_R^0$, and the adjoint operator $d_R^* : C^1(G, R) \to C^0(G, R)$ is defined by $d_R^* = i_R^0 \circ \partial \circ (i_R^1)^{-1}$. For simplicity of notation, we also write $d_{\mathbb{R}}$ and $d_{\mathbb{R}}^*$ as d and d^* respectively. We say d^*d is the Laplacian on functions and dd^* is the Laplacian on 1-forms, both denoted by Δ .

In the following, we will introduce the notion of harmonic 1-form and present a proposition of "Hodge" orthogonal decomposition which states that any 1-form can be uniquely decomposed into a harmonic 1-form and an exact 1-form. More detailed discussions can be found in [BF11].

Definition 2.1. We say $\omega \in \Omega(G)$ is a harmonic 1-form if $\Delta \omega = 0$. For each $f \in C^0(G, \mathbb{R})$, we say df is an exact 1-form.

Lemma 2.2. $\omega = \sum_{\mathbf{e} \in \mathbf{E}_O(G)} \omega_{\mathbf{e}} \cdot d\mathbf{e}$ is a harmonic 1-form if and only if $d^*\omega = 0$ if and only if $\sum_{\mathbf{e} \in \mathbf{E}_O(G), \mathbf{e}(1) = v} \omega_{\mathbf{e}} = \sum_{\mathbf{e} \in \mathbf{E}_O(G), \mathbf{e}(0) = v} \omega_{\mathbf{e}}$ for all $v \in V(G)$.

Proof. Note that the kernel of d is made of constant functions and $d^*\omega$ can be a constant function only if $d^*\omega = 0$.

Denote the space of all harmonic 1-forms by $\mathcal{H}^1(G)$ and the space of all exact 1-forms by $\operatorname{Im}(d)$. Clearly $\mathcal{H}^1(G)$ and $\operatorname{Im}(d)$ are both linear subspaces of $\Omega(G)$. The following proposition says that they are orthogonal complements of each other.

Proposition 2.3 (Hodge orthogonal decomposition). There is a canonical Hodge decomposition $\Omega(G) = \mathcal{H}^1(G) \oplus \operatorname{Im}(d)$. That is, any 1-form ω can be written uniquely as $\omega = \phi_1(\omega) + \phi_2(\omega)$ where $\phi_1(\omega)$ is a harmonic 1-form and $\phi_2(\omega)$ is an exact 1-form. In addition, $\langle \phi_1(\omega), \phi_2(\omega) \rangle = 0$.

Proof. Consider the function $d^*\omega$. It is clear that $\sum_{v \in V(G)} d^*\omega(v) = 0$. Therefore, the equation $\Delta f = d^*df = d^*\omega$ has a unique solution of f up to addition by a constant, since the Laplacian matrix Δ has rank n-1 (the sum of each column of Δ is 0, and G is connected) whose null space is consisted of the constant functions. Then df is an exact form and $\omega - df$ is a harmonic 1-form by Lemma 2.2. Let $df = \sum_{\mathbf{e} \in \mathbf{E}_O(G)} \omega'_{\mathbf{e}} \cdot d\mathbf{e}$ and $\omega - df = \sum_{\mathbf{e} \in \mathbf{E}_O(G)} \omega'_{\mathbf{e}} \cdot d\mathbf{e}$. It remains to show that $\langle df, \omega - df \rangle = \sum_{\mathbf{e} \in \mathbf{E}_O(G)} \omega'_{\mathbf{e}} \omega''_{\mathbf{e}} = 0$. Note that $\omega'_{\mathbf{e}} = f(\mathbf{e}(1)) - f(\mathbf{e}(0))$. Then

$$\sum_{\mathbf{e}\in\mathbf{E}_{O}(G)}\omega_{\mathbf{e}}'\omega_{\mathbf{e}}'' = \sum_{\mathbf{e}\in\mathbf{E}_{O}(G)}\left(f(\mathbf{e}(1)) - f(\mathbf{e}(0))\right)\omega_{\mathbf{e}}''$$
$$= \sum_{v\in V(G)}f(v)\left(\sum_{\mathbf{e}\in\mathbf{E}_{O}(G),\mathbf{e}(1)=v}\omega_{\mathbf{e}}'' - \sum_{\mathbf{e}\in\mathbf{E}_{O}(G),\mathbf{e}(0)=v}\omega_{\mathbf{e}}''\right) = 0$$

Remark 2.4. Since the first real cohomology group is computed as $H^1(G, \mathbb{R}) = \Omega(G) / \operatorname{Im}(d)$, we have $H^1(G, \mathbb{R}) \simeq \mathcal{H}^1(G)$. This means that any 1-cocycle in $H^1(G, \mathbb{R})$ can be represented canonically by a unique harmonic 1-form.

The first real and integral homology groups are $H_1(G, \mathbb{R}) = \ker \partial_{\mathbb{R}}$ and $H_1(G, \mathbb{Z}) = \ker \partial_{\mathbb{Z}}$ respectively. In particular, $H_1(G, \mathbb{Z}) = H_1(G, \mathbb{R}) \bigcap C_1(G, \mathbb{Z})$. Note that integral 1-cycles (elements of $H_1(G, \mathbb{Z})$) are also called *circulations* in graph theory.

Lemma 2.5. We have the following properties of $H_1(G, R)$:

- (a) $H_1(G, R)$ is isomorphic to R^g ;
- (b) $H_1(G,\mathbb{Z})$ is a full-rank lattice in $H_1(G,\mathbb{R})$;
- (c) $H_1(G, \mathbb{Z}/t\mathbb{Z})$ is canonically isomorphic to $H_1(G, \mathbb{Z})/tH_1(G, \mathbb{Z})$.

Proof. Let T be a spanning tree of G. Then $G \setminus T$ contains g edges. Fix an orientation of G and without loss of generality, we may label the positively oriented edges in $G \setminus T$ to be $\mathbf{e}_1, \cdots, \mathbf{e}_g$ and the positively oriented edges in T to be $\mathbf{e}_{g+1}, \cdots, \mathbf{e}_m$. Let K be the free R-module on $\{\mathbf{e}_1, \cdots, \mathbf{e}_g\}$. Clearly $K \simeq R^g$ and we claim that $H_1(G, R) \simeq K$. Consider the restriction homomorphism $\rho: H_1(G, R) \to K$ sending $\sum_{i=1}^m c_i \mathbf{e}_i \in H_1(G, R)$ to $\sum_{i=1}^g c_i \mathbf{e}_i \in K$. For each \mathbf{e}_i , consider its initial vertex $\mathbf{e}_i(0)$ and terminal vertex $\mathbf{e}_i(1)$. Then there is a unique path P_i from $\mathbf{e}_i(1)$ to $\mathbf{e}_i(0)$ contained in the spanning tree *T*. Let $u_i = \mathbf{e}_i + P_i^{ab}$ for $i = 1, \dots, g$. It is clear u_i 's are elements of $H_1(G, \mathbb{Z})$, and $\rho(\sum_{i=1}^g c_i u_i) = \sum_{i=1}^g c_i \mathbf{e}_i$. This implies that ρ is surjective. Moreover, it can be easily shown that $\sum_{i=g+1}^m c_i \mathbf{e}_i = 0$ if and only if $c_i = 0$ for all $i = g + 1, \dots, m$. The injectivity of ρ follows.

(b) is straightforward to verify.

For (c), consider the canonical homomorphism $\phi : H_1(G, \mathbb{Z}) \to H_1(G, \mathbb{Z}/t\mathbb{Z})$ which sends $\sum_{i=1}^m c_i \mathbf{e}_i$ to $\sum_{i=1}^m \overline{c}_i \mathbf{e}_i$ where $\overline{c}_i = c_i \mod t$. Then it is easy to see that ker $\phi = tH_1(G, \mathbb{Z})$ and it remains to show that ϕ is surjective. As in the proof of (1), we let T be a spanning tree where the oriented edges of $G \setminus T$ are $\mathbf{e}_1, \cdots \mathbf{e}_g$. By the argument in the proof of (a), we see that there is an isomorphism between $H_1(G, \mathbb{R})$ and the free \mathbb{R} -module on $\{\mathbf{e}_1, \cdots \mathbf{e}_g\}$. This means we can lift each $\sum_{i=1}^m \overline{c}_i \mathbf{e}_i \in H_1(G, \mathbb{Z}/t\mathbb{Z})$ to some $\sum_{i=1}^m c_i \mathbf{e}_i \in H_1(G, \mathbb{Z})$ by first lifting $\overline{c}_1, \cdots \overline{c}_g \in \mathbb{Z}/t\mathbb{Z}$ to $c_1, \cdots c_g \in \mathbb{Z}$ freely.

Remark 2.6. Since $H_1(G, \mathbb{Z})$ is a full-rank lattice in $H_1(G, \mathbb{R})$ (or a discrete cocompact subgroup of $H_1(G, \mathbb{R})$) and the pairing $\mathcal{H}^1(G) \times H_1(G, \mathbb{R}) \to \mathbb{R}$ is perfect, the dual lattice $H_1(G, \mathbb{Z})^{\vee} =$ $\operatorname{Hom}(H_1(G, \mathbb{Z}), \mathbb{Z})$ of $H_1(G, \mathbb{Z})$ is a full-rank lattice in $\mathcal{H}^1(G)$. It can also be easily verified that $i_{\mathbb{Z}}^2(H_1(G, \mathbb{Z}))$ is a subgroup of $H_1(G, \mathbb{Z})^{\vee}$ of finite index.

As shown in the proof of Lemma 2.5, the u_i 's form a basis of $H_1(G, \mathbb{Z})$. The following proposition shows how to derive a dual basis of $H_1(G, \mathbb{Z})^{\vee}$.

Proposition 2.7. For a graph G of genus g and T a spanning tree of G, let $\mathbf{e}_1, \cdots, \mathbf{e}_g \in \mathbf{E}_O(G)$ be the positively oriented edges in $G \setminus T$ with respect to some orientation O on G. Then $\{\phi_1(d\mathbf{e}_1), \cdots, \phi_1(d\mathbf{e}_g)\}$ is a basis of $H_1(G, \mathbb{Z})^{\vee}$ in $\mathcal{H}^1(G)$. In addition, $H_1(G, \mathbb{Z})^{\vee} = \phi_1(C^1(G, \mathbb{Z}))$.

Proof. Using the orthogonal decomposition in Proposition 2.3, we have $d\mathbf{e}_i = \phi_1(d\mathbf{e}_i) + \phi_2(d\mathbf{e}_i)$ where $\phi_1(d\mathbf{e}_i) \in \mathcal{H}^1(G)$ is a harmonic 1-form and $\phi_2(d\mathbf{e}_i) \in \text{Im}(d)$ is an exact form.

For a spanning tree T of G, let $u_i = \mathbf{e}_i + P_i^{ab}$ for $i = 1, \dots, g$ as in the proof of Lemma 2.5. Then $\{u_1, \dots, u_g\}$ is a basis of $H_1(G, \mathbb{Z})$. Since e_i is always disjoint from P_j , we must have $d\mathbf{e}_i(\mathbf{u}_j) = d\mathbf{e}_i(\mathbf{e}_j) + d\mathbf{e}_i(P_i^{ab}) = d\mathbf{e}_i(\mathbf{e}_j) = \delta_{ij}$ for all $i, j = 1, \dots, g$.

Note that $d\mathbf{e}_i(u) = \phi_1(d\mathbf{e}_i)(u)$ for all $u \in H_1(G, \mathbb{Z})$. This implies $\phi_1(d\mathbf{e}_i)(u_j) = \delta_{ij}$, i.e., $\{\phi_1(d\mathbf{e}_1), \dots, \phi_1(d\mathbf{e}_g)\}$ is a basis of $H_1(G, \mathbb{Z})^{\vee}$ as desired, which is the dual to the basis $\{u_1, \dots, u_g\}$ of $H_1(G, \mathbb{Z})$.

It remains to show that $H_1(G,\mathbb{Z})^{\vee} = \phi_1(C^1(G,\mathbb{Z}))$. We just need to show that $\phi_1(d\mathbf{e}_i) \in H_1(G,\mathbb{Z})^{\vee}$ for $i = 1, \dots, m$. By the above discussion, this follows from the fact that $\phi_1(d\mathbf{e}_i) = 0$ (meaning that $d\mathbf{e}_i$ is an exact 1-form), if and only if e_i is a bridge (i.e., $G \setminus e_i$ is disjoint), if and only if e_i is contained in all spanning trees of G.

2.2. The character group. A (unitary) character $\chi_{\omega} : \mathcal{T} \to S^1$ of \mathcal{T} can be associated to each 1-form $\omega \in \Omega$ by letting $\chi_{\omega}(\alpha) = e(\omega(\alpha))$ for all $\alpha \in \mathcal{T}$. Here $e(x) := \exp(2\pi\sqrt{-1}x)$ and we will use this conventional notation throughout the paper. Note that $\chi_{-\omega} = \overline{\chi_{\omega}}$ where $\overline{\chi_{\omega}}$ is the conjugate character of χ_{ω} .

For a locally compact group Γ , denote by $\widehat{\Gamma}$ the *Pontryagin dual* of Γ , which is the group of unitary characters (continuous group homomorphisms from Γ to the unit circle S^1) on Γ .

Lemma 2.8. We have the following commutative diagram where the bottom row is exact and $\mathcal{X}(G) = \mathcal{H}^1(G)/H_1(G,\mathbb{Z})^{\vee}$:



Proof. We will concretize the well-known results about Pontryagin duals in our setting: the Pontryagin dual of a finite dimensional real vector space is a real vector space of the same dimension, and the Pontryagin dual of a rank-*q* lattice is a *q*-dimensional real torus.

First note that since the unit circle $S^1 \simeq \mathbb{R}/\mathbb{Z}$ is an injective \mathbb{Z} -module, the sequence $H_1(G,\mathbb{Z}) \hookrightarrow$ $H_1(G,\mathbb{R}) \hookrightarrow \mathcal{T}(G)$ induces the sequence on their Pontryagin duals $\mathcal{T}(G) \twoheadrightarrow H_1(G,\mathbb{R}) \twoheadrightarrow H_1(G,\mathbb{Z})$.

Now we will show that the homomorphism from $\Omega(G)$ to $\mathcal{T}(G)$ defined by $\omega \mapsto \chi_{\omega}$ is actually an isomorphism. Let $\omega = \sum_{i=1}^{m} \omega_i \cdot d\mathbf{e}_i$ be a nontrivial 1-form and we may assume ω_1 is nonzero. Choose a real number c such that $c\omega_1$ is not an integer. Then $\chi_{\omega}(c\mathbf{e}_1) = e(\omega(c\mathbf{e}_1)) = e(c\omega_1) \neq 1$ which means χ_{ω} cannot be the identity in $\widehat{\mathcal{T}(G)}$. It remains to show that for any character $\chi \in \widehat{\mathcal{T}(G)}$. There exists a 1-form such that $\chi = \chi_{\omega}$. Consider the function $f(c_1, \dots, c_m) = \chi(\sum_{i=1}^m c_i \mathbf{e}_i)$. Then χ being a character of \mathcal{T} guarantees that $\frac{\partial f}{\partial c_i} = 2\pi\sqrt{-1}\omega_i f$ where ω_i 's are constant real numbers. Let $\omega = \sum_{i=1}^m \omega_i \cdot d\mathbf{e}_i$ and it follows clearly that $\chi = \chi_{\omega}$. By Proposition 2.3, $\omega = \phi_1(\omega) + \phi_2(\omega)$ where $\phi_1(\omega) \in \mathcal{H}^1(G)$ and $\phi_2(\omega) \in \text{Im}(d)$. Then

 $\chi_{\omega}|_{H_1(G,\mathbb{R})} = \chi_{\phi_1(\omega)}|_{H_1(G,\mathbb{R})}$, which induces a compatible isomorphism between $\mathcal{H}^1(G)$ and $\widehat{H_1(G,\mathbb{R})}$.

It remains to show that there is a compatible isomorphism between $\mathcal{X}(G)$ and $H_1(G,\mathbb{Z})$. For $\omega, \omega' \in \Omega(G), \ \chi_{\omega}|_{H_1(G,\mathbb{Z})} = \chi_{\omega'}|_{H_1(G,\mathbb{Z})}, \text{ if and only if } e(\omega(\alpha)) = e(\omega'(\alpha)) \text{ for all } \alpha \in H_1(G,\mathbb{Z}), \text{ if } \alpha \in H_1(G,\mathbb{Z}), \text{ or } \alpha \in H$ and only if $\omega(\alpha) - \omega'(\alpha) \in \mathbb{Z}$ for all $\alpha \in H_1(G, \mathbb{Z})$, if and only if $\phi_1(\omega) - \phi_1(\omega') \in H_1(G, \mathbb{Z})^{\vee}$. Thus, we have $H_1(G,\mathbb{Z}) \simeq \mathcal{H}^1(G)/H_1(G,\mathbb{Z})^{\vee}$.

Remark 2.9. For the rest of the paper, we will identify $\widehat{\mathcal{T}(G)}$, $\widehat{H_1(G,\mathbb{R})}$ and $\widehat{H_1(G,\mathbb{Z})}$ with $\Omega(G)$, $\mathcal{H}^1(G)$ and $\mathcal{X}(G)$ respectively. In particular, we call $\mathcal{X}(G)$ the character group of G, which is a g-dimensional real torus. Provided that G is known, we also simply write $\mathcal{X}(G)$ as \mathcal{X} . For $\omega \in \Omega(G)$, we write $\underline{\omega} := \phi_1(\omega) + H_1(G, \mathbb{Z})^{\vee} \in \mathcal{X}$. By Lemma 2.8, when restricted to $H_1(G, \mathbb{Z})$, we also write χ_{ω} as $\chi_{\underline{\omega}}$ for all $\omega \in \Omega(G)$. In comparison to the character group, it is worth mentioning some other related groups. Note that both being lattices in $H_1(G,\mathbb{R}), H_1(G,\mathbb{Z})$ is subgroup of $(i_{\mathbb{Z}}^1)^{-1}(H_1(G,\mathbb{Z})^{\vee})$ of finite index (Remark 2.6). Then $(i_{\mathbb{Z}}^1)^{-1}(H_1(G,\mathbb{Z})^{\vee})/H_1(G,\mathbb{Z})$ is the Jacobian group $\operatorname{Jac}(G)$ of the graph G [BN07], and the g-dimensional real torus $H_1(G,\mathbb{R})/H_1(G,\mathbb{Z})$ is the (tropical) Jacobian group $Jac(\Gamma)$ of the metric graph Γ obtained from G by identifying edges with the unit interval [BF11].

For elements x and y in some set X, let δ_{xy} be the delta function, i.e., $\delta_{xy} = 1$ if x = y, and $\delta_{xy} = 0$ otherwise.

Proposition 2.10. Fixing $\alpha, \beta \in H_1(G, \mathbb{Z})$, we have the following orthogonal relation

$$\frac{1}{\operatorname{vol}(\mathcal{X})} \int_{\mathcal{X}} \chi_{\omega}(\alpha) \overline{\chi_{\omega}(\beta)} dV_{\omega} = \delta_{\alpha\beta}$$

where $\chi_{\omega}(\alpha)\chi_{\omega}(\beta)$ is considered as a function on \mathcal{X} with variable ω , dV_{ω} is the volume g-form on \mathcal{X} and $\operatorname{vol}(\mathcal{X})$ is the total volume of \mathcal{X} .

Proof. We identify the domain of integration \mathcal{X} with the fundamental domain of the lattice $H_1(G, \mathbb{Z})^{\vee}$ in $\mathcal{H}^1(G)$.

Let $\{u_1, \dots, u_g\}$ be a basis of $H_1(G, \mathbb{Z})$ and $\{u^1, \dots, u^g\}$ be the corresponding dual basis of $H_1(G, \mathbb{Z})^{\vee}$, i.e., $u^i(u_j) = \delta_{ij}$. Then we can write $\alpha = \sum_{i=1}^g \alpha_i u_i$, $\beta = \sum_{i=1}^g \beta_i u_i$ and $\omega = \sum_{i=1}^g \omega_i u^i$. Now

$$\chi_{\omega}(\alpha)\overline{\chi_{\omega}(\beta)} = e\left(\sum_{i=1}^{g}\omega_i(\alpha_i - \beta_i)\right)$$

and

$$dV_{\omega} = \sqrt{|\det((\langle u^i, u^j \rangle)_{i,j})|} d\omega_1 \cdots d\omega_g.$$

Since the integration is over the fundamental domain of the lattice $H_1(G,\mathbb{Z})^{\vee}$, we have

$$\int_{\mathcal{X}} \chi_{\omega}(\alpha) \overline{\chi_{\omega}(\beta)} dV_{\omega}$$

$$= \int_{\omega_1=0}^{1} \cdots \int_{\omega_g=0}^{1} e\left(\sum_{i=1}^{g} \omega_i(\alpha_i - \beta_i)\right) \sqrt{|\det((\langle u^i, u^j \rangle)_{i,j})|} d\omega_1 \cdots d\omega_g$$

$$= \operatorname{vol}(\mathcal{X}) \delta_{\alpha\beta}.$$

Now let us consider a subgroup Λ of $H_1(G, \mathbb{Z})$ of finite index. Then Λ is also a rank-g lattice in $H_1(G, \mathbb{R})$, and the corresponding dual lattice Λ^{\vee} in $\mathcal{H}^1(G)$ contains $H_1(G, \mathbb{Z})^{\vee}$ as a sub-lattice. Let $Q_{\Lambda} := H_1(G, \mathbb{Z})/\Lambda$ and $\mathcal{X}_{\Lambda} := \mathcal{H}^1(G)/\Lambda^{\vee}$. For $\alpha \in H_1(G, \mathbb{Z})$ and $\omega \in \Omega(G)$, we write $\underline{\alpha} = \alpha + \Lambda \in Q_{\Lambda}$ and $\underline{\omega} := \phi_1(\omega) + \Lambda^{\vee} \in \mathcal{X}_{\Lambda}$.

Lemma 2.11. There exists a commutative diagram with the rows being exact:

Proof. Since the unit circle is $S^1 \simeq \mathbb{R}/\mathbb{Z}$ is an injective \mathbb{Z} -module, the exact sequence $0 \to \Lambda \hookrightarrow H_1(G,\mathbb{Z}) \twoheadrightarrow Q_\Lambda \to 0$ induces an exact sequence on their Pontryagin duals $0 \to \widehat{Q_\Lambda} \hookrightarrow \widehat{H_1(G,\mathbb{Z})} \twoheadrightarrow \widehat{\Lambda} \to 0$. Since $\mathcal{X} = \mathcal{H}^1(G)/H_1(G,\mathbb{Z})^{\vee}$ and $\mathcal{X}_{\Lambda} := \mathcal{H}^1(G)/\Lambda^{\vee}$, we have the exact sequence $0 \to \Lambda^{\vee}/H_1(G,\mathbb{Z})^{\vee} \hookrightarrow \mathcal{X} \twoheadrightarrow \mathcal{X}_{\Lambda} \to 0$.

The isomorphism $\mathcal{X} \simeq H_1(G, \mathbb{Z})$ defined by $\underline{\omega} \mapsto \chi_{\underline{\omega}}$ is the same as in Lemma 2.8, which naturally induces compatible isomorphisms $\Lambda^{\vee}/H_1(G, \mathbb{Z})^{\vee} \simeq \widehat{Q_{\Lambda}}$ and $\mathcal{X}_{\Lambda} \simeq \widehat{\Lambda}$. In particular, for each $\omega \in \Lambda^{\vee}$, $\chi_{\underline{\omega}}$ is a character of $H_1(G, \mathbb{Z})$ such that for each $\alpha, \alpha' \in H_1(G, \mathbb{Z})$ such that $\alpha - \alpha' \in \Lambda$, we have $\chi_{\underline{\omega}}(\alpha) = \chi_{\underline{\omega}}(\alpha')$. This means that $\chi_{\underline{\omega}}$ can be considered as a character of Q_{Λ} . Moreover, for $\omega, \omega' \in \Omega(G), \chi_{\omega}|_{\Lambda} = \chi_{\omega'}|_{\Lambda}$ if and only if $\phi_1(\omega) - \phi_1(\omega') \in \Lambda^{\vee}$.

Remark 2.12. As in Remark 2.9, for the rest of the paper, we will identify \widehat{Q}_{Λ} and $\widehat{\Lambda}$ with $\Lambda^{\vee}/H_1(G,\mathbb{Z})^{\vee}$ and \mathcal{X}_{Λ} respectively. Note that \widehat{Q}_{Λ} is finite abelian group non-canonically isomorphic to Q_{Λ} . In addition, by Lemma 2.11, when restricted to Λ , we also write χ_{ω} as $\chi_{\underline{\omega}}$ for all $\omega \in \Omega(G)$.

The following proposition is a generalization of Proposition 2.10.

Proposition 2.13. (a) Fixing $\alpha, \beta \in \Lambda$, we have the following orthogonal relation

$$\frac{1}{\operatorname{vol}(\mathcal{X}_{\Lambda})} \int_{\mathcal{X}_{\Lambda}} \chi_{\omega}(\alpha) \overline{\chi_{\omega}(\beta)} dV_{\omega} = \delta_{\alpha\beta}$$

where $\chi_{\omega}(\alpha)\chi_{\omega}(\overline{\beta})$ is considered as a function on \mathcal{X}_{Λ} with variable ω , dV_{ω} is the volume g-form on \mathcal{X}_{Λ} and $\operatorname{vol}(\mathcal{X}_{\Lambda})$ is the total volume of \mathcal{X}_{Λ} .

(b) Fixing $\underline{\alpha}, \beta \in Q_{\Lambda}$, we have the following orthogonal relation

$$\frac{1}{|Q_{\Lambda}|} \sum_{\underline{\omega} \in \widehat{Q_{\Lambda}}} \chi_{\underline{\omega}}(\alpha) \overline{\chi_{\underline{\omega}}(\beta)} = \delta_{\underline{\alpha}\underline{\beta}}$$

Proof. (a) follows from an analogous argument as in the proof of Proposition 2.10.

Since $\sum_{\underline{\omega}\in\widehat{Q_{\Lambda}}}\chi_{\underline{\omega}}(\alpha)\overline{\chi_{\underline{\omega}}(\beta)} = \sum_{\underline{\omega}\in\widehat{Q_{\Lambda}}}\chi_{\underline{\omega}}(\alpha-\beta)$, (b) follows from the classical fact of characters on finite abelian groups.

Remark 2.14. By Lemma 2.11, one can see that $|Q_{\Lambda}| \cdot \operatorname{vol}(\mathcal{X}_{\Lambda}) = \operatorname{vol}(\mathcal{X})$. Actually, $\operatorname{vol}(\mathcal{X})$ has an interpretation with respect to the total number of spanning trees of G. Note that $\mathcal{X} = \mathcal{H}^1(G)/\mathcal{H}_1(G,\mathbb{Z})^{\vee} = H_1(G,\mathbb{R})/\mathcal{H}_1(G,\mathbb{Z})^{\vee}$. As mentioned in Remark 2.9, the Jacobian Jac(G) of the graph G is $(i_{\mathbb{Z}}^1)^{-1}(\mathcal{H}_1(G,\mathbb{Z})^{\vee})/\mathcal{H}_1(G,\mathbb{Z})$, and the Jabobian Jac(Γ) of the metric graph Γ corresponding to G is $H_1(G,\mathbb{R})/\mathcal{H}_1(G,\mathbb{Z})$. It follows that $\operatorname{vol}(\mathcal{X})\cdot\operatorname{vol}(\operatorname{Jac}(\Gamma)) = 1$ and $\operatorname{vol}(\operatorname{Jac}(\Gamma)) = \operatorname{vol}(\mathcal{X}) \cdot |\operatorname{Jac}(G)|$. It is well known that Jac(G), also called the critical group of G, is bijective to the set of spanning trees of G. A geometric interpretation of such a bijection is provided in the dual version of Kirchhoff's matrix-tree theorem first discovered by An, Baker, KuperBerg and Shokrieh (see Theorem 1.5 and Corollary 1.6 of [ABKS14]). Summarizing the above arguments, we give the following proposition.

Proposition 2.15. Let w(G) be complexity of G, i.e., the number of spanning trees of G. Then $vol(\mathcal{X}) = 1/\sqrt{w(G)}$ and $vol(\mathcal{X}_{\Lambda}) = 1/(\sqrt{w(G)} \cdot |Q_{\Lambda}|)$.

2.3. Twisted adjacency matrices and twisted edge adjacency matrices. For $\omega \in \Omega(G)$, the adjacency and edge adjacency matrices can be twisted by the corresponding character χ_{ω} .

- **Definition 2.16.** (i) The *(vertex)* adjacency matrix of G twisted by χ_{ω} , denoted by A_{ω} , is an $n \times n$ matrix defined as follows: for each pair of (not necessarily distinct) vertices $v, w \in V(G)$, $(A_{\omega})_{vw} = \sum_{\mathbf{e} \in \mathbf{E}(G)} e^{(0)=v} e^{(1)=w} \chi_{\omega}(\mathbf{e}).$
- $\begin{aligned} (A_{\omega})_{vw} &= \sum_{\mathbf{e} \in \mathbf{E}(G), \mathbf{e}(0) = v, \mathbf{e}(1) = w} \chi_{\omega}(\mathbf{e}). \\ (\text{ii) The edge adjacency matrix of } G \text{ twisted by } \chi_{\omega}, \text{ denoted by } W_{1,\omega}, \text{ is defined as follows:} \\ (W_{1,\omega})_{\mathbf{ab}} &= \chi_{\omega}(\mathbf{b}) = e(\omega(\mathbf{b})), \text{ and } (W_{1,\omega})_{\mathbf{ab}} = 0 \text{ otherwise.} \end{aligned}$

We also define the variants $W'_{1,\omega}$ and $W''_{1,\omega}$ of $W_{1,\omega}$ as: $(W'_{1,\omega})_{\mathbf{ab}} = \chi_{\omega}(\mathbf{a})$ and $(W''_{1,\omega})_{\mathbf{ab}} = \chi_{\omega/2}(\mathbf{a})\chi_{\omega/2}(\mathbf{b})$ if $\mathbf{a} \to \mathbf{b}$, and $(W'_{1,\omega})_{\mathbf{ab}} = (W''_{1,\omega})_{\mathbf{ab}} = 0$ otherwise. Let B_{ω} be a $2m \times 2m$ diagonal matrix whose diagonal entry corresponding to $\mathbf{e} \in \mathbf{E}(G)$ is

Let B_{ω} be a $2m \times 2m$ diagonal matrix whose diagonal entry corresponding to $\mathbf{e} \in \mathbf{E}(G)$ is $e(\omega(\mathbf{e})/2)$. The following lemma says that $W_{1,\omega}$, $W'_{1,\omega}$ and $W''_{1,\omega}$ are similar matrices.

Lemma 2.17.
$$W_{1,\omega} = W_1 B_{2\omega}, W'_{1,\omega} = B_{2\omega} W_1 = B_{2\omega} W_{1,\omega} B_{2\omega}^{-1}$$
 and $W''_{1,\omega} = B_{\omega} W_1 B_{\omega} = B_{\omega} W_{1,\omega} B_{\omega}^{-1}$.
Proof. Straightforwardly verifiable by definitions of $W_{1,\omega}, W'_{1,\omega}$ and $W''_{1,\omega}$.

Remark 2.18. When $\omega = 0$, A_{ω} and $W_{1,\omega}$ specialize to the conventional (vertex) adjacency matrix A and edge adjacency matrix W_1 respectively. In this case, if G has multiple edges or loops, then for distinct vertices v and w, A_{vw} is exactly the number of edges between v and w, and A_{vv} is twice the number of loops at v (corresponding to the two orientations for each loop).

Remark 2.19. A_{ω} and $W_{1,\omega}$ can also be considered as the vertex and edge adjacency matrices of the complex unit gain graph over G with each oriented edge $\mathbf{e} \in \mathbf{E}(G)$ associated with a complex unit gain $\chi_{\omega}(\mathbf{e})$.

As in convention, the *spectrum* of a square matrix M, denoted by Spec M, is the multiset of eigenvalues of M counting multiplicity. For two square matrices M and N, we say M is *isospectral* to N if Spec M = Spec N, and we say the spectrum of M is the complex conjugate (resp. negation) of

the spectrum of N, written as Spec $M = \overline{\text{Spec }N}$ (resp. Spec M = -Spec N), if for each $\lambda \in \text{Spec }M$ of multiplicity m, we have $\overline{\lambda} \in \text{Spec }N$ of multiplicity m (resp. $-\lambda \in \text{Spec }N$ of multiplicity m). We say Spec M is invariant under complex conjugate (resp. negation), if Spec $M = \overline{\text{Spec }M}$ (resp. Spec M = -Spec M).

Some basic properties of A_{ω} and $W_{1,\omega}$ are stated in the following lemma.

Lemma 2.20. (a) A_{ω} is a Hermitian matrix and the eigenvalues of A_{ω} are all real. In particular, $\overline{A_{\omega}} = A_{-\omega} = A_{\omega}^{T}$, and A_{ω} and $A_{-\omega}$ are similar matrices.

- (b) $W_{1,\omega}$ and $W_{1,-\omega}$ are similar matrices.
- (c) Spec $W_{1,\omega}$ is invariant under complex conjugate.

Proof. For (a), we note that

$$(\overline{A_{\omega}})_{vw} = (A_{-\omega})_{vw} = \sum_{\mathbf{e}\in\mathbf{E}(G),\mathbf{e}(0)=v,\mathbf{e}(1)=w} e(-\omega(\mathbf{e})) = \sum_{\mathbf{e}\in\mathbf{E}(G),\mathbf{e}(0)=v,\mathbf{e}(1)=w} e(\omega(\mathbf{e}^{-1}))$$
$$= \sum_{\mathbf{e}\in\mathbf{E}(G),\mathbf{e}(0)=w,\mathbf{e}(1)=v} e(\omega(\mathbf{e})) = (A_{\omega})_{wv}.$$

Hence $\overline{A_{\omega}} = A_{-\omega} = A_{\omega}^T$, which means that A_{ω} is a Hermitian matrix and the eigenvalues of A_{ω} are all real, and . It also follows that A_{ω} and $A_{-\omega}$ are similar since a matrix is similar to its transpose.

For (b), by Lemma 2.17, we only need to show that $W_{1,\omega}''$ and $W_{1,-\omega}''$ are similar. Note that whenever **a** feeds into **b**, $(W_{1,-\omega}'')_{\mathbf{a}\mathbf{b}} = e(-\omega(\mathbf{a} + \mathbf{b})/2) = (W_{1,\omega}'')_{\mathbf{b}^{-1}\mathbf{a}^{-1}} = (W_{1,\omega}'')_{\mathbf{a}^{-1}\mathbf{b}^{-1}}$. Consequently, $W_{1,-\omega}''$ and $W_{1,-\omega}''$ are similar, since $W_{1,-\omega}''$ and $W_{1,\omega}'''$ are similar by a permutation matrix switching **e** and \mathbf{e}^{-1} for all $\mathbf{e} \in \mathbf{E}(G)$.

For (c), note that $W_{1,-\omega} = \overline{W_{1,\omega}}$ which means that $\operatorname{Spec} W_{1,\omega} = \operatorname{Spec} W_{1,-\omega} = \operatorname{Spec} \overline{W_{1,\omega}} = \overline{\operatorname{Spec} W_{1,\omega}}$.

2.4. Basic properties of the L-functions. To simplify the notation, let $\chi_{\omega}(P) := \chi_{\omega}(P^{ab})$.

Definition 2.21. The *L*-function with respect to χ_{ω} is defined to be the following function:

$$L(u,\chi_{\omega}) = L_G(u,\chi_{\omega}) = \prod_{[P]\in\overline{\mathcal{P}}} \left(1 - \chi_{\omega}(P)u^{\tau(P)}\right)^{-1}$$

Lemma 2.22.

$$L(u, \chi_{\omega}) = \exp\left(\sum_{[C]\in\overline{\mathcal{C}}} \frac{\chi_{\omega}(C)}{r(C)} u^{\tau(C)}\right)$$

and in particular

$$z(u) = \exp\left(\sum_{[C]\in\overline{\mathcal{C}}} \frac{1}{r(C)} u^{\tau(C)}\right).$$

Proof.

$$\begin{split} L(u,\chi_{\omega}) &= \prod_{[P]\in\overline{\mathcal{P}}} \left(1 - \chi_{\omega}(P)u^{\tau(P)}\right)^{-1} \\ &= \exp\left(\sum_{[P]\in\overline{\mathcal{P}}} \log\left(\left(1 - \chi_{\omega}(P)u^{\tau(P)}\right)^{-1}\right)\right) \right) \\ &= \exp\left(\sum_{[P]\in\overline{\mathcal{P}}} \sum_{n=1}^{\infty} \frac{\left(\chi_{\omega}(P)u^{\tau(P)}\right)^{n}}{n}\right) \\ &= \exp\left(\sum_{[P]\in\overline{\mathcal{P}}} \sum_{n=1}^{\infty} \frac{\chi_{\omega}(P^{n})u^{\tau(P^{n})}}{n}\right) \\ &= \exp\left(\sum_{[C]\in\overline{\mathcal{C}}} \frac{\chi_{\omega}(C)}{r(C)}u^{\tau(C)}\right), \end{split}$$

which also yields the second identity since $z(u) = L(u, \chi_0)$.

Lemma 2.23. $L(u, \chi_{\omega}) = L(u, \chi_{-\omega}).$

Proof. Note that $[P] \in \overline{\mathcal{P}}$ if and only if $[P^{-1}] \in \overline{\mathcal{P}}$. Then

$$L(u, \chi_{\omega}) = \prod_{[P]\in\overline{\mathcal{P}}} \left(1 - \chi_{\omega}(P)u^{\tau(P)}\right)^{-1}$$

$$= \prod_{[P^{-1}]\in\overline{\mathcal{P}}} \left(1 - \chi_{\omega}(P^{-1})u^{\tau(P^{-1})}\right)^{-1}$$

$$= \prod_{[P]\in\overline{\mathcal{P}}} \left(1 - \chi_{-\omega}(P)u^{\tau(P)}\right)^{-1}$$

$$= L(u, \chi_{-\omega}).$$

Proposition 2.24. Consider $\omega, \omega' \in \Omega$.

- (a) $L(u, \chi_{\omega}) = z(u)$ if and only if $\omega \in \ker(\Omega \twoheadrightarrow \mathcal{X}) = \operatorname{Im}(d_{\mathbb{R}}) \oplus H_1(G, \mathbb{Z})^{\vee}$ if and only if $\phi_1(\omega) \in \operatorname{Im}(d_{\mathbb{R}})$ $H_1(G,\mathbb{Z})^{\vee}.$
- (b) $L(u, \chi_{\omega}) = L(u, \chi_{\omega'})$ if $\phi_1(\omega) \phi_1(\omega') \in H_1(G, \mathbb{Z})^{\vee}$.

Proof. For (a), first note that for any exact 1-form df and any circuit C, we must have $\int_C (\omega + df) =$ $\int_C \omega + \int_C df = \int_C \omega, \text{ which means that } \chi_{\omega}(C) = \chi_{\omega+df}(C).$ Choose g prime paths C_1, \dots, C_g such that $C_1^{ab}, \dots, C_g^{ab}$ form a basis of $H_1(G, \mathbb{Z})$. Then we

have the equivalence of the following statements:

(1) $\omega \in \ker(\Omega \twoheadrightarrow \mathcal{X}) = \operatorname{Im}(d_{\mathbb{R}}) \oplus H_1(G, \mathbb{Z})^{\vee};$ (2) $\phi_1(\omega) \in H_1(G,\mathbb{Z})^{\vee};$ (3) $\chi_{\omega}|_{H_1(G,\mathbb{Z})} = 1;$ (4) $\chi_{\omega}|_{\{C_1,\cdots,C_g\}} = 1;$ (5) $\chi_{\omega}(P) = 1$ for all $P \in \mathcal{P}.$ (6) $\chi_{\omega}(C) = 1$ for all $C \in \mathcal{C}$.

Now we claim that the statements (1)-(6) are equivalent to (7): $L(u, \chi_{\omega}) = z(u)$. It is easy to see by definition that the statements (1)-(6) imply (7). Now suppose $L(u, \chi_{\omega}) = z(u)$. Note that

$$\log L(u,\chi_{\omega}) = \sum_{[C]\in\overline{\mathcal{C}}} \frac{\operatorname{Re}(\chi_{\omega}(C))}{r(C)} u^{\tau(C)} = \sum_{l=1}^{\infty} \left(\sum_{\tau(C)=l, [C]\in\overline{\mathcal{C}}} \frac{\operatorname{Re}(\chi_{\omega}(C))}{r(C)} \right) u^{l}$$

This means for all l, we have $\sum_{\tau(C)=l,[C]\in\overline{\mathcal{C}}} \frac{\operatorname{Re}(\chi_{\omega}(C))}{r(C)} = \sum_{\tau(C)=l,[C]\in\overline{\mathcal{C}}} \frac{1}{r(C)}$. But since $|\chi_{\omega}(C)| = 1$ for all $C \in \mathcal{C}$, this actually means that $\chi_{\omega}(C) = 1$ for all $C \in \mathcal{C}$. As a conclusion, statements (1) - (6) are equivalent to each other.

For (b), similarly we see that $\phi_1(\omega) - \phi_1(\omega') \in H_1(G, \mathbb{Z})^{\vee}$ if and only if $\chi_{\omega}(C) = \chi_{\omega'}(C)$ for all $C \in \mathcal{C}$ if and only if $\chi_{\omega}(P) = \chi_{\omega'}(P)$ for all $P \in \mathcal{P}$. Therefore, $\phi_1(\omega) - \phi_1(\omega') \in H_1(G, \mathbb{Z})^{\vee}$ implies $L(u, \chi_{\omega}) = L(u, \chi_{\omega'})$. (It should be emphasized that the converse statement is not true, unlike in (a). For example, by Lemma 2.23, $L(u, \chi_{\omega}) = L(u, \chi_{-\omega})$)

Remark 2.25. Proposition 2.24 is employed in the proof of spectral antisymmetry in Section 4 (see Lemma 4.7 and Lemma 4.8).

2.5. The Ihara determinant formula and computation of the L-functions. The Ihara zeta function z(u) of a graph G of genus g can be computed from its edge adjacency matrix W_1 and adjacency matrix A by $z(u)^{-1} = \det(I - uW_1)^{-1}$ and the celebrated Ihara determinant formula $z(u)^{-1} = (1 - u^2)^{g-1} \det(I - Au + Qu^2)$ respectively. Here Q is the diagonal matrix with the j-th diagonal entry being $\deg(v_j) - 1$ where $\deg(v_j)$ is the degree of the vertex $v_j \in V(G)$. Such determinant formulas are also true for the L-functions, as summarized in the following theorem (an elementary proof is available in [Ter11]).

Theorem 2.26 (Hashimoto [Has89], Bass [Bas92]). *L*-functions have the following determinantal expressions:

(a) Let p_{ω} be the characteristic polynomial of $W_{1,\omega}$. Then

$$L(u, \chi_{\omega})^{-1} = \det(I - uW_{1,\omega}) = u^{2m} p_{\omega}(u^{-1}).$$

(b) (The Ihara Determinant Formula)

 $L(u, \chi_{\omega})^{-1} = (1 - u^2)^{g-1} \det (I - A_{\omega}u + Qu^2).$

3. Main results

We stick to the terms and notations introduced in Subsection 2.1 and 2.2.

3.1. The canonical character and spectral antisymmetry. For a square matrix M, the spectral radius $\rho(M)$ of M is defined as $\max\{|\lambda| : \lambda \in \text{Spec } M\}$ as in convention.

Theorem 3.1 (Spectral antisymmetry of twisted graph adjacency). For a graph G of genus g, let A be its adjacency matrix, W_1 its edge adjacency matrix and \mathcal{X} its character group. For any $\omega \in \Omega$, let A_{ω} and $W_{1,\omega}$ be respectively the adjacency matrix and edge adjacency matrix of G twisted by χ_{ω} .

- (a) There exists a unique character $\theta \in \mathcal{X}$ such that for all $\omega, \omega' \in \Omega$ with $\underline{\omega} + \underline{\omega'} = \theta$, we have
 - (i) $A_{\omega'}$ is similar to $-A_{\omega}$, Spec $A_{\omega'}$ is the negation of Spec A_{ω} , and
 - (ii) Spec $W_{1,\omega'}$ is the negation of Spec $W_{1,\omega}$.

Moreover, for each $\omega \in \Omega$, we have $\underline{\omega} = \theta$ if Spec $W_{1,\omega} = -$ Spec W_1 .

- (b) θ is a 2-torsion of \mathcal{X} .
- (c) $\theta = 0$ if and only if G is bipartite. In this case, for each $\omega \in \Omega$, Spec A_{ω} and Spec $W_{1,\omega}$ are both invariant under negation.
- (d) $\rho(A_{\omega}) \leq \rho(A)$ and $\rho(W_{1,\omega}) \leq \rho(W_1)$ for all $\omega \in \Omega$.
- (e) $\rho(A_{\omega}) < \rho(A)$ for all $\omega \in \Omega$ such that $\underline{\omega} \notin \{0, \theta\}$.

- (f) If g = 1, then $\rho(W_{1,\omega}) = 1$ for all $\omega \in \Omega$.
- (g) If G is non-bipartite and has genus $g \ge 2$, then $\rho(W_{1,\omega}) < \rho(W_1)$ for all $\omega \in \Omega$ such that $\underline{\omega} \notin \{0, \theta\}$.
- (h) If G is bipartite and has genus $g \ge 2$, then $\rho(W_{1,\omega}) < \rho(W_1)$ for all $\omega \in \Omega$ such that $\underline{\omega}$ is not a 2-torsion of \mathcal{X} .

Remark 3.2. We call θ the canonical character of G and $\omega \in \Omega(G)$ a canonical 1-form of G if $\underline{\omega} = \theta$. For any character $\underline{\omega} \in \mathcal{X}$, we say $\underline{\omega'} = \theta - \underline{\omega}$ is the dual character of $\underline{\omega}$, and ω' is a dual 1-form of ω . Therefore, θ itself is the dual of the trivial character. Note that the canonical character is unique, while there are infinitely many canonical 1-forms.

Remark 3.3. Note that unlike the vertex adjacency matrices, which are all Hermitian, the edge adjacency matrices do not necessarily have real spectra general. They exhibit a more intricate spectral behavior, as can be seen by comparing Theorem 3.1(e) to Theorem 3.1(f)(g)(h). In particular, additional investigations are required to provide a more refined version of Theorem 3.1(h) in the scenario where G is bipartite, accounting for the possible non-trivial 2-torsions $\underline{\omega} \in \mathcal{X}$ such that $\rho(W_{1,\omega}) = \rho(W_1)$ (see Figure 2(c) for an illustrative example).

We defer the technical and lengthy proof of Theorem 3.1 to Section 4.

For a special case of G being a regular graph, as a corollary of Theorem 3.1, we have the following concrete characterization of the spectra of the twisted adjacency matrices of G.

Theorem 3.4. Let G be a (q+1)-regular graph of genus g, and θ be the canonical character of G. Then for $\omega \in \Omega$, we have $\rho(A_{\omega}) \leq q+1$ and $\rho(W_{1,\omega}) \leq q$. Specifically,

- (a) if $\underline{\omega} = 0$, then $q + 1 \in \operatorname{Spec} A_{\omega}$ and $q \in \operatorname{Spec} W_{1,\omega}$, both with multiplicity 1;
- (b) if $\underline{\omega} = \theta$, then $-(q+1) \in \operatorname{Spec} A_{\omega}$ and $-q \in \operatorname{Spec} W_{1,\omega}$, both with multiplicity 1;
- (c) if $\underline{\omega} \notin \{0, \theta\}$, then $\rho(A_{\omega}) < q+1$, and if in addition $g \ge 2$, then $\rho(W_{1,\omega}) < q$;
- (d) if G is bipartite, then $\theta = 0$, and if in addition $g \ge 2$, then $|\lambda| < q$ for all $\lambda \in \operatorname{Spec} W_{1,\omega} \setminus \{\pm q\}$ with $\underline{\omega} = 0$;
- (e) if G is non-bipartite, then $\theta \neq 0$, and we have
 - (i) if $\underline{\omega} = 0$, then $-(q+1) \notin \operatorname{Spec} A_{\omega}$, and if in addition $g \geq 2$, then $|\lambda| < q$ for all $\lambda \in \operatorname{Spec} W_{1,\omega} \setminus \{q\}$, and
 - (ii) if $\underline{\omega} = \theta$, then $q + 1 \notin \operatorname{Spec} A_{\omega}$, and if in addition $g \ge 2$, then $|\lambda| < q$ for all $\lambda \in \operatorname{Spec} W_{1,\omega} \setminus \{-q\}$.

Proof. Let us first consider the twisted adjacency matrices A_{ω} . Note that A_{ω} is Hermitian and its eigenvalues are all real (Lemma 2.20(a)). Now since G is (q + 1)-regular, we get (see, e.g., [Ter11])

(1)
$$\rho(A) = q + 1$$
,

- (2) q+1 is an eigenvalue of A of multiplicity 1, and
- (3) $-(q+1) \in \operatorname{Spec} A$ if and only if G is bipartite.

Consider $\omega, \omega' \in \Omega$ such that $\underline{\omega} = 0$ and $\underline{\omega'} = \theta$. Now since $\underline{\omega + \omega'} = \underline{\omega'} = \theta$, we must have Spec $A_{\omega} =$ Spec A = -Spec $A_{\omega'}$ by Theorem 3.1(a). Then it follows that

- (1) $q + 1 \in \operatorname{Spec} A_{\omega}$ with multiplicity 1,
- (2) $-(q+1) \in \operatorname{Spec} A_{\omega'}$ with multiplicity 1, and
- (3) $-(q+1) \in \operatorname{Spec} A_{\omega}$ if and only if $q+1 \in \operatorname{Spec} A_{\omega'}$ if and only if G is bipartite if and only if $\theta = 0$ (Theorem 3.1(c)).

In addition, by Theorem 3.1(e), $\rho(A_{\omega}) < q+1$ for all $\omega \in \Omega$ such that $\underline{\omega} \notin \{0, \theta\}$.

Now let us consider the twisted edge adjacency matrices $W_{1,\omega}$.

Using the determinant formulas for L-functions in Theorem 2.26, since G is (q+1)-regular which implies $Q = qI_n$ where I_n is the $n \times n$ identity matrix, we have the following relation between Spec $W_{1,\omega}$ and Spec A_{ω} for each $\omega \in \Omega$:

$$\prod_{\kappa \in \operatorname{Spec} W_{1,\omega}} (1 - \kappa u) = (1 - u^2)^{g-1} \prod_{\lambda \in \operatorname{Spec} A_{\omega}} (1 - \lambda u + qu^2).$$

Let $(1 - \kappa_1 u)(1 - \kappa_2 u) = 1 - \lambda u + qu^2$ where $|\kappa_1| \ge |\kappa_2|$. Then by the quadratic formula, we get (1) if $\lambda = q + 1$, then $\kappa_1 = q$ and $\kappa_2 = 1$;

(2) if $\lambda = -(q+1)$, then $\kappa_1 = -q$ and $\kappa_2 = -1$;

(3) if
$$2\sqrt{q} < |\lambda| < q+1$$
, then $\kappa_{1,2} = 2q/(\lambda \pm \sqrt{\lambda^2 - 4q})$, $|\kappa_1| \in (\sqrt{q}, q)$, and $|\kappa_2| \in (1, \sqrt{q})$; and

(4) if $|\lambda| \leq 2\sqrt{q}$, then $\kappa_{1,2} = 2q/(\lambda \pm \sqrt{4q - \lambda^2}\sqrt{-1})$ and $|\kappa_1| = |\kappa_2| = \sqrt{q}$.

Therefore, we further conclude that

(1) if $q + 1 \in \operatorname{Spec} A_{\omega}$ with multiplicity 1, then $q \in \operatorname{Spec} W_{1,\omega}$ with multiplicity 1;

- (2) if $-(q+1) \in \operatorname{Spec} A_{\omega}$ with multiplicity 1, then $-q \in \operatorname{Spec} W_{1,\omega}$ with multiplicity 1; and
- (3) if $\rho(A_{\omega}) < q+1$ and $g \ge 2$, then $\rho(W_{1,\omega}) < q$.

Then the properties of Spec $W_{1,\omega}$ stated in the theorem follow from the properties of Spec A_{ω} as shown above.

Remark 3.5. For some irregular graphs G of genus at least 2, the condition $\underline{\omega} \notin \{0, \theta\}$ can only guarantee that $\rho(A_{\omega}) < \rho(A)$ (Theorem 3.1(e)), but cannot fully guarantee that $\rho(W_{1,\omega}) < \rho(W_1)$ if G is bipartite (Theorem 3.1(h)). (In Subsection 3.4, we give an example of this elusive case, illustrated in Figure 2(c) and Figure 3(c).) However, this assurance is unexceptionally true for regular bipartite graphs of genus at least 2 (Theorem 3.4(c)).

3.2. Some trace formulas.

Definition 3.6. We define the following counting functions of circuits:

- (i) For $\alpha \in H_1(G, \mathbb{Z})$, $N(\alpha, l) = \#\{\text{circuits } C \in \mathcal{C} \mid \tau(C) = l, C^{ab} = \alpha\}.$
- (ii) For $\underline{\alpha} \in Q_{\Lambda}$, $N(\underline{\alpha}, l) = \#\{\text{circuits } C \in \mathcal{C} \mid \tau(C) = l, \underline{C^{\text{ab}}} = \underline{\alpha}\}.$

Definition 3.7. For each $\omega \in \Omega$, define $\mathcal{K}(\omega, l)$ as the *trace* of the twisted edge adjacency matrix $W_{1,\omega}^l$, which we call the *trace distribution* function of order l.

Let C_l be the set of all circuits of length n.

Lemma 3.8. The following are some equivalent definitions of $\mathcal{K}(\omega, n)$:

(a) $\mathcal{K}(\omega, l) = \sum_{C \in \mathcal{C}_l} \chi_{\omega}(C);$ (b) $\mathcal{K}(\omega, l) = \sum_{\lambda \in \operatorname{Spec} W_{1,\omega}} \lambda^l.$

To prove this Lemma we need the notion of *edge-walk*. An *edge-walk* Δ of length N which is a sequence of oriented edges $\mathbf{a}_0 \triangleright \mathbf{a}_1 \triangleright \cdots \triangleright \mathbf{a}_N$ such that $\mathbf{a}_{i+1}(0) = \mathbf{a}_i(1)$ for $i = 0, 1, \cdots, N-1$, of which the *initial oriented edge* is $\Delta(0) := \mathbf{a}_0$ and the *terminal oriented edge* is $\Delta(1) := \mathbf{a}_N$. We also say the walk $\mathbf{a}_1 \cdots \mathbf{a}_N$ is the *associated walk* of Δ , and the walk $\mathbf{a}_1 \cdots \mathbf{a}_{N-1}$ is the *interior walk* of Δ . The *inverse* of Δ is the edge-walk $\Delta^{-1} := \mathbf{a}_N^{-1} \triangleright \cdots \triangleright \mathbf{a}_0^{-1}$. An edge-walk $\Delta' = \mathbf{b}_0 \triangleright \cdots \triangleright \mathbf{b}_M$ is a *sub-edge-walk* of Δ if $\mathbf{b}_j = \mathbf{a}_{i+j}$ for all $j = 0, \cdots, M$ and some $i \in \{0, \cdots, N\}$. We say Δ has a *backtrack* if $\mathbf{a}_{i+1} = \mathbf{a}_i^{-1}$ for some $0 \le i \le N-1$. We say Δ is a *closed edge-walk* at the *base oriented edge* \mathbf{e} if $\mathbf{e} = \mathbf{a}_0 = \mathbf{a}_N$. Note that the associated walk of a closed edge-walk (resp. a closed edge-walk without a backtrack) is always a closed walk (resp. a circuit). (An alternative way of describing edge-walks is to use the notion of oriented line graphs. See, e.g., Kotani and Sunada [KS00]. A non-backtracking edge-walk on G is a walk on the oriented line graph associated with G. Here we will stick to the term "edge-walk" throughout the paper). We also write $\chi_{\omega}(\Delta) = \chi_{\omega}(\Delta) := \chi_{\omega}(\Delta^{\mathrm{ab}})$ for a walk Δ and an edge-walk Δ whose associated walk is Δ . Clearly, $\chi_{\omega}(\Delta^{-1}) = \chi_{-\omega}(\Delta) = \overline{\chi_{\omega}(\Delta)}$. proof of Lemma 3.8. For (a), First note that by definition of the twisted edge adjacency matrix $W_{1,\omega}$, the (e, e)-entry of Spec $W_{1,\omega}^l$ is the sum over the circuits at the base oriented edge e, each twisted by χ_{ω} . Therefore, being the trace of $W_{1,\omega}^l$, $\mathcal{K}(\omega, l)$ is exactly $\sum_{C \in \mathcal{C}_l} \chi_{\omega}(C)$.

(b) follows from the facts that $\mathcal{K}(\omega, l) = \sum_{\lambda' \in \operatorname{Spec} W_{1,\omega}^l} \lambda'$ and $\operatorname{Spec} W_{1,\omega}^l = \{\lambda^l \mid \lambda \in \operatorname{Spec} W_{1,\omega}\}$ (as a multiset).

By Lemma 3.8(a), it is clear that as a function of $\omega \in \Omega$, $\mathcal{K}(\omega, l)$ is continuous and differentiable.

Lemma 3.9. For $\omega, \omega' \in \Omega$, if $\underline{\omega} = \underline{\omega'}$, then $\mathcal{K}(\omega, l) = \mathcal{K}(\omega', l)$ for all l.

Proof. We see from Lemma 4.7 that $W_{1,\omega}$ and $W_{1,\omega'}$ are isospectral whenever $\underline{\omega} = \underline{\omega'}$. Therefore, $\mathcal{K}(\omega, l) = \mathcal{K}(\omega', l)$ by Lemma 3.8(b).

By this lemma, for each $\underline{\omega} \in \mathcal{X}$, we may also write $\mathcal{K}(\underline{\omega}, l) = \mathcal{K}(\omega, l)$, considered as a function over the character group \mathcal{X} .

Lemma 3.10. The following are some basic properties of $\mathcal{K}(\omega, l)$ of a graph G with canonical character θ .

(a) $\max_{\omega \in \Omega} |\mathcal{K}(\omega, l)| = \mathcal{K}(0, l) = N(l)$, which is a nonnegative integer.

- (b) $\mathcal{K}(\omega, l)$ is real and $\mathcal{K}(\omega, l) = \mathcal{K}(-\omega, l)$ for all $\omega \in \Omega$.
- (c) $\mathcal{K}(\theta \underline{\omega}, l) = (-1)^n \mathcal{K}(\underline{\omega}, l)$ for all $\omega \in \Omega$. In particular, $\mathcal{K}(\theta, l) = \mathcal{K}(0, l)$ when l is even, and $\mathcal{K}(\theta, l) = -\mathcal{K}(0, l)$ when l is odd.
- (d) If G is bipartite, then $\mathcal{K}(\omega, l) = 0$ for all $\omega \in \Omega$ when l is odd.

Proof. For (a), note that $\mathcal{K}(\omega, l) = \sum_{C \in \mathcal{C}_l} \chi_{\omega}(C)$ (Lemma 3.8(a)). Therefore, $\mathcal{K}(0, l) = |\mathcal{C}_l| = N(l)$ and $|\mathcal{K}(\omega, l)| \leq \mathcal{K}(0, l)$.

- (b) follows from Lemma 2.20(b)(c).
- (c) follows from Theorem 3.1(a).
- (d) follows from Theorem 3.1(c).

Theorem 3.11 (Trace formulas). Let h be a function analytic on the open disk $D = \{z \in \mathbb{C} \mid z < \rho(W_{1,\omega}) + \epsilon\}$ for some $\epsilon > 0$. Let \hat{h} be the inverse Z-transform of h, i.e., $\hat{h}(l) = 1/(2\pi\sqrt{-1}) \oint_C h(z) z^{-l-1} dz$ where C is a counterclockwise simple closed path entirely in D and about the origin. We have the following identities:

$$\begin{aligned} (a) \ \sum_{\lambda \in \operatorname{Spec} W_{1,\omega}} (h(\lambda) - h(0)) &= \sum_{l \ge 1} \sum_{\alpha \in H_1(G,\mathbb{Z})} \chi_{\omega}(\alpha) N(\alpha, l) \hat{h}(l), \\ (b) \ \int_{\mathcal{X}} \chi_{-\omega}(\alpha) \left(\sum_{\lambda \in \operatorname{Spec} W_{1,\omega}} (h(\lambda) - h(0)) \right) dV_{\omega} &= \operatorname{vol}(\mathcal{X}) \cdot \sum_{l \ge 1} N(\alpha, l) \hat{h}(l), \\ (c) \ for \ \underline{\omega} \in \widehat{Q_{\Lambda}}, \ \sum_{\lambda \in \operatorname{Spec} W_{1,\omega}} (h(\lambda) - h(0)) &= \sum_{l \ge 1} \sum_{\underline{\alpha} \in Q_{\Lambda}} \chi_{\underline{\omega}}(\underline{\alpha}) N(\underline{\alpha}, l) \hat{h}(l), \ and \\ (d) \ for \ \underline{\alpha} \in Q_{\Lambda}, \ \sum_{\underline{\omega} \in \widehat{Q_{\Lambda}}} \chi_{-\underline{\omega}}(\underline{\alpha}) \sum_{\lambda \in \operatorname{Spec} W_{1,\omega}} (h(\lambda) - h(0)) &= |Q_{\Lambda}| \cdot \sum_{l \ge 1} N(\underline{\alpha}, l) \hat{h}(l). \end{aligned}$$

Applying Theorem 3.11 to a special case of $h(z) = z^l$, we derive some Fourier transform formulas

Applying Theorem 3.11 to a special case of $h(z) = z^2$, we derive some Fourier transform formulas as follows.

Theorem 3.12. We have the following identities:

- (a) The Fourier and inverse Fourier transforms between $N(\alpha, l)$ and $\mathcal{K}(\omega, l)$:
 - (i) $\mathcal{K}(\omega, l) = \sum_{\alpha \in H_1(G,\mathbb{Z})} \chi_{\omega}(\alpha) N(\alpha, l)$, and
 - (*ii*) $N(\alpha, l) = \frac{1}{\operatorname{vol}(\mathcal{X})} \int_{\mathcal{X}} \chi_{-\omega}(\alpha) \mathcal{K}(\omega, l) dV_{\omega}.$
- (b) The Fourier and inverse Fourier transforms between $N(\underline{\alpha}, l)$ and $\mathcal{K}(\underline{\omega}, l)$ for $\underline{\alpha} \in Q_{\Lambda}$ and $\underline{\omega} \in \widehat{Q_{\Lambda}}$:

(i)
$$\mathcal{K}(\underline{\omega}, l) = \sum_{\underline{\alpha} \in Q_{\Lambda}} \chi_{\underline{\omega}}(\underline{\alpha}) N(\underline{\alpha}, l), and$$

(*ii*)
$$N(\underline{\alpha}, l) = \frac{1}{|Q_{\Lambda}|} \sum_{\underline{\omega} \in \widehat{Q_{\Lambda}}} \chi_{-\underline{\omega}}(\underline{\alpha}) \mathcal{K}(\underline{\omega}, l)$$

Proof of Theorem 3.11 and Theorem 3.12. We prove Theorem 3.12 first. Note that $\mathcal{K}(\omega, l) = \sum_{C \in \mathcal{C}_l} \chi_{\omega}(C)$ by Lemma 3.8(a), and $\chi_{\omega}(C) = \chi_{\omega}(C^{ab})$ where $C^{ab} \in H^1(G, \mathbb{Z})$ is the abelianization of C. As a result, for all $C^{ab} = \alpha$, we have the identity $\sum_{C \in \mathcal{C}_l, C^{ab} = \alpha} \chi_{\omega}(C) = \chi_{\omega}(\alpha)N(\alpha, l)$. This implies (a)(i), i.e., $\mathcal{K}(\omega, l) = \sum_{\alpha \in H_1(G, \mathbb{Z})} \chi_{\omega}(\alpha)N(\alpha, l)$. If in addition, $\underline{\omega} \in \widehat{Q_\Lambda}$, then for each $\alpha, \alpha' \in H^1(G, \mathbb{Z})$ such that $\underline{\alpha} = \underline{\alpha'} \in Q_\Lambda$ (meaning that $\alpha - \alpha' \in \Lambda$), we have $\chi_{\omega}(\alpha) = \chi_{\omega}(\alpha')$, which can be written also as $\chi_{\underline{\omega}}(\underline{\alpha})$. Hence (b)(i) follows, i.e., $\mathcal{K}(\underline{\omega}, l) = \sum_{\underline{\alpha} \in Q_\Lambda} \chi_{\underline{\omega}}(\underline{\alpha})N(\underline{\alpha}, l)$.

For (a)(ii), applying (a)(i) and the orthogonal relation in Proposition 2.10, we have

$$\begin{split} &\frac{1}{\operatorname{vol}(\mathcal{X})} \int_{\mathcal{X}} \chi_{-\omega}(\alpha) \mathcal{K}(\omega, l) dV_{\omega} \\ = &\frac{1}{\operatorname{vol}(\mathcal{X})} \int_{\mathcal{X}} \chi_{-\omega}(\alpha) \left(\sum_{\beta \in H_1(G, \mathbb{Z})} \chi_{\omega}(\beta) N(\beta, l) \right) dV_{\omega} \\ = &\sum_{\beta \in H_1(G, \mathbb{Z})} N(\beta, l) \left(\frac{1}{\operatorname{vol}(\mathcal{X})} \int_{\mathcal{X}} \chi_{-\omega}(\alpha) \chi_{\omega}(\alpha) dV_{\omega} \right) \\ = &\sum_{\beta \in H_1(G, \mathbb{Z})} N(\beta, l) \delta_{\alpha\beta} \\ = &N(\alpha, l). \end{split}$$

Analogously, for (b)(ii), applying (b)(i) and the orthogonal relation in Proposition 2.13(b), we have

$$\begin{split} &\frac{1}{|Q_{\Lambda}|}\sum_{\underline{\omega}\in\widehat{Q_{\Lambda}}}\chi_{-\underline{\omega}}(\underline{\alpha})\mathcal{K}(\underline{\omega},l)\\ =&\frac{1}{|Q_{\Lambda}|}\sum_{\underline{\omega}\in\widehat{Q_{\Lambda}}}\chi_{-\underline{\omega}}(\underline{\alpha})\left(\sum_{\underline{\beta}\in Q_{\Lambda}}\chi_{\underline{\omega}}(\underline{\beta})N(\underline{\beta},l)\right)\\ =&\sum_{\underline{\beta}\in Q_{\Lambda}}N(\underline{\beta},l)\left(\frac{1}{|Q_{\Lambda}|}\sum_{\underline{\omega}\in\widehat{Q_{\Lambda}}}\chi_{-\underline{\omega}}(\underline{\alpha})\chi_{\underline{\omega}}(\underline{\beta})\right)\\ =&\sum_{\underline{\beta}\in Q_{\Lambda}}N(\underline{\beta},l)\delta_{\underline{\alpha}\underline{\beta}}\\ =&N(\underline{\alpha},l). \end{split}$$

Now we can prove the trace formulas in Theorem 3.11. Note that on the open disk D, we can write $h(z) = h(0) + \sum_{n \ge 1} \hat{h}(l) z^l$. Since $\operatorname{Spec} W_{1,\omega} \subset D$, we have $h(\lambda) - h(0) = \sum_{l \ge 1} \hat{h}(l) \lambda^l$ for each $\lambda \in \operatorname{Spec} W_{1,\omega}$. Note that $\mathcal{K}(\omega, l) = \sum_{\lambda \in \operatorname{Spec} W_{1,\omega}} \lambda^l$ by Lemma 3.8(b). Hence, using

Theorem 3.12(a)(i),

$$\sum_{\lambda \in \operatorname{Spec} W_{1,\omega}} (h(\lambda) - h(0)) = \sum_{\lambda \in \operatorname{Spec} W_{1,\omega}} \left(\sum_{l \ge 1} \hat{h}(l) \lambda^l \right)$$
$$= \sum_{l \ge 1} \left(\sum_{\lambda \in \operatorname{Spec} W_{1,\omega}} \lambda^l \right) \hat{h}(l)$$
$$= \sum_{l \ge 1} \mathcal{K}(\omega, l) \hat{h}(l)$$
$$= \sum_{l \ge 1} \sum_{\alpha \in H_1(G,\mathbb{Z})} \chi_{\omega}(\alpha) N(\alpha, l) \hat{h}(l).$$

For $\alpha \in H_1(G, \mathbb{Z})$, by Theorem 3.12(a)(ii),

$$\int_{\mathcal{X}} \chi_{-\omega}(\alpha) \left(\sum_{\lambda \in \operatorname{Spec} W_{1,\omega}} (h(\lambda) - h(0)) \right) dV_{\omega}$$
$$= \int_{\mathcal{X}} \chi_{-\omega}(\alpha) \left(\sum_{l \ge 1} \mathcal{K}(\omega, l) \hat{h}(l) \right) dV_{\omega}$$
$$= \operatorname{vol}(\mathcal{X}) \cdot \sum_{l \ge 1} N(\alpha, l) \hat{h}(l).$$

For $\underline{\omega} \in \widehat{Q_{\Lambda}}$, by Theorem 3.12(b)(i),

$$\sum_{\lambda \in \operatorname{Spec} W_{1,\omega}} (h(\lambda) - h(0)) = \sum_{l \ge 1} \mathcal{K}(\omega, l) \hat{h}(l)$$
$$= \sum_{l \ge 1} \sum_{\underline{\alpha} \in Q_{\Lambda}} \chi_{\underline{\omega}}(\underline{\alpha}) N(\underline{\alpha}, l) \hat{h}(l).$$

For $\underline{\alpha} \in Q_{\Lambda}$, by Theorem 3.12(b)(ii),

$$\sum_{\underline{\omega}\in\widehat{Q_{\Lambda}}}\chi_{-\underline{\omega}}(\underline{\alpha})\sum_{\lambda\in\operatorname{Spec}W_{1,\omega}}(h(\lambda)-h(0))$$
$$=\sum_{\underline{\omega}\in\widehat{Q_{\Lambda}}}\chi_{-\underline{\omega}}(\underline{\alpha})\sum_{l\geq 1}\mathcal{K}(\omega,l)\hat{h}(l)$$
$$=\sum_{l\leq 1}\left(\sum_{\underline{\omega}\in\widehat{Q_{\Lambda}}}\chi_{-\underline{\omega}}(\underline{\alpha})\mathcal{K}(\omega,l)\right)\hat{h}(l)$$
$$=|Q_{\Lambda}|\cdot\sum_{l\leq 1}N(\underline{\alpha},l)\hat{h}(l).$$

As a corollary of Theorem 3.1 and Theorem 3.12, we have the following result about some vanishing behaviors of $N(\alpha, l)$ and $N(\underline{\alpha}, l)$.

Corollary 3.13. Let θ be the canonical divisor of G.

(a) For $\alpha \in H_1(G,\mathbb{Z})$, if $\chi_{\theta}(\alpha) = 1$, then $N(\alpha, l) = 0$ when l is odd, and if $\chi_{\theta}(\alpha) = -1$, then $N(\alpha, l) = 0$ when l is even.

- (b) If $\theta \in \widehat{Q_{\Lambda}}$, then for $\underline{\alpha} \in Q_{\Lambda}$, if $\chi_{\theta}(\underline{\alpha}) = 1$, then $N(\underline{\alpha}, l) = 0$ when l is odd, and if $\chi_{\theta}(\underline{\alpha}) = -1$, then $N(\underline{\alpha}, l) = 0$ when l is even.
- (c) If G is bipartite, then for all $\alpha \in H_1(G,\mathbb{Z})$, $N(\alpha, l) = 0$ when l is odd, and for all $\underline{\alpha} \in Q_\Lambda$, $N(\underline{\alpha}, l) = 0$ when l is odd.

Proof. By Theorem 3.12(a)(ii) and (b)(ii), we have

$$N(\alpha, l) = \frac{1}{\operatorname{vol}(\mathcal{X})} \int_{\mathcal{X}} \chi_{-\underline{\omega}}(\alpha) \mathcal{K}(\underline{\omega}, l) dV_{\omega}$$

and

$$N(\underline{\alpha}, l) = \frac{1}{|Q_{\Lambda}|} \sum_{\underline{\omega} \in \widehat{Q_{\Lambda}}} \chi_{-\underline{\omega}}(\underline{\alpha}) \mathcal{K}(\underline{\omega}, l).$$

By Lemma 3.10(b)(c), we have $\mathcal{K}(\theta - \underline{\omega}, l) = \mathcal{K}(\underline{\omega} - \theta, l) = (-1)^n \mathcal{K}(\underline{\omega}, l) = (-1)^l \mathcal{K}(-\underline{\omega}, l)$. Therefore,

$$\chi_{-(\theta-\underline{\omega})}(\alpha)\mathcal{K}(\theta-\underline{\omega},l) + \chi_{-(\underline{\omega}-\theta)}(\alpha)\mathcal{K}(\underline{\omega}-\theta,l)$$
$$= (-1)^{l}\chi_{\theta}(\alpha)\left(\chi_{-\underline{\omega}}(\alpha)\mathcal{K}(\underline{\omega},l) + \chi_{\underline{\omega}}(\alpha)\mathcal{K}(-\underline{\omega},l)\right).$$

This just means that if $(-1)^l \chi_{\theta}(\alpha) = -1$, then $N(\alpha, l) = 0$, and (a) follows.

Analogously, if in addition $\theta \in \widehat{Q}_{\Lambda}$, then using the same argument, (b) follows. If G is bipartite, then $\theta = 0$ by Theorem 3.1(c), and (c) follows from (a) and (b). Note that (c) is also a direct consequence of the combinatorial fact that there are no circuits of odd length on a bipartite graph.

3.3. Counting cycles in homology classes.

Definition 3.14. Let $\alpha \in H_1(G, \mathbb{Z})$ and then $\underline{\alpha} \in Q_{\Lambda}$. We define the following counting functions of cycles and prime cycles:

- (i) $\pi(l) = \#\{\text{prime cycles } [P] \in \overline{\mathcal{P}} \mid \tau(P) = l\},\ \pi_c(l) = \#\{\text{cycles } [C] \in \overline{\mathcal{C}} \mid \tau(C) = l\}.$
- (ii) $\pi(\alpha, l) = \#\{\text{prime cycles } [P] \in \overline{\mathcal{P}} \mid \tau(P) = l, P^{ab} = \alpha\}, \\ \pi_c(\alpha, l) = \#\{\text{cycles } [C] \in \overline{\mathcal{C}} \mid \tau(C) = l, C^{ab} = \alpha\}.$
- (iii) $\pi(\underline{\alpha}, l) = \#\{\text{prime cycles } [P] \in \overline{\mathcal{P}} \mid \tau(P) = l, \underline{P^{ab}} = \underline{\alpha}\}, \\ \pi_c(\underline{\alpha}, l) = \#\{\text{cycles } [C] \in \overline{\mathcal{C}} \mid \tau(C) = l, \underline{C^{ab}} = \underline{\alpha}\}.$

Let $[\alpha|k] := \{\beta \in H_1(G, \mathbb{Z}) \mid k\beta = \alpha\}$ and $[\underline{\alpha}|k] := \{\underline{\beta} \in Q_\Lambda \mid k\underline{\beta} = \underline{\alpha}\}$. Let $\mu(n)$ be the Möbius function.

Proposition 3.15. We have the following identities:

$$(a) \ \pi_{c}(\alpha, l) = \sum_{d|l} \left(\sum_{\beta \in [\alpha|(l/d)]} \pi(\beta, d) \right);$$

$$(b) \ \pi_{c}(\underline{\alpha}, l) = \sum_{d|l} \left(\sum_{\underline{\beta} \in [\alpha|(l/d)]} \pi(\underline{\beta}, d) \right);$$

$$(c) \ N(\alpha, l) = \sum_{d|l} d \cdot \left(\sum_{\beta \in [\alpha|(l/d)]} \pi(\beta, d) \right);$$

$$(d) \ N(\underline{\alpha}, l) = \sum_{d|l} d \cdot \left(\sum_{\underline{\beta} \in [\alpha|(l/d)]} \pi(\underline{\beta}, d) \right);$$

$$(e) \ \pi(\alpha, l) = \frac{1}{l} \sum_{d|l} \mu \left(\frac{l}{d} \right) \sum_{\beta \in [\alpha|(l/d)]} N(\beta, d);$$

$$(f) \ \pi(\underline{\alpha}, l) = \frac{1}{l} \sum_{d|l} \mu \left(\frac{n}{d} \right) \sum_{\underline{\beta} \in [\alpha|(l/d)]} N(\underline{\beta}, d).$$

Proof. Note that when $[\alpha|(l/d)]$ is nonempty, for each $\beta \in [\alpha|(l/d)]$ and $P \in \mathcal{P}$ such that $\tau(P) = d$ and $P^{ab} = \beta$ (resp. $\underline{P^{ab}} = \underline{\beta}$), we have $P^{l/d} \in \mathcal{C}$, $\tau(P^{l/d}) = l$ and $(P^{l/d})^{ab} = \alpha$ (resp. $\underline{(P^{l/d})}^{ab} = \underline{\alpha}$). Then (a)-(d) follow from the definitions of the counting functions.

(e) and (f) are Möbius inversions of (c) and (d) respectively. More precisely, for (e),

$$\begin{split} \frac{1}{l} \sum_{d|l} \mu\left(\frac{l}{d}\right) \sum_{\beta \in [\alpha|(l/d)]} N(\beta, d) \\ &= \frac{1}{l} \sum_{d|l} \mu\left(\frac{l}{d}\right) \sum_{\beta \in [\alpha|(l/d)]} \sum_{d'|d} d' \left(\sum_{\gamma \in [\beta|(d/d')]} \pi(\gamma, d')\right) \\ (\text{With } l' = l/d) &= \frac{1}{l} \sum_{d'|l} \sum_{\gamma \in [\alpha|(l/d')]} \sum_{l'|(l/d')} \mu(l') \cdot d' \cdot \pi(\gamma, d') \\ &= \frac{1}{l} \sum_{d'|l} d' \sum_{\gamma \in [\alpha|(l/d')]} \pi(\gamma, d') \left(\sum_{l'|(l/d')} \mu(l')\right) \\ &= \pi(\alpha, l). \end{split}$$

The last equality follows from the property of Möbius function that if l = 1, then $\sum_{d|l} \mu(d) = 1$, and otherwise $\sum_{d|l} \mu(d) = 0$.

(f) follows from an analogous argument.

Remark 3.16. By (a)-(d) of Proposition 3.15, we see that

$$N(\alpha, l) = 0 \quad \Leftrightarrow \quad \pi_c(\alpha, l) = 0 \quad \Leftrightarrow \quad \pi(\beta, d) = 0 \text{ for all } d|l \text{ and } \beta \in [\alpha|(l/d)]$$

and

$$N(\underline{\alpha}, l) = 0 \quad \Leftrightarrow \quad \pi_c(\underline{\alpha}, l) = 0 \quad \Leftrightarrow \quad \pi(\beta, d) = 0 \text{ for all } d|l \text{ and } \underline{\beta} \in [\underline{\alpha}|(l/d)].$$

Recall that by Theorem 3.1(b), the canonical character θ of G must be a 2-torsion of the character group \mathcal{X} of G, or equivalently $\chi_{\theta}(H_1(G,\mathbb{Z})) \subseteq \{\pm 1\}$. Again, for a full-rank sublattice Λ of $H_1(G,\mathbb{Z})$, let $Q_{\Lambda} = H_1(G,\mathbb{Z})/\Lambda$, and accordingly $\widehat{Q_{\Lambda}} = \Lambda^{\vee}/H_1(G,\mathbb{Z})^{\vee}$ is a finite subgroup of \mathcal{X} . Let $\rho_G := \rho(W_1)$ be the spectral radius of the edge adjacency matrix W_1 of G, $M_G = \{\lambda \in \text{Spec } W_1 \mid |\lambda| = \rho_G\}$, and ν_G the cardinality of M_G . Suppose the genus of G is at least 2. Note that ν_G can be interpreted as the greatest common divisor of the lengths of all circuits on G, and by Perron-Frobenius theorm for non-negative matrices, there are exactly ν_G eigenvalues of W_1 with modulus ρ_G and the spectrum of W_1 is invariant under multiplication by $e(1/\nu_G)$ [KS00]. In particular, if G is (q + 1)-regular, then $\rho_G = q$; if G is bipartite, then ν_G is even; if G is regular and non-bipartite, then $\nu_G = 1$; if Gis regular and bipartite, then $\nu_G = 2$.

Theorem 3.17. Suppose the genus g of the graph G is at least 2. Let $\alpha \in H_1(G, \mathbb{Z})$ and correspondingly $\underline{\alpha} \in Q_{\Lambda}$.

(a) π(<u>α</u>, l) = π_c(<u>α</u>, l) = N(<u>α</u>, l) = 0 if one of the following cases occurs:

(i) ν_G ∤ l;
(ii) θ ∈ Q_Λ, χ_θ(<u>α</u>) = 1 and l is odd;
(iii) θ ∈ Q_Λ, χ_θ(<u>α</u>) = -1 and l is even.

(b) As l goes to infinity,

$$\pi(\underline{\alpha}, l) \sim \pi_c(\underline{\alpha}, l) \sim \frac{N(\underline{\alpha}, l)}{l} \sim \frac{\nu_G \rho_G^l}{l |Q_\Lambda|}$$

if one of the following cases occurs:

- (i) $\nu_G \mid l \text{ and } \theta \notin Q_\Lambda$;
- (ii) G is bipartite, $\nu_G \mid l$, and every nonzero element in $\widehat{Q_{\Lambda}}$ is not a 2-torsion of \mathcal{X} ;
- (iii) G is regular and bipartite, and $\nu_G \mid l$.

(c) As l goes to infinity,

$$\pi(\underline{\alpha}, l) \sim \pi_c(\underline{\alpha}, l) \sim \frac{N(\underline{\alpha}, l)}{l} \sim \frac{2\nu_G \rho_G^l}{l|Q_\Lambda|}$$

if one of the following cases occurs:

(i) G is not bipartite, $\nu_G \mid l, \theta \in Q_\Lambda, \chi_\theta(\underline{\alpha}) = 1$ and l is even;

(ii) G is not bipartite, $\nu_G \mid l, \theta \in \widehat{Q}_{\Lambda}, \chi_{\theta}(\underline{\alpha}) = -1$ and l is odd.

Proof. First, we observe that the vanishing behavior of $\pi(\underline{\alpha}, l)$ and $\pi_c(\underline{\alpha}, l)$ is determined by that of $N(\underline{\alpha}, l)$ (Remark 3.16).

Note that $N(l) = \mathcal{K}(0, l) = \sum_{\lambda \in \text{Spec } W_1} \lambda^l$. The following facts are known:

- (1) If $\nu_G \nmid l$, then N(l) = 0, which also implies $N(\alpha, l) = N(\underline{\alpha}, l) = 0$ as stated in (a)(i);
- (2) If $\nu_G \mid l$, then $\mathcal{K}(0, l) \sim \sum_{\lambda \in M_G} \lambda^l = \nu_G \rho_G^l$ as l goes to infinity.

(a)(ii) and (a)(iii) follow from Corollary 3.13(b).

For all the three cases of (b), note that for each $\underline{\omega} \in Q_{\Lambda} \setminus \{\underline{0}\}$, we must have $\rho(W_{1,\omega}) < \rho_G$ by Theorem 3.1 and Theorem 3.4. Therefore, by Theorem 3.11, in either of the three cases of (b),

$$N(\underline{\alpha}, l) = \frac{1}{|Q_{\Lambda}|} \sum_{\underline{\omega} \in \widehat{Q_{\Lambda}}} \chi_{-\underline{\omega}}(\underline{\alpha}) \mathcal{K}(\underline{\omega}, l) \sim \frac{\mathcal{K}(0, l)}{|Q_{\Lambda}|} \sim \frac{\nu_{G} \rho_{G}^{l}}{|Q_{\Lambda}|}$$

as l goes to infinity. Now since $N(\underline{\alpha}, l)$ goes to infinity exponentially as a function of l, by Proposition 3.15, (b) follows.

For both cases of (c), note that the canonical character θ is nontrivial and in \widehat{Q}_{Λ} . Then by Theorem 3.1, $\mathcal{K}(\theta) = (-1)^l \mathcal{K}(0, l)$, and for each $\underline{\omega} \in \widehat{Q}_{\Lambda} \setminus \{\underline{0}, \theta\}$, we must have $\rho(W_{1,\omega}) < \rho_G$. Then in either case of c,

$$N(\underline{\alpha}, l) = \frac{1}{|Q_{\Lambda}|} \sum_{\underline{\omega} \in \widehat{Q_{\Lambda}}} \chi_{-\underline{\omega}}(\underline{\alpha}) \mathcal{K}(\underline{\omega}, l) \sim \frac{(1 + \chi_{\theta}(\underline{\alpha})(-1)^n) \mathcal{K}(0, l)}{|Q_{\Lambda}|} \sim \frac{2\nu_G \rho_G^l}{|Q_{\Lambda}|}$$

as l goes to infinity, and again by Proposition 3.15, (c) follows.

For a special case, now we consider G to be a regular graph and $\Lambda = tH_1(G,\mathbb{Z})$. As can be seen from Lemma 2.5(c), $H_1(G,\mathbb{Z}/t\mathbb{Z}) = H_1(G,\mathbb{Z})/tH_1(G,\mathbb{Z}) = Q_{\Lambda}$. As in convention, elements in $H_1(G,\mathbb{Z}/t\mathbb{Z})$ are called *t*-circulations on G. Consider $\alpha \in H_1(G,\mathbb{Z})$ and its corresponding *t*circulation $\underline{\alpha} \in H_1(G,\mathbb{Z}/t\mathbb{Z})$.

Corollary 3.18. Let G be a (q + 1)-regular graph of genus $g \ge 2$. Let $\alpha \in H_1(G, \mathbb{Z})$ and its corresponding t-circulation $\underline{\alpha} \in H_1(G, \mathbb{Z}/t\mathbb{Z})$.

(a) If G is non-bipartite and t is odd, then

$$\pi(\underline{\alpha}, l) \sim \pi_c(\underline{\alpha}, l) \sim \frac{N(\underline{\alpha}, l)}{l} \sim \frac{q^l}{l \cdot t^g}$$

as l goes to ∞ .

(b) If G is non-bipartite and t is even, then

(i) if $\chi_{\theta}(\alpha) = 1$, then $\chi_{\theta}(\alpha') = 1$ for all $\alpha' \in \underline{\alpha}$ and

$$\begin{cases} \pi(\underline{\alpha}, l) = \pi_c(\underline{\alpha}, l) = N(\underline{\alpha}, l) = 0, & \text{when } l \text{ is odd;} \\ \pi(\underline{\alpha}, l) \sim \pi_c(\underline{\alpha}, l) \sim \frac{N(\underline{\alpha}, l)}{l} \sim \frac{2q^n}{l \cdot t^g}, & \text{when } l \text{ is even and goes to } \infty; \end{cases}$$

(ii) if
$$\chi_{\theta}(\alpha) = -1$$
, then $\chi_{\theta}(\alpha') = -1$ for all $\alpha' \in \underline{\alpha}$ and
 $\int \pi(\alpha, l) = \pi_c(\alpha, l) = N(\alpha, l) = 0$, when l is even:

$$\begin{cases} \pi(\underline{\alpha},l) \sim \pi_c(\underline{\alpha},l) \sim \frac{N(\underline{\alpha},l)}{l} \sim \frac{2q^l}{l \cdot t^g}, & \text{when } l \text{ is odd and goes to } \infty \end{cases}$$



FIGURE 1. A genus-2 graph: two vertices v and w are connected by three paths Δ_0 , Δ_1 and Δ_2 of length l_0 , l_1 , and l_2 respectively.

(c) If G is bipartite, then

$$\begin{cases} \pi(\underline{\alpha}, l) = \pi_c(\underline{\alpha}, l) = N(\underline{\alpha}, l) = 0, & \text{when } l \text{ is odd;} \\ \pi(\underline{\alpha}, l) \sim \pi_c(\underline{\alpha}, l) \sim \frac{N(\underline{\alpha}, l)}{l} \sim \frac{2q^l}{l \cdot t^g}, & \text{when } l \text{ is even and goes to } \infty \end{cases}$$

Proof. It is clear that $|Q_{\Lambda}| = t^g$, and if G is non-bipartite, then the 2-torsion subgroup of χ is a subgroup of $\widehat{Q_{\Lambda}}$ if t is even, and intersects $\widehat{Q_{\Lambda}}$ only at $\underline{0}$ if t is odd. Therefore, the statement is a direct consequence of Theorem 3.4 and Theorem 3.17.

Next, we will explore two examples: the first example is to demonstrate the spectral antisymmetry (Theorem 3.1) and its impacts on the distribution of $\rho(W_{1,\omega})$ and $\mathcal{K}(\omega, n)$ for certain genus-2 graphs, and the second example is about some concrete identities based on the trace formulas and counting functions for the complete graph K_4 .

3.4. An example on distributions of spectral radii and trace function. For this example, note that a genus-2 graph G is made of three paths Δ_0 , Δ_1 and Δ_2 with identical initial vertex vand terminal vertex w (Figure 1). Let l_j be the length of Δ_i for i = 0, 1, 2. Let v_1 and v_2 be the vertices adjacent to v on Δ_1 and Δ_2 respectively. Correspondingly, let \mathbf{e}_1 and \mathbf{e}_2 be the oriented edges from v to v_1 and v_2 respectively. Then $\{\phi_1(d\mathbf{e}_1), \phi_1(d\mathbf{e}_2)\}$ is a basis of $H_1(G, \mathbb{Z})^{\vee}$ in $\mathcal{H}^1(G)$ (Proposition 2.7) and the character group $\mathcal{X} = \mathcal{H}^1(G)/H_1(G, \mathbb{Z})^{\vee}$ is a 2-dimensional real torus such that each $\underline{\omega} \in \mathcal{X}$ has a unique representative $\omega = \omega_1 \cdot \phi_1(d\mathbf{e}_1) + \omega_2 \cdot \phi_1(d\mathbf{e}_2)$ with $0 \leq \omega_{1,2} < 1$.

We consider three cases of this type of graphs: a non-bipartite graph G_1 with $l_0 = 1$, $l_1 = 2$ and $l_2 = 3$; a bipartite graph G_2 with $l_0 = 1$, $l_1 = 3$ and $l_2 = 5$; and a bipartite graph G_3 with $l_0 = 2$, $l_1 = 2$ and $l_2 = 4$.

Distribution of $\rho(W_{1,\omega})$. Figure 2(a), 2(b) and 2(c) show the distributions of spectral radii $\rho(W_{1,\omega})$ over the character groups $\mathcal{X}(G_1)$, $\mathcal{X}(G_2)$ and $\mathcal{X}(G_3)$ respectively. For each of the three cases $\mathcal{X} = \mathcal{X}(G_1)$, $\mathcal{X}(G_2)$, or $\mathcal{X}(G_3)$, the horizontal and vertial axes correspond to $\phi_1(d\mathbf{e}_1)$ and $\phi_2(d\mathbf{e}_2)$ respectively, i.e., each $\omega = \omega_1 \cdot \phi_1(d\mathbf{e}_1) + \omega_1 \cdot \phi_1(d\mathbf{e}_2) \in \mathcal{X}$ has coordinates (ω_1, ω_2) with $0 \leq \omega_{1,2} < 1$.

For the graph G_1 , by spectral antisymmetry (Theorem 3.1), there is a unique canonical character $\theta \in \mathcal{X}(G_1)$ which is a nontrivial 2-torsion since G_1 is non-bipartite. Actually the canonical θ has coordinates (0.5, 0) by Construction 4.2, which can also be observed in Figure 2(a). In particular, $\rho(W_{1,\omega}) = \rho(W_1)$ (of value approximately 1.42405) if $\underline{\omega} \in \{0, \theta\}$, and $\rho(W_{1,\omega}) < \rho(W_1)$ otherwise. Also, one can observe that the distribution of $\rho(W_{1,\omega})$ is symmetric with respect to either the trivial character or the canonical character, as expected from Lemma 2.20(b) and Theorem 3.1(a).

Graphs G_2 and G_3 are both bipartite, which implies by Theorem 3.1(c) that their canonical characters are both trivial. On the other hand, since G_2 and G_3 are not regular graphs (meaning



FIGURE 2. Distributions of spectral radii $\rho(W_{1,\omega})$ over the character groups of graphs (a) G_1 with $l_0 = 1, l_1 = 2, l_2 = 3$, (b) G_2 with $l_0 = 1, l_1 = 3, l_2 = 5$ and (c) G_3 with $l_0 = 2, l_1 = 2, l_2 = 4$. For each case, the horizontal and vertial axes correspond to $\phi_1(d\mathbf{e}_1)$ and $\phi_2(d\mathbf{e}_2)$ respectively.

that Theorem 3.4(c) is not applicable), $\rho(W_{1,\omega}) < \rho(W_1)$ is not guaranteed for all nontrivial characters $\underline{\omega}$ but only for characters which are not 2-torsions by Theorem 3.1(h). For the 2-torsions $\underline{\omega}$ of \mathcal{X} , either $\rho(W_{1,\omega}) < \rho(W_1)$ or $\rho(W_{1,\omega}) = \rho(W_1)$ can happen. In our example, as shown in Figure 2(b)(c), for graph G_2 , $\rho(W_{1,\omega}) < \rho(W_1)$ (of value approximately 1.27065) is true for all nontrivial characters $\underline{\omega}$, while for graph G_3 , $\rho(W_{1,\omega}) = \rho(W_1)$ (of value approximately 1.30216) is satisfied for a nontrivial 2-torsion character $\underline{\omega}$ with coordinates (0, 0.5).

Distribution of $\mathcal{K}(\omega, l)$. Figure 3 shows the distributions of $\mathcal{K}(\omega, l)$ (defined as the trace of $W_{1,\omega}^l$, see Subsection 3.2) over the character groups of graphs G_1 , G_2 and G_3 .

For G_1 (Figure 3(a)), we show the cases of l = 3, 4, 20, 21. In particular, $\mathcal{K}(0,3) = 6, \mathcal{K}(0,4) = 8$, $\mathcal{K}(0,20) = 1278$, and $\mathcal{K}(0,21) = 1574$, which correspond to the number of circuits of length 3, 4, 20 and 21 respectively (Lemma 3.10(a)). One may observe that the distribution of $\mathcal{K}(\omega, l)$ is symmetric with respect to the trivial character for all cases (Lemma 3.10(b)). Note that as shown previously, the canonical character θ has coordinates (0.5,0). One can also observe that the distribution of $\mathcal{K}(\omega, l)$ is symmetric with respect to $\theta/2$ at (0.25, 0) when l = 4, 20, and antisymmetric with respect to $\theta/2$ when l = 3, 21 (Lemma 3.10(c)). It is noticeable that when n is small (3 and 4 in this example), it is possible that $|\mathcal{K}(\omega, l)| = \mathcal{K}(0, l)$ for $\omega \notin \{0, \theta\}$: in the case l = 3, $\mathcal{K}(\omega, 3) = 6$ for all characters ω with the first coordinate 0, and $\mathcal{K}(\omega,3) = -6$ for all characters ω with the first coordinate 0.5; in the case l = 4, $\mathcal{K}(\omega, 4) = 8$ for all characters $\underline{\omega}$ with the second coordinate 0, and $\mathcal{K}(\omega,4) = -8$ for all characters $\underline{\omega}$ with the second coordinate 0.5. Note that for G_1 , there is exactly one dominant eigenvalue $\rho(W_1)$ of W_1 , which means by Theorem 3.1 that there is exactly one dominant eigenvalue $-\rho(W_1)$ of $W_{1,\omega}$ for $\underline{\omega} = \theta$, and $\rho(W_{1,\omega}) < \rho(W_1)$ for all other characters $\omega \notin \{0, \theta\}$. Then as l goes to infinity, $\mathcal{K}(0, l) \sim \rho(W_1)^l$ and $\mathcal{K}(\theta, l) = (-1)^l \mathcal{K}(0, l) \sim (-1)^l \rho(W_1)^l$. As a result, for l sufficiently large (20 and 21 in this example), only when $\underline{\omega} \in \{0, \theta\}$, we can have $|\mathcal{K}(\omega, l)| = \mathcal{K}(0, l).$

Graphs G_2 and G_3 are bipartite. Then the corresponding canonical characters are both trivial, and by Lemma 3.10(d)), $\mathcal{K}(\omega, l)$ vanishes for all characters $\underline{\omega}$ whenever l is odd. Therefore, here we only show cases for even l's. For G_2 (Figure 3(b)) and G_3 (Figure 3(c)), we show the cases of l = 4, 6, 8, 30 and of l = 4, 6, 20, 22 respectively. Recall that we've shown in our explanation of Figure 2 that for G_2 , $\rho(W_{1,\omega}) < \rho(W_1)$ for all nontrivial characters $\underline{\omega}$, and for G_3 , there exist a nontrivial 2-torsion character $\underline{\omega}$ with coordinates (0.5,0) such that $\rho(W_{1,\omega}) = \rho(W_1)$ (actually Spec $W_{1,\omega} = -$ Spec W_1). As a result, for G_2 , when l is a sufficiently large even number (30 in



FIGURE 3. Distributions of $\mathcal{K}(\omega, l)$ over the character groups of graphs G_1 , G_2 and G_3 . For each case, the horizontal and vertial axes correspond to $\phi_1(d\mathbf{e}_1)$ and $\phi_2(d\mathbf{e}_2)$ respectively.

this example), $|\mathcal{K}(\omega, l)| < \mathcal{K}(0, l)$ for all nontrivial characters $\underline{\omega}$, while for G_3 , we see a similar alternating behavior of the distribution of $\mathcal{K}(\omega, l)$ with respect to consecutive large enough even numbers (20 and 22 in this example) as for G_1 with respect to consecutive large enough natural numbers.

3.5. An example on trace formulas and cycle counting. For this example, we consider the complete graph $G = K_4$ with 4 vertices v_1 , v_2 , v_3 and v_4 as shown in Figure 4.

Denote by \mathbf{e}_{ij} the oriented edge from v_i to v_j . Consider a 1-form $\omega = \frac{\varphi}{2\pi} (d\mathbf{e}_{12} + d\mathbf{e}_{23} + d\mathbf{e}_{34} + d\mathbf{e}_{41})$ and the corresponding character χ_{ω} with $\chi_{\omega}(\mathbf{e}_{12}) = \chi_{\omega}(\mathbf{e}_{23}) = \chi_{\omega}(\mathbf{e}_{34}) = \chi_{\omega}(\mathbf{e}_{41}) = e^{\sqrt{-1}\varphi}$ and $\chi_{\omega}(\mathbf{e}_{13}) = \chi_{\omega}(\mathbf{e}_{24}) = 1$. Then the twisted adjacency matrix A_{ω} (indexed with respect to the vertex ordering v_1, v_2, v_3, v_4) is

$$\begin{pmatrix} 0 & e^{\sqrt{-1}\varphi} & 1 & e^{-\sqrt{-1}\varphi} \\ e^{-\sqrt{-1}\varphi} & 0 & e^{\sqrt{-1}\varphi} & 1 \\ 1 & e^{-\sqrt{-1}\varphi} & 0 & e^{\sqrt{-1}\varphi} \\ e^{\sqrt{-1}\varphi} & 1 & e^{-\sqrt{-1}\varphi} & 0 \end{pmatrix}$$

Remark 3.19 (Eigenvalues of A_{ω} and Spec $W_{1,\omega}$). Note that A_{ω} is a circulant matrix, whose eigenvalues can be easily computed as $\mu_1 = 1 + 2\cos\varphi$, $\mu_2 = -1 - 2\sin\varphi$, $\mu_3 = 1 - 2\cos\varphi$, and



FIGURE 4. An example for K_4 .

 $\mu_4 = -1 + 2 \sin \varphi$. Based on the determinantal expressions of the L-functions in Theorem 2.26, since K_4 is 3-regular which implies Q = 2I where I is the 4×4 identity matrix, we see that $\operatorname{Spec} W_{1,\omega} = \{4/(\mu_i \pm \sqrt{\mu_i^2 - 8}) \mid i = 1, \cdots, 4\} \bigcup \{1, 1, -1, -1\}$ as multisets.

Some identities based on the trace formulas. First consider the case $\varphi = 0$, which implies $\mu_1 = 3$, $\mu_2 = \mu_3 = \mu_4 = -1$, and $W_{1,\omega} = W_1$ has eigenvalues 2 of multiplicity 1, $-1/2 + \sqrt{-7/2}$ of multiplicity 3, $-1/2 - \sqrt{-7/2}$ of multiplicity 3, 1 of multiplicity 3 and -1 of multiplicity 2. Now the trace formula in Theorem 3.11(a) degenerates to $\sum_{\lambda \in \text{Spec } W_1} (h(\lambda) - h(0)) = \sum_{l \ge 1} N(l)\hat{h}(l)$ where $N(l) = \mathcal{K}(0, l) = \sum_{\lambda \in \text{Spec } W_1} \lambda^l$ (Lemma 3.8(b) and Lemma 3.10(a)). By the above computation of Spec W_1 , we see that

$$N(l) = \mathcal{K}(0, l)$$

= $2^{l} + 3\left(-\frac{1}{2} + \frac{\sqrt{-7}}{2}\right)^{l} + 3\left(-\frac{1}{2} - \frac{\sqrt{-7}}{2}\right)^{l} + 3 + 2(-1)^{l}$
= 0, 0, 24, 24, 0, 96, 168, 168, 528, 1200, 1848, 3960, 8736, 16128, 31944, ...
for $l = 1, \dots, 15, \dots$.

Consider h(z) to be the exponential function $h(z) = e^z$ where $\hat{h}(l) = 1/(l!)$. Then the trace formula affords the identity

$$e^{2} + 3e + 2e^{-1} + 6e^{-1/2}\cos(\sqrt{7}/2) - 12$$

= $\sum_{l=1}^{\infty} N(l)/(l!) = \frac{24}{3!} + \frac{24}{4!} + \frac{96}{6!} + \frac{168}{7!} + \cdots$
= 5.172675227....

Now let $\omega_0 = (1/2) \cdot (d\mathbf{e}_{12} + d\mathbf{e}_{23} + d\mathbf{e}_{34} + d\mathbf{e}_{41}), \ \omega_1 = (1/6) \cdot (d\mathbf{e}_{12} + d\mathbf{e}_{23} + d\mathbf{e}_{34} + d\mathbf{e}_{41})$ and $\omega_2 = (1/3) \cdot (d\mathbf{e}_{12} + d\mathbf{e}_{23} + d\mathbf{e}_{34} + d\mathbf{e}_{41})$ (corresponding to $\phi = \pi, \pi/3, 2\pi/3$ respectively). Then first we note that $\omega_0 \in H_1(G, \mathbb{Z})^{\vee}$ which means $\underline{\omega}_0 = 0$ and $\underline{\omega}_2 = -\underline{\omega}_1$. Actually using the expressions of the eigenvalues of A_{ω} and $W_{1,\omega}$ in Remark 3.19, we derive Spec $A_{\omega_1} = \text{Spec } A_{\omega_2} = \{0, 2, -1 + \sqrt{3}, -1 - \sqrt{3}\}$ and

Spec
$$W_{1,\omega_1}$$
 = Spec W_{1,ω_2} = { $\pm\sqrt{-2}, 1 \pm \sqrt{-1},$
 $-\frac{1}{2} + \frac{\sqrt{3}}{2} \pm \sqrt{1 + \frac{\sqrt{3}}{2}}\sqrt{-1}, -\frac{1}{2} - \frac{\sqrt{3}}{2} \pm \sqrt{1 - \frac{\sqrt{3}}{2}}\sqrt{-1}, 1, 1, -1, -1$ }

Thus

$$\begin{aligned} \mathcal{K}(\omega_1, l) &= \mathcal{K}(\omega_2, l) \\ &= (1 + (-1)^l)(2^{l/2} + 2) + (1 + \sqrt{-1})^l + (1 - \sqrt{-1})^l + \\ &\left(-\frac{1}{2} + \frac{\sqrt{3}}{2} + \sqrt{1 + \frac{\sqrt{3}}{2}}\sqrt{-1} \right)^l + \left(-\frac{1}{2} + \frac{\sqrt{3}}{2} - \sqrt{1 + \frac{\sqrt{3}}{2}}\sqrt{-1} \right)^l + \\ &\left(-\frac{1}{2} - \frac{\sqrt{3}}{2} + \sqrt{1 - \frac{\sqrt{3}}{2}}\sqrt{-1} \right)^l + \left(-\frac{1}{2} - \frac{\sqrt{3}}{2} - \sqrt{1 - \frac{\sqrt{3}}{2}}\sqrt{-1} \right)^l \\ &= 0, 0, -12, 12, 0, -12, 0, 36, 96, -60, 0, -252, 0, -252, 768, \cdots \\ &\text{for } l = 1, \cdots, 15, \cdots . \end{aligned}$$

Let $\Lambda = \{ \alpha \in H_1(G, \mathbb{Z}) \mid \chi_{\omega_1}(\alpha) = 1 \}$, $\alpha_1 = \mathbf{e}_{12} + \mathbf{e}_{23} + \mathbf{e}_{31}$, and $\alpha_2 = \mathbf{e}_{14} + \mathbf{e}_{43} + \mathbf{e}_{31}$. Then one may verify that Λ is a sublattice of $H_1(G, \mathbb{Z})$ of index 3, $Q_{\Lambda} = \{\underline{\alpha}_0, \underline{\alpha}_1, \underline{\alpha}_2\}$ where $\underline{\alpha}_0 = \Lambda$, $\underline{\alpha}_1 = \alpha_1 + \Lambda$ and $\underline{\alpha}_2 = \alpha_2 + \Lambda$, and $\widehat{Q_{\Lambda}} = \{0, \underline{\omega}_1, \underline{\omega}_2\}$. Moreover, $\chi_{\underline{\omega}_1}(\underline{\alpha}_1) = \chi_{\underline{\omega}_2}(\underline{\alpha}_2) = e^{\sqrt{-1} \cdot 2\pi/3}$ and $\chi_{\underline{\omega}_1}(\underline{\alpha}_2) = \chi_{\underline{\omega}_2}(\underline{\alpha}_1) = e^{\sqrt{-1} \cdot 4\pi/3}$. Then evaluating the formula $N(\underline{\alpha}, l) = \frac{1}{|Q_{\Lambda}|} \sum_{\underline{\omega} \in \widehat{Q_{\Lambda}}} \chi_{-\underline{\omega}}(\underline{\alpha}) \mathcal{K}(\underline{\omega}, l)$ (Theorem 3.12(b)(ii)) in this setting, we get

$$N(\underline{\alpha_0}, l) = (\mathcal{K}(0, l) + 2\mathcal{K}(\omega_1, l))/3$$

= 0, 0, 0, 16, 0, 24, 56, 80, 240, 360, 616, 1152, 2912, 5208, 11160, ...

and

$$N(\underline{\alpha_1}, l) = N(\underline{\alpha_2}, l) = (\mathcal{K}(0, l) - \mathcal{K}(\omega_1, l))/3$$

= 0, 0, 12, 4, 0, 36, 56, 44, 144, 420, 616, 1404, 2912, 5460, 10392, ...

for $l = 1, \dots, 15, \dots$.

Again let $h(z) = e^z$. Applying the trace formula in Theorem 3.11(d) to α_0 , we get

$$\sum_{\omega \in \{0,\omega_1,\omega_2\}} \sum_{\lambda \in \operatorname{Spec} W_{1,\omega}} (h(\lambda) - h(0)) = 3 \sum_{l \ge 1} N(\underline{\alpha_0}, l) \hat{h}(l).$$

Using the computation results of Spec W_1 , Spec W_{1,ω_1} , Spec W_{1,ω_2} and $N(\underline{\alpha_0}, l)$ derived above, we come to the following concrete identity

$$\left(e^2 + 3e + 2e^{-1} + 6e^{-1/2}\cos\left(\sqrt{7}/2\right) - 12\right) + 2\left(2\cos(\sqrt{2}) + 2e\cos(1) + 2e^{(-1+\sqrt{3})/2}\cos\left(\sqrt{1+\sqrt{3}/2}\right) + 2e^{(-1-\sqrt{3})/2}\cos\left(\sqrt{1-\sqrt{3}/2}\right) + 2e + 2e^{-1} - 12\right)$$

= $3\sum_{n\geq 1} \frac{N(\alpha_0, l)}{l!} = 3\left(\frac{16}{4!} + \frac{24}{6!} + \frac{56}{7!} + \frac{80}{8!} + \cdots\right) = 2.141622583\cdots$

Cycle counting. To compute the prime-cycle counting function $\pi(\underline{\alpha}, l)$ and the closed-cycle counting function $\pi_c(\underline{\alpha}, l)$ for $\underline{\alpha} \in Q_{\Lambda} = \{\underline{\alpha}_0, \underline{\alpha}_1, \underline{\alpha}_2\}$, we will use the identities

$$\pi(\underline{\alpha}, l) = \frac{1}{l} \sum_{d|l} \mu\left(\frac{l}{d}\right) \sum_{\underline{\beta} \in [\underline{\alpha}|(l/d)]} N(\underline{\beta}, d)$$

and

$$\pi_c(\underline{\alpha}, l) = \sum_{d|l} \left(\sum_{\underline{\beta} \in [\underline{\alpha}|(l/d)]} \pi(\underline{\beta}, d) \right)$$

in Proposition 3.15 respectively.

Define $\delta_m(a,b) = 1$ if $a \equiv b \mod m$ and $\delta_m(a,b) = 0$ otherwise. Then actually $[\underline{\alpha}_i|k] = \left\{\underline{\alpha}_j|\delta_3(i,jk)\right\}$ for i, j = 0, 1, 2 in this example. Therefore, we get

$$\pi(\underline{\alpha_i}, l) = \frac{1}{l} \sum_{d|n} \mu\left(\frac{n}{d}\right) \sum_{j=0,1,2} \delta_3(i, jl/d) N(\underline{\alpha_j}, d)$$

and

$$\pi_c(\underline{\alpha_i},l) = \sum_{d|l} \sum_{j=0,1,2} \delta_3(i,jl/d) \pi(\underline{\alpha_i},d)$$

for i = 0, 1, 2. In particular,

$$\begin{aligned} \pi(\underline{\alpha_0}, l) &= 0, 0, 0, 4, 0, 4, 8, 8, 24, 36, 56, 92, 224, 368, 744, \cdots, \\ \pi_c(\underline{\alpha_0}, l) &= 0, 0, 0, 4, 0, 4, 8, 12, 32, 36, 56, 102, 224, 376, 744, \cdots, \\ \pi(\underline{\alpha_1}, l) &= \pi(\underline{\alpha_2}, l) = 0, 0, 4, 1, 0, 4, 8, 5, 16, 42, 56, 114, 224, 386, 692, \cdots, \\ \pi_c(\underline{\alpha_1}, l) &= \pi_c(\underline{\alpha_2}, l) = 0, 0, 4, 1, 0, 8, 8, 6, 16, 42, 56, 122, 224, 394, 696, \cdots \end{aligned}$$

for $l = 1, \dots, 15, \dots$.

For the asymptotic behavior of $\pi(\underline{\alpha}, l)$ and $\pi_c(\underline{\alpha}, l)$ for $\underline{\alpha} \in Q_\Lambda$, we note that Case (b)(i) of Theorem 3.17 applies, since $\nu_G = 1$ and $\widehat{Q_\Lambda}$ does not contain the canonical character θ (elements of $\widehat{Q_\Lambda}$ are all 3-torsions of the character group, while θ is a 2-torison). More precisely, we conclude that as l goes to infinity, $\pi(\underline{\alpha}, l)$ (or $\pi_c(\underline{\alpha}, l)$) approximates an equal distribution of value $\nu_G \rho_G^l/(l|Q_\Lambda|) = 2^l/(3l)$ over all $\underline{\alpha} \in Q_\Lambda$.

Now let us consider a case where the canonical character θ is taken into account. Let $\omega_3 = (1/4) \cdot (d\mathbf{e}_{12} + d\mathbf{e}_{23} + d\mathbf{e}_{34} + d\mathbf{e}_{41}), \ \omega_4 = (1/12) \cdot (d\mathbf{e}_{12} + d\mathbf{e}_{23} + d\mathbf{e}_{34} + d\mathbf{e}_{41}) \ \text{and} \ \omega_5 = (5/12) \cdot (d\mathbf{e}_{12} + d\mathbf{e}_{23} + d\mathbf{e}_{34} + d\mathbf{e}_{41}) \ \text{(corresponding to } \phi = \pi/2, \pi/6, 5\pi/6 \ \text{respectively}).$ Then it is easily verifiable that

- (i) $2 \cdot \underline{\omega_4} = \underline{\omega_1}, \ 3 \cdot \underline{\omega_4} = \underline{\omega_3}, \ 4 \cdot \underline{\omega_4} = \underline{\omega_2}, \ 5 \cdot \underline{\omega_4} = -\underline{\omega_4} = \underline{\omega_5},$
- (ii) ω_3 is a canonical 1-form, i.e., $\underline{\omega_3} = \theta$ (guaranteed by the uniqueness of canonical character and the fact that Spec $W_{1,\omega_3} = -$ Spec W_1 which can be derived from the formulas in Remark 3.19),
- (iii) ω_1 is a dual 1-form of ω_4 , and ω_2 is a dual 1-form of ω_5 , and
- (iv) Spec $W_{1,\omega_4} = \operatorname{Spec} W_{1,\omega_5} = -\operatorname{Spec} W_{1,\omega_1} = -\operatorname{Spec} W_{1,\omega_2}$ (by Theorem 3.1(a) and our computation of Spec W_{1,ω_1} and Spec W_{1,ω_2} above).

Thus we have

$$\mathcal{K}(\omega_3, l) = (-1)^l \mathcal{K}(0, l)$$

= 0, 0, -24, 24, 0, 96, -168, 168, -528, 1200, -1848, 3960, -8736, 16128, -31944, ...

and

$$\mathcal{K}(\omega_4, l) = \mathcal{K}(\omega_5, l) = (-1)^l \mathcal{K}(\omega_1, l) = (-1)^l \mathcal{K}(\omega_2, l)$$

= 0, 0, 12, 12, 0, -12, 0, 36, -96, -60, 0, -252, 0, -252, -768, ...

for $l = 1, \dots, 15, \dots$.

Let $\Lambda' = \{ \alpha \in H_1(G, \mathbb{Z}) \mid \chi_{\omega_4}(\alpha) = 1 \}$, and $\underline{\alpha'_i} = i \cdot \alpha_1 + \Lambda'$ for $i = 0, 1, \dots, 5$. Then Λ' is a sublattice of Λ of index 2, $Q_{\Lambda'} = \{\underline{\alpha'_0}, \underline{\alpha'_1}, \dots, \underline{\alpha'_5}\}$ is a cyclic group generated by $\underline{\alpha'_1}$, and

 $\widehat{Q_{\Lambda'}} = \{0, \underline{\omega_1}, \cdots, \underline{\omega_5}\} \text{ is a cyclic group generated by } \underline{\omega_4}. \text{ Moreover, } \chi_{\underline{\omega_4}}(\underline{\alpha'_1}) = e^{\sqrt{-1} \cdot \pi/3}, \text{ and } \underline{\alpha_i} = \alpha'_i \bigsqcup \alpha'_{i+3} \text{ for } i = 0, 1, 2.$

Again, using the formula in Theorem 3.12(b)(ii), we get

$$\begin{split} N(\underline{\alpha_0'},l) &= \frac{1+(-1)^l}{6} (\mathcal{K}(0,l)+2\mathcal{K}(\omega_1)) \\ &= 0, 0, 0, 16, 0, 24, 0, 80, 0, 360, 0, 1152, 0, 5208, 0, \cdots, \\ N(\underline{\alpha_1'},l) &= N(\underline{\alpha_5'},l) = \frac{1-(-1)^l}{6} (\mathcal{K}(0,l)-\mathcal{K}(\omega_1)) \\ &= 0, 0, 12, 0, 0, 0, 56, 0, 144, 0, 616, 0, 2912, 0, 10392, \cdots, \\ N(\underline{\alpha_2'},l) &= N(\underline{\alpha_4'},l) = \frac{1+(-1)^l}{6} (\mathcal{K}(0,l)-\mathcal{K}(\omega_1)) \\ &= 0, 0, 0, 4, 0, 36, 0, 44, 0, 420, 0, 1404, 0, 5460, 0, \cdots, \text{ and} \\ N(\underline{\alpha_3'},l) &= \frac{1-(-1)^l}{6} (\mathcal{K}(0,l)+2\mathcal{K}(\omega_1)) \\ &= 0, 0, 0, 0, 0, 0, 56, 0, 240, 0, 616, 0, 2912, 0, 11160, \cdots \end{split}$$

for $l = 1, \dots, 15, \dots$.

Analogous to the computation in the previous case, we have

$$\pi(\underline{\alpha'_i}, l) = \frac{1}{l} \sum_{d|l} \mu\left(\frac{l}{d}\right) \sum_{j=0,\cdots,5} \delta_6(i, jl/d) N(\underline{\alpha'_j}, d)$$

and

$$\pi_c(\underline{\alpha'_i}, l) = \sum_{d|l} \sum_{j=0,\cdots,5} \delta_3(i, jl/d) \pi(\underline{\alpha'_i}, d)$$

for $i = 0, \dots, 5$. In particular,

$$\begin{aligned} \pi(\underline{\alpha'_0},l) &= 0, 0, 0, 4, 0, 4, 0, 8, 0, 36, 0, 92, 0, 368, 0, \cdots, \\ \pi_c(\underline{\alpha'_0},l) &= 0, 0, 0, 4, 0, 4, 0, 12, 0, 36, 0, 102, 0, 376, 0, \cdots, \\ \pi(\underline{\alpha'_1},l) &= \pi(\underline{\alpha'_5},l) = 0, 0, 4, 0, 0, 0, 8, 0, 16, 0, 56, 0, 224, 0, 692, \cdots, \\ \pi_c(\underline{\alpha'_1},l) &= \pi_c(\underline{\alpha'_5},l) = 0, 0, 4, 0, 0, 0, 8, 0, 16, 0, 56, 0, 224, 0, 696, \cdots, \\ \pi(\underline{\alpha'_2},l) &= \pi(\underline{\alpha'_4},l) = 0, 0, 0, 1, 0, 4, 0, 5, 0, 42, 0, 114, 0, 386, 0, \cdots, \\ \pi_c(\underline{\alpha'_2},l) &= \pi_c(\underline{\alpha'_4},l) = 0, 0, 0, 1, 0, 8, 0, 6, 0, 42, 0, 122, 0, 394, 0, \cdots, \\ \pi(\underline{\alpha'_3},l) &= 0, 0, 0, 0, 0, 0, 8, 0, 32, 0, 56, 0, 224, 0, 744, \cdots, \end{aligned}$$

for $l = 1, \dots, 15, \dots$.

Now since $\widehat{Q_{\Lambda'}}$ contains the canonical character $\theta = \underline{\omega_3}$, we have $\chi_{\theta}(\underline{\alpha'_0}) = \chi_{\theta}(\underline{\alpha'_2}) = \chi_{\theta}(\underline{\alpha'_4})$ and $\chi_{\theta}(\underline{\alpha'_1}) = \chi_{\theta}(\underline{\alpha'_3}) = \chi_{\theta}(\underline{\alpha'_5}) = -1$. Using Theorem 3.17(a) and (c), we obtain the asymptotic behavior of $\pi(\underline{\alpha'}, l)$ and $\pi_c(\underline{\alpha'}, l)$ for $\underline{\alpha'} \in Q_{\Lambda'}$:

(i)
$$\pi(\underline{\alpha'_i}, l) = \pi_c(\underline{\alpha'_i}, l) = 0$$
 if $i + l$ is odd;
(ii) as l goes to infinity, $\pi(\underline{\alpha'_i}, l)$ (or $\pi_c(\underline{\alpha'_i}, l)$) approaches $2\nu_G \rho_G^l/(l|Q_{\Lambda'}|) = 2^l/(3l)$ if $i + l$ is even.

We will break down the proof of Theorem 3.1 into the construction of certain special canonical 1-forms in Construction 4.1 and Construction 4.2, and the proofs of a series of lemmas (Lemma 4.7 to 4.17).

Construction 4.1. (Find a canonical 1-form with respect to an orientation).

Input: A graph G, an orientation O of G, and the set $\mathbf{E}_O(G)$ of edges positively oriented with respect to O.

Output: A canonical 1-form $\omega_O = (1/2) \sum_{\mathbf{e} \in \mathbf{E}_O(G)} d\mathbf{e}$ of G. (Its canonicity will be shown in the proof of Theorem 3.1.)

For a spanning tree T of G, denote by $E(G) \setminus T$ the set of edges of G not contained in T, $\mathbf{E}(G) \setminus T$ the set of oriented edges of G not contained in T, and $\mathbf{E}_O(G) \setminus T$ the set of oriented edges of G not contained in T which are positively oriented with respect to O.

Construction 4.2. (Find a canonical 1-form with respect to a spanning tree and an orientation). **Input:** A graph G of genus g, a spanning tree T of G, and an orientation O of G.

Output: A canonical 1-form $\omega_{(T,O)}$ of G derived as follows: (Its canonicity will be shown in the proof of Theorem 3.1.)

- (1) Since T is a tree, there exists a unique partition $\{V_1, V_2\}$ of V(G) with respect to which T is bipartite. (Such a partition can be derived using the parity of the distance between each two vertices of T. If V(G) is a singleton, then we let $V_1 = V(G)$ and $V_2 = \emptyset$.)
- (2) Construct a function $\kappa : \mathbf{E}_O(G) \setminus T \to \{0, 1/2\}$ in the following way: for each $\mathbf{e} \in \mathbf{E}_O(G) \setminus T$, if \mathbf{e} is not a loop and the two end vertices of \mathbf{e} lie in V_1 and V_2 respectively, then $\kappa(\mathbf{e}) = 0$; otherwise, $\kappa(\mathbf{e}) = 1/2$.

(3)
$$\omega_{(T,O)} = \sum_{\mathbf{e} \in \mathbf{E}_O(G) \setminus T} \kappa(\mathbf{e}) d\mathbf{e}$$

Remark 4.3. Let $\mathcal{X}_{(T,O)} := \{\sum_{i=1}^{g} \omega_i d\mathbf{e}_i \mid 0 \leq \omega_i < 1\}$ where $\{\mathbf{e}_1, \cdots, \mathbf{e}_g\} = \mathbf{E}_O(G) \setminus T$. Then ϕ_1 restricted to $\mathcal{X}_{(T,O)}$ is a bijection from $\mathcal{X}_{(T,O)}$ to \mathcal{X} by Proposition 2.7. Since $\omega_{(T,O)} \in \mathcal{X}_{(T,O)}$, using $\omega_{(T,O)}$ instead of ω_O as a canonical 1-form is more effective in many applications (e.g., in the proofs of Lemma 4.12 and Lemma 4.17).

Lemma 4.4. Let ω_O be the output of Construction 4.1 and $\omega_{(T,O)}$ be the output of Construction 4.2. Then $\underline{\omega_O} = \omega_{(T,O)}$.

Proof. As in the proof of Proposition 2.7, we construct a basis $\{u_1, \dots, u_g\}$ of $H_1(G, \mathbb{Z})$ as $u_i = \mathbf{e}_i + P_i^{ab} \in H_1(G, \mathbb{Z})$ for $i = 1, \dots, g$. Here $\{\mathbf{e}_1, \dots, \mathbf{e}_g\} = \mathbf{E}_O(G) \setminus T$ and P_i is the unique path from $\mathbf{e}_i(1)$ to $\mathbf{e}_i(0)$ contained in the spanning tree T.

To prove $\underline{\omega_O} = \underline{\omega_{(T,O)}}$, it suffices to show that for $i = 1, \dots, g$, $\omega_O(u_i) - \underline{\omega_{(T,O)}}(u_i)$ is an integer. By Construction 4.1 and Construction 4.2, we have $\omega_O = (1/2) \sum_{\mathbf{e} \in \mathbf{E}_O(G)} d\mathbf{e}$ and $\underline{\omega_{(T,O)}} = \sum_{\mathbf{e} \in \mathbf{E}_O(G) \setminus T} \kappa(\mathbf{e}) d\mathbf{e}$. Then $\omega_O(u_i) = 1/2 + \omega_O(P_i^{ab})$ and $\underline{\omega_{(T,O)}}(u_i) = \kappa(\mathbf{e}_i)$. Note that $2 \cdot \omega_O(P_i^{ab})$ is an integer with the same parity as $\tau(P_i)$.

If \mathbf{e}_i is not a loop and its two end vertices lie in V_1 and V_2 respectively, then $\tau(P_i)$ must be odd, which implies $\omega_O(u_i)$ is an integer. By Construction 4.2, $\omega_{(T,O)}(u_i) = \kappa(\mathbf{e}_i) = 0$ in this case. Otherwise, both the end vertices of \mathbf{e}_i lie in either V_1 or V_2 . In this case, $\tau(P_i)$ must be even. Then $\omega_O(u_i) - 1/2$ is an integer and $\omega_{(T,O)}(u_i) = \kappa(\mathbf{e}_i) = 1/2$. Again, $\omega_O(u_i) - \omega_{(T,O)}(u_i)$ is an integer as desired.

Lemma 4.5. Let ω_O be the output of Construction 4.1. Then for all $\omega, \omega' \in \Omega$ such that $\omega' = \omega + \omega_O$, we have $A_{\omega'} = -A_{\omega}$ and $W_{1,\omega'} = -W_{1,\omega}$.

Proof. Since $\omega_O = (1/2) \sum_{\mathbf{e} \in \mathbf{E}_O(G)} d\mathbf{e}$, we have $\chi_{\omega_O}(\mathbf{e}) = -1$ for all $\mathbf{e} \in \mathbf{E}(G)$. Then $\chi_{\omega'}(\mathbf{e}) = \chi_{\omega_O}(\mathbf{e})\chi_{\omega}(\mathbf{e}) = -\chi_{\omega}(\mathbf{e})$.

Then $(A_{\omega'})_{vw} = \sum_{\mathbf{e}\in \mathbf{E}(G), \mathbf{e}(0)=v, \mathbf{e}(1)=w} \chi_{\omega'}(\mathbf{e}) = -\sum_{\mathbf{e}\in \mathbf{E}(G), \mathbf{e}(0)=v, \mathbf{e}(1)=w} \chi_{\omega}(\mathbf{e}) = -(A_{\omega})_{vw}$ for all $v, w \in V(G)$. Moreover, $(W_{1,\omega'})_{\mathbf{ab}} = \chi_{\omega'}(\mathbf{b}) = \chi_{\omega_O}(\mathbf{b})\chi_{\omega}(\mathbf{b}) = -\chi_{\omega}(\mathbf{b}) = -(W_{1,\omega})_{\mathbf{ab}}$ if **a** feeds into **b**; $(W_{1,\omega'})_{\mathbf{ab}} = -(W_{1,\omega})_{\mathbf{ab}} = 0$ otherwise.

Lemma 4.6. Let $\omega_{(T,O)}$ be the output of Construction 4.2. Let $\omega = \sum_{i=1}^{g} \omega_i d\mathbf{e}_i$ where $\{\mathbf{e}_1, \cdots, \mathbf{e}_g\} = \mathbf{E}_O(G) \setminus T$, and $\omega' = \omega_{(T,O)} - \omega$. Then

(a) $A_{\omega'}$ is similar to $-A_{\omega}$, and

(b) Spec $W_{1,\omega'} = -$ Spec $W_{1,\omega}$.

Proof. Let V_1, V_2 be as in Construction 4.2 where $V_1 = \{v_1, \dots, v_k\}$ and $V_2 = \{v_{k+1}, \dots, v_n\}$; let $\mathbf{E}_1 = \{\mathbf{e} \in \mathbf{E}(G) \mid \mathbf{e}(1) \in V_1\} = \{\mathbf{a}_1, \dots, \mathbf{a}_h\}$ and $\mathbf{E}_2 = \{\mathbf{e} \in \mathbf{E}(G) \mid \mathbf{e}(1) \in V_2\} = \{\mathbf{a}_{h+1}, \dots, \mathbf{a}_{2m}\}$. Specifically, $[x_1, \dots, x_n]^T$ is an eigenvector of A_{ω} with eigenvalue λ , if and only if

$$[x_1,\cdots,x_k,-x_{k+1},\cdots,-x_n]^2$$

is an eigenvector of $A_{\omega'}$ with eigenvalue $-\lambda$.

Specifically, $[x_1, \dots, x_{2m}]^T$ is an eigenvector of $W_{1,-\omega}$ with eigenvalue λ , if and only if

$$[x_1,\cdots,x_h,-x_{h+1},\cdots,-x_{2m}]^T$$

is an eigenvector of $W_{1,\omega'}$ with eigenvalue $-\lambda$.

Note that by construction, $\omega_{(T,O)} = \sum_{i=1}^{g} \kappa(\mathbf{e}_i) d\mathbf{e}_i$, $\omega = \sum_{i=1}^{g} \omega_i d\mathbf{e}_i$ and $\omega' = \sum_{i=1}^{g} (\kappa(\mathbf{e}_i) - \omega_i) d\mathbf{e}_i$.

For (a), consider an eigenvector $[x_1, \dots, x_n]^T$ of $A_{-\omega}$ with λ being the associated eigenvalue. Then $\sum_{j=1}^n (A_{-\omega})_{ij} x_j = \lambda x_i$ for each $i = 1, \dots, n$. We claim that $[x'_1, \dots, x'_n]^T = [x_1, \dots, x_k, -x_{k+1}, \dots, -x_n]^T$ is an eigenvector of $A_{\omega'}$ whose associated eigenvalue is $-\lambda$.

Consider an oriented edge $\mathbf{e} \in \mathbf{E}(G)$ such that $\mathbf{e}(0) = v_i$ and $\mathbf{e}(1) = v_j$. Then by construction, one may verify that

$$\chi_{\omega_{(T,O)}}(\mathbf{e}) = e\left(\kappa(\mathbf{e})\right) = t_{ij} := \begin{cases} 1, & \text{one of } v_i \text{ and } v_j \text{ lies in } V_1 \text{ and the other in } V_2; \\ -1, & \text{both } v_i \text{ and } v_j \text{ lie in either } V_1 \text{ or } V_2. \end{cases}$$

Now we have $(A_{\omega'})_{ij} = \sum_{\mathbf{e}\in \mathbf{E}(G), \mathbf{e}(0)=v_i, \mathbf{e}(1)=v_j} \chi_{-\omega}(\mathbf{e})\chi_{\omega_{(T,O)}}(\mathbf{e}) = (A_{-\omega})_{ij}t_{ij}$. For all $1 \le j \le n$, one can see that $t_{ij}x'_j = -x_j$ for $1 \le i \le k$, and $t_{ij}x'_j = x_j$ for $k+1 \le i \le n$.

Therefore,

$$\sum_{j=1}^{n} (A_{\omega'})_{ij} x'_{j} = \sum_{j=1}^{n} (A_{-\omega})_{ij} t_{ij} x'_{j} = \begin{cases} -\sum_{j=1}^{n} (A_{-\omega})_{ij} x_{j} = -\lambda x_{i} = -\lambda x'_{i}, & \text{for } 1 \le i \le k; \\ \sum_{j=1}^{n} (A_{-\omega})_{ij} x_{j} = \lambda x_{i} = -\lambda x'_{i}, & \text{for } k+1 \le i \le n. \end{cases}$$

Hence we conclude that $[x'_1, \dots, x'_n]^T$ is an eigenvector of $A_{\omega'}$ whose associated eigenvalue is $-\lambda$ as claimed. Note that the procedure also works reversely. Therefore, $\operatorname{Spec} A_{\omega'} = -\operatorname{Spec} A_{-\omega}$. By Lemma 2.20, $A_{\omega'}$ and A_{ω} are Hermitian matrices, and A_{ω} is similar to $A_{-\omega}$. Then $A_{\omega'}$ is similar to $-A_{\omega}$.

An analogous argument can be applied to (b). Let $[x_1, \dots, x_{2m}]^T$ be an eigenvector of $W_{1,-\omega}$ whose associated eigenvalue is λ . Then $\sum_j (W_{1,-\omega})_{ij} x_j = \sum_{\mathbf{a}_i \to \mathbf{a}_j} \chi_{-\omega}(\mathbf{a}_j) x_j = \lambda x_i$. We verify that

$$[x'_1, \cdots, -x'_{2m}]^T = [x_1, \cdots, x_h, -x_{h+1}, \cdots, -x_{2m}]^T$$

is an eigenvector of $W_{1,\omega'}$ whose associated eigenvalue is $-\lambda$ as follows.

By construction, $\chi_{\omega_{(T,O)}}(\mathbf{a}_j) = -1$ if $\{\mathbf{a}_j(0), \mathbf{a}_j(1)\} \subset V_1$ or $\{\mathbf{a}_j(0), \mathbf{a}_j(1)\} \subset V_2$; otherwise, $\chi_{\omega_{(T,O)}}(\mathbf{a}_j) = 1$. Note that if $\mathbf{a}_i \to \mathbf{a}_j$, then $\mathbf{a}_i(1) = \mathbf{a}_j(0)$. Hence we conclude that for $\mathbf{a}_i \to \mathbf{a}_j$,

$$\chi_{\omega'}(\mathbf{a}_j) = \chi_{\omega_{(T,O)}}(\mathbf{a}_j)\chi_{-\omega}(\mathbf{a}_j) = \begin{cases} \chi_{-\omega}(\mathbf{a}_j), & \text{if } \mathbf{a}_i \in \mathbf{E}_1, \, \mathbf{a}_j \in \mathbf{E}_2 \text{ or } \mathbf{a}_i \in \mathbf{E}_2, \, \mathbf{a}_j \in \mathbf{E}_1; \\ -\chi_{-\omega}(\mathbf{a}_j), & \text{if } \{\mathbf{a}_i, \mathbf{a}_j\} \in \mathbf{E}_1 \text{ or } \{\mathbf{a}_i, \mathbf{a}_j\} \in \mathbf{E}_2. \end{cases}$$

Now for each $i = 1, \dots, h$, Therefore,

$$\sum_{j} (W_{1,\omega'})_{ij} x'_{j} = \sum_{\mathbf{a}_{i} \to \mathbf{a}_{j}} \chi_{\omega'}(\mathbf{a}_{j}) x'_{j} = \sum_{\mathbf{a}_{i} \to \mathbf{a}_{j}} \chi_{\omega_{(T,O)}}(\mathbf{a}_{j}) \chi_{-\omega}(\mathbf{a}_{j}) x'_{j}$$
$$= \begin{cases} \sum_{\mathbf{a}_{i} \to \mathbf{a}_{j}} \chi_{-\omega}(\mathbf{a}_{j}) x_{j} = \lambda x_{i} = -\lambda x'_{i}, & \text{for } 1 \le i \le h; \\ -\sum_{\mathbf{a}_{i} \to \mathbf{a}_{j}} \chi_{-\omega}(\mathbf{a}_{j}) x_{j} = -\lambda x_{i} = -\lambda x'_{i}, & \text{for } h+1 \le i \le 2m. \end{cases}$$

Hence we conclude that $[x'_1, \dots, x'_{2m}]^T$ is an eigenvector of $W_{1,\omega'}$ whose associated eigenvalue is $-\lambda$. As in the proof of (a), this procedure also works reversely. Since $W_{1,-\omega}$ is similar to $W_{1,\omega}$ by Lemma 2.20, this means Spec $W_{1,\omega'} = -$ Spec $W_{1,\omega}$.

Lemma 4.7. For $\omega, \omega' \in \Omega$, if $\underline{\omega} = \underline{\omega'}$, then

- (a) A_{ω} is similar to $A_{\omega'}$;
- (b) $W_{1,\omega}$ is isospectral to $W_{1,\omega'}$.

Proof. For (a), we first consider the case $\phi_1(\omega) = \phi_1(\omega')$, i.e., $\omega' = \omega + df$ where df is an exact 1-form. Then note that for $v_i, v_j \in V(G)$, $(A_{\omega})_{ij} = \sum_{\mathbf{e} \in \mathbf{E}(G), \mathbf{e}(0) = v_i, \mathbf{e}(1) = v_j} \chi_{\omega}(\mathbf{e})$ and

$$(A_{\omega'})_{ij} = \sum_{\mathbf{e}\in\mathbf{E}(G),\mathbf{e}(0)=v_i,\mathbf{e}(1)=v_j} \chi_{\omega+df}(\mathbf{e})$$
$$= \sum_{\mathbf{e}\in\mathbf{E}(G),\mathbf{e}(0)=v_i,\mathbf{e}(1)=v_j} e(\omega(\mathbf{e}) + df(\mathbf{e}))$$
$$= \sum_{\mathbf{e}\in\mathbf{E}(G),\mathbf{e}(0)=v_i,\mathbf{e}(1)=v_j} e(\omega(\mathbf{e}))e(f(v_j) - f(v_i))$$
$$= e(-f(v_i))(A_{\omega})_{vw}e(f(v_j))$$

Let B_f be an $n \times n$ diagonal matrix with the (i, i)-entry being $e(f(v_i))$. Then it follows that A_{ω} is similar to $A_{\omega'}$ by $A_{\omega'} = B_f^{-1} A_{\omega} B_f$.

Now we assume in general that $\underline{\omega} = \underline{\omega}'$. By Proposition 2.7, this means that there exist an exact form df such that $(\omega + df) - \omega' = \sum_{i=1}^{g} c_i d\mathbf{e}_i$ with $c_i \in \mathbb{Z}$. Hence, $A_{\omega+df} = A_{\omega'}$, and consequently A_{ω} is similar to $A_{\omega'}$ since A_{ω} is similar to $A_{\omega+df}$ as shown above.

For (b), we note that by Proposition 2.24(b), $L(u, \chi_{\omega}) = L(u, \chi_{\omega'})$. Then $W_{1,\omega}$ and $W_{1,\omega'}$ are isospectral since they must have the same characteristic polynomial by Theorem 2.26(a).

Lemma 4.8. Consider a graph G and its space of characters \mathcal{X} . For each character $\underline{\omega} \in \mathcal{X}$, if $\operatorname{Spec} W_{1,\omega} = \operatorname{Spec} W_1$, then we must have $\underline{\omega} = 0$.

Proof. If Spec $W_{1,\omega} = \text{Spec } W_1$, then $L(u, \chi_{\omega}) = z(u)$ by Theorem 2.26(a). Hence by Proposition 2.24(a), we conclude that $\underline{\omega} = 0$.

Remark 4.9. Lemma 4.8 actually also follows from the twisted Perron-Frobenius theorem by Adachi and Sunada [AS87b]. An alternative proof of the above lemma comes from Theorem A(3)(4) and of [AS87b] which can be restated in our setting as follows: Let R be the spectral radius of W_1 . Then

(i) R is an eigenvalue of W_1 of multiplicity 1; and

(ii) for $\omega \in \Omega(G)$, $W_{1,\omega}$ has an eigenvalue at $R \cdot e(t)$ if and only if $\chi_{\omega}(P) = e(t \cdot \tau(P))$ for all $P \in \mathcal{P}$. In this case, Spec $W_{1,\omega} = e(t) \cdot \text{Spec } W_1$ and every eigenvalue of $W_{1,\omega}$ of absolute value R has multiplicity 1.

By this theorem, if Spec $W_{1,\omega}$ = Spec W_1 , then by (i), Spec $W_{1,\omega}$ has an eigenvalue of multiplicity 1 at R, and by (ii), $\chi_{\omega}(P^{ab}) = 1$ for all $P \in \mathcal{P}$. Since we can always choose $P_1, \dots, P_g \in \mathcal{P}$ such that $\{P_1^{ab}, \dots, P_g^{ab}\}$ is a basis of $H_1(G, \mathbb{Z})$, we conclude that $\chi_{\omega}(\alpha) = 1$ for all $\alpha \in H_1(G, \mathbb{Z})$. (For example, we may let $P_i = \mathbf{e}_i A_i$ as in the proof of Proposition 2.7.) As a result, $\underline{\omega}$ must be the trivial character of $H_1(G, \mathbb{Z})$ as claimed.

Lemma 4.10. Consider a graph G and an arbitrary path Δ on G. There exists $a \in \mathbb{N}$ such that for each pair of vertices v and v' (not necessarily distinct), and each non-negative integer b, we have either

- (i) there exists no walk from v to v' of length ax + b for all $x \in \mathbb{N}$; or
- (ii) there exists some $x_0 \in \mathbb{N}$ such that for all $x > x_0$, there exists a walk from v to v' of length ax + b which contains Δ as a sub-walk.

Proof. First, we claim that there exists $a \in \mathbb{N}$ such that for each vertex v, there exists a closed walk Υ_v at v of length a which contains Δ as a sub-walk.

For each vertex v, let Υ'_v be a closed walk at v which is constituted of a walk from v to $\Delta(0)$, the walk Δ , and a walk from $\Delta(1)$ to v. Then we may let a be a common multiple of the lengths of all Υ'_v , and $a_v = a/\tau(\Upsilon'_v)$. As a result, $\Upsilon_v = (\Upsilon'_v)^{a_v}$ is a closed walk at v of length a which contains Δ as a sub-walk as desired.

Now for each $v, v' \in V(G)$, suppose (i) is not satisfied, i.e., for some x_0 , there exists a walk $\Upsilon'_{v,v'}$ from v to v' of length $ax_0 + b$. Then for all $x > x_0$, the walk $\Upsilon_{(v,v'),ax+b}$ constituted of $(\Upsilon_v)^{x-x_0}$ followed by $\Upsilon'_{v,v'}$ is a walk from v to v' of length ax + b which contains Δ as a sub-walk. \Box

We say two walks Δ and Δ' of the same length with the same initial vertex and the same terminal vertex are *coherent* with respect to χ_{ω} if $\chi_{\omega}(\Delta) = \chi_{\omega}(\Delta')$, and *incoherent* with respect to χ_{ω} if $\chi_{\omega}(\Delta) \neq \chi_{\omega}(\Delta')$.

Lemma 4.11. $\rho(A_{\omega}) \leq \rho(A)$ for all $\omega \in \Omega$. In addition, $\rho(A_{\omega}) < \rho(A)$ if and only if there exist a pair of incoherent walks with respect to χ_{ω} .

Proof. We will employ Gelfand's formula which is an identity relating the spectral radius of a matrix to a limit of matrix norms, namely,

(Gelfand's formula.) for any square matrix M and any matrix norm $\|\cdot\|$, we have $\rho(M) = \lim_{k \to \infty} \|M^k\|^{1/k}$.

For convenience, we will use the ∞ -norm of A^k_{ω} ,

$$||A_{\omega}^{k}|| := \max_{1 \le i \le n} \sum_{j=1}^{n} |(A_{\omega}^{k})_{ij}|.$$

This affords

$$\rho(A_{\omega}) = \lim_{k \to \infty} \|A_{\omega}^k\|^{1/k}$$
$$= \lim_{k \to \infty} \left(\max_{1 \le i \le n} \sum_{j=1}^n |(A_{\omega}^k)_{ij}| \right)^{1/k}.$$

33

1

Denote the set of all walks from v_i to v_j by $\Gamma(i, j)$ and the set of all walks from v_i to v_j of length k by $\Gamma_k(i, j)$. Then we observe that

$$(A^k_{\omega})_{ij} = \sum_{\Delta \in \Gamma_k(i,j)} \chi_{\omega}(\Delta).$$

Immediately, we can get that $|(A_{\omega}^{k})_{ij}| \leq \#\Gamma_{k}(i,j) = A_{ij}^{k}$, which implies that $\rho(A_{\omega}) \leq \rho(A)$ for all $\omega \in \Omega$. In addition, note that if all walks in $\Gamma_{k}(i,j)$ are coherent with respect to χ_{ω} , then $|(A_{\omega}^{k})_{ij}| = \#\Gamma_{k}(i,j) = A_{ij}^{k}$. Therefore, by Gelfand's formula, we conclude that $\rho(A_{\omega}) = \rho(A)$ if there are no pairs of incoherent paths with respect to χ_{ω} .

It remains to show that conversely we only need to find one pair of incoherent walks to make $\rho(A_{\omega}) < \rho(A)$.

Suppose now that walks $\Delta, \Delta' \in \Gamma_k(s, t)$ are incoherent for some $v_s, v_t \in V(G)$. Then by Lemma 4.10, there exists $a \in \mathbb{N}$ such that for all $v_i, v_j \in V(G)$, we have either

(i) $\Gamma_{ax}(i,j) = \emptyset$ for all $x \in \mathbb{N}$, or

(ii) there exists a walk $\Upsilon \in \Gamma_{ax}(i, j)$ containing Δ as a sub-walk for all x large enough.

For Case (i), we have $(A_{\omega}^{ax})_{ij} = A_{ij}^{ax} = 0$ for all $x \in \mathbb{N}$. For Case (ii), we note that we can always get another walk $\Upsilon' \in \Gamma_{ax}(i, j)$ by replacing a sub-walk Δ in Υ (if there are several copies of Δ contained in Υ , just choose one) by Δ' and leaving the remaining part unchanged. By this construction, Υ and Υ' must be incoherent, since Δ and Δ' are incoherent. As a result, we see that $|(A_{\omega}^{ax})_{ij}| < A_{ij}^{ax}$ for all x large enough. Furthermore, combining Case (i) and Case (ii), we conclude that, there exists 0 < c < 1 and a large enough $y \in \mathbb{N}$ such that $|(A_{\omega}^{y})_{ij}| \leq cA_{ij}^{y}$ for all $i, j = 1, \dots, n$.

Therefore,
$$|(A_{\omega}^{ky})_{ij}| \le c^k A_{ij}^{ky}$$
 and $||A_{\omega}^{ky}|| = \max_{1 \le i \le n} \sum_{j=1}^n |(A_{\omega}^{ky})_{ij}| \le c^k ||A^{ky}||.$

Finally, we have

1

$$\rho(A_{\omega}) = \lim_{k \to \infty} \|A_{\omega}^{ky}\|^{1/(ky)} \le \lim_{k \to \infty} \left(c^k \|A^{ky}\|\right)^{1/(ky)} = c^{1/y}\rho(A) < \rho(A)$$

as stated.

Lemma 4.12.
$$\rho(A_{\omega}) < \rho(A)$$
 for all $\omega \in \mathcal{X}_{(T,O)} \setminus \{0, \omega_{(T,O)}\}$ (see Remark 4.3).

Proof. We only consider $\omega \in \mathcal{X}_{(T,O)}$ here, i.e., $\omega = \sum_{\mathbf{e} \in \mathbf{E}_O(G) \setminus T} \omega(\mathbf{e}) d\mathbf{e}$ with $0 \leq \omega(\mathbf{e}) < 1$ for each $\mathbf{e} \in \mathbf{E}_O(G) \setminus T$. First note that it follows from Lemma 4.6 that $\rho(A_{\omega_{(T,O)}}) = \rho(A)$. We will show that for all other ω , we have a strict inequality $\rho(A_\omega) < \rho(A)$.

By Lemma 4.11, to make $\rho(A_{\omega}) < \rho(A)$, it will be enough to find a pair of incoherent walks with respect to χ_{ω} , or equivalently, find a closed walk C of even length such that $\chi_{\omega}(C) \neq 1$. Let us inspect the closed walks containing edges in $\mathbf{e} \in \mathbf{E}(G) \setminus T$. For vertices v and v', denote by T(v, v')the unique path from v to v' on T. For $\mathbf{e} \in \mathbf{E}(G) \setminus T$, let $C_{\mathbf{e}}$ be the closed walk constituted of \mathbf{e} and $T(\mathbf{e}(1), \mathbf{e}(0))$. Let V_1, V_2 be as in Construction 4.2.

- (i) Let $\mathbf{E}' = \{\mathbf{e} \in \mathbf{E}(G) \setminus T \mid \{\mathbf{e}(0), \mathbf{e}(1)\} \cap V_1 \neq \emptyset$ or $\{\mathbf{e}(0), \mathbf{e}(1)\} \cap V_2 \neq \emptyset\}$ and $\mathbf{E}'' = \{\mathbf{e} \in \mathbf{E}(G) \setminus T \mid \{\mathbf{e}(0), \mathbf{e}(1)\} \subseteq V_1$ or $\{\mathbf{e}(0), \mathbf{e}(1)\} \subseteq V_2\}$. Then for $\omega \in \mathcal{X}_{(T,O)}$, we have $\omega \in \{0, \omega_{(T,O)}\}$ if and only if $\{\chi_{\omega}(\mathbf{e}) \mid \mathbf{e} \in \mathbf{E}'\} = \{1\}$ and $\{\chi_{\omega}(\mathbf{e}) \mid \mathbf{e} \in \mathbf{E}''\}$ is a singleton $\{1\}$ or $\{-1\}$.
- and only if $\{\chi_{\omega}(\mathbf{e}) \mid \mathbf{e} \in \mathbf{E}'\} = \{1\}$ and $\{\chi_{\omega}(\mathbf{e}) \mid \mathbf{e} \in \mathbf{E}''\}$ is a singleton $\{1\}$ or $\{-1\}$. (ii) For $\mathbf{e} \in \mathbf{E}(G) \setminus T$, note that $2\tau(C_{\mathbf{e}})$ is even and we have $\chi_{\omega}(C_{\mathbf{e}}^2) = \chi_{\omega}(C_{\mathbf{e}})^2 = \chi_{\omega}(\mathbf{e})^2 = 1$ if and only if $\chi_{\omega}(\mathbf{e}) = \pm 1$.

¹We note that Lemma 4.12 is recognized in the context of complex unit gain graphs (see Theorem 4.4 in [MKS22]), provided Lemma 4.4 and Lemma 4.7(a). Their proof was based on Perron-Frobenius theory, while a similar proof is not applicable to twisted edge adjacency matrices.

- (iii) For $\mathbf{e} \in \mathbf{E}'$, then the length of $T(\mathbf{e}(1), \mathbf{e}(0))$ is odd. Then $\tau(C_{\mathbf{e}}) = \tau(T(\mathbf{e}(1), \mathbf{e}(0))) + 1$ is even. Note that $\chi_{\omega}(C_{\mathbf{e}}) = \chi_{\omega}(\mathbf{e})$. Thus $\chi_{\omega}(C_{\mathbf{e}}) = 1$ if and only if $\chi_{\omega}(\mathbf{e}) = 1$.
- (iv) For $\mathbf{e}, \mathbf{e}' \in \mathbf{E}''$, consider a closed walk $C_{\mathbf{e},\mathbf{e}'}$ constituted of $\mathbf{e}, T(\mathbf{e}(1), \mathbf{e}'(0)), \mathbf{e}'$ and $T(\mathbf{e}'(1), \mathbf{e}(0))$. Since $\tau(T(\mathbf{e}(1), \mathbf{e}'(0)))$ and $T(\mathbf{e}'(1), \mathbf{e}(0))$ are both even or both odd, the length of $C_{\mathbf{e},\mathbf{e}'}$ must be even. Note that $\chi_{\omega}(C_{\mathbf{e},\mathbf{e}'}) = \chi_{\omega}(\mathbf{e})\chi_{\omega}(\mathbf{e}')$. Thus $\chi_{\omega}(C_{\mathbf{e},\mathbf{e}'}) = 1$ if and only if $\chi_{\omega}(\mathbf{e})\chi_{\omega}(\mathbf{e}') = 1$.

It follows from (i)-(iv) that if $\omega \in \mathcal{X}_{(T,O)} \setminus \{0, \omega_{(T,O)}\}$, we may also find a closed walk C of even length such that $\chi_{\omega}(C) \neq 1$, which means that $\rho(A_{\omega}) < \rho(A)$.

We say a graph G is *edge-walk-connected*² if for each pair of oriented edges $\mathbf{e}, \mathbf{e}' \in \mathbf{E}(G)$ (not necessarily distinct), there exists a non-backtracking edge-walk from \mathbf{e} to \mathbf{e}' .

Lemma 4.13. Consider an edge-walk-connected graph G and an arbitrary non-backtracking edgewalk Δ on G. There exists $a \in \mathbb{N}$ such that for each $\mathbf{e}, \mathbf{e}' \in \mathbf{E}(G)$ (not necessarily distinct) and each non-negative integers b, we have either

- (i) there exists no non-backtracking edge-walk from \mathbf{e} to \mathbf{e}' of length ax + b for all $x \in \mathbb{N}$; or
- (ii) there exists some $x_0 \in \mathbb{N}$ such that for all $x > x_0$, there exists a non-backtracking edge-walk from \mathbf{e} to \mathbf{e}' of length ax + b which contains $\boldsymbol{\Delta}$ as a sub-edge-walk.

Proof. Analogous to the proof of Lemma 4.10, we first claim that there exists $a \in \mathbb{N}$ such that for each $\mathbf{e} \in \mathbf{E}(G)$, there exists a non-backtracking edge-walk $\Upsilon_{\mathbf{e}}$ from \mathbf{e} to \mathbf{e} (equivalently this means that $\Upsilon_{\mathbf{e}}$ is a circuit at \mathbf{e}) of length a which contains $\boldsymbol{\Delta}$ as a sub-edge-walk.

For each $\mathbf{e} \in \mathbf{E}(G)$, let $\mathbf{\Upsilon}'_{\mathbf{e}}$ be a closed edge-walk at \mathbf{e} which is constituted of a non-backtracking edge-walk from \mathbf{e} to $\mathbf{\Delta}(0)$, the non-backtracking edge-walk $\mathbf{\Delta}$, and a non-backtracking edge-walk from $\mathbf{\Delta}(1)$ to \mathbf{e} . Then we may let a be a common multiple of the lengths of all $\mathbf{\Upsilon}'_{\mathbf{e}}$, and $a_{\mathbf{e}} = a/\tau(\mathbf{\Upsilon}'_{\mathbf{e}})$. As a result, $\mathbf{\Upsilon}_{\mathbf{e}} = (\mathbf{\Upsilon}'_{\mathbf{e}})^{a_{\mathbf{e}}}$ is a closed edge-walk at \mathbf{e} of length a which contains $\mathbf{\Delta}$ as a sub-edge-walk as desired.

Now for each $\mathbf{e}, \mathbf{e}' \in \mathbf{E}(G)$, suppose (i) is not satisfied, i.e., for some x_0 , there exists a non-backtracking edge-walk $\Upsilon'_{\mathbf{e},\mathbf{e}'}$ from \mathbf{e} to \mathbf{e}' of length $ax_0 + b$. Then for all $x > x_0$, the edge-walk $\Upsilon'_{\mathbf{e},\mathbf{e}'}$, followed by $\Upsilon'_{\mathbf{e},\mathbf{e}'}$ is a non-backtracking edge-walk from \mathbf{e} to \mathbf{e}' of length ax + b which contains Δ as a sub-edge-walk.

Clearly graphs of genus at most 1 cannot be edge-walk-connected. The following lemma says that a graph of genus at least 2 is "almost" edge-walk-connected.

Lemma 4.14. A graph G of genus at least 2 contains uniquely a subgraph of the same genus which is edge-walk-connected.

Proof. We perform repeatedly the following trimming process on G: whenever there is a degree 1 vertex, remove this vertex and the unique edge adjacent to the vertex. So when the process terminates, we derive a subgraph G' of G which has no degree 1 vertex and has the same genus as G. Note that each component of $G \setminus G'$ is a tree with an open end corresponding to the attaching vertex at G'. This construction guarantees the uniqueness of G', since if there is another G'' derived in this way, then $G \setminus G''$ must be disjoint from G' and $G \setminus G'$ must be disjoint from G''.

We claim that G' is edge-walk-connected. For a tree T which is a subgraph of G' and vertices $v, v' \in V(T)$, let T(v, v') denote the unique path on T from v to v'. For a graph G'' of genus at least 1 which is a subgraph of G', it is also clear that for any $v \in V(G'')$, there is a non-backtracking (not necessarily tailless) walk from v to v on G''. We use G''_v to denote the set of such walks.

Consider two oriented edges $\mathbf{e}, \mathbf{e}' \in \mathbf{E}(G')$ with e and e' being the underlying (non-oriented) edges respectively. First note that by our construction of G',

(1) if e is a cut-edge of G', then $G' \setminus e$ must have exactly two connected components G_1 and G_2 , each of genus at least 1;

²Equivalently, this means that the oriented line graph [KS00] associated to G is strongly connected.

- (2) if both e and e' are cut-edges of G', then $G' \setminus \{e, e'\}$ must have exactly three connected components G_1 , G_2 and G_3 such that G_1 and G_3 have genus at least 1, and both e and e' are in contact with G_2 .
- (3) if $\{e, e'\}$ is a cut of G' while neither e nor e' is a cut-edge of G', then $G' \setminus \{e, e'\}$ must have exactly two connected components G_1 and G_2 , while at least one of G_1 and G_2 has genus at least 1.

For each of the following cases, we will construct explicitly a non-backtracking edge-walk $\Delta_{\mathbf{e},\mathbf{e}'}$ from \mathbf{e} to \mathbf{e}' .

First, consider the case $\mathbf{e}' = \mathbf{e}$ or \mathbf{e}^{-1} .

- (a) If $G'' = G' \setminus e$ is connected, then G'' must have genus at least 1. We can let $\Delta_{\mathbf{e},\mathbf{e}}^{\times}$ (the interior of $\Delta_{\mathbf{e},\mathbf{e}}$) be $T(\mathbf{e}(1),\mathbf{e}(0))$ where T is a spanning tree of G'', and $\Delta_{\mathbf{e},\mathbf{e}^{-1}}^{\times}$ be any walk in $G''_{\mathbf{e}(1)}$.
- (b) If e is a cut-edge of G', then $G' \setminus e = G_1 \sqcup G_2$ where G_1 and G_2 are both connected with genus at least 1. Suppose $\mathbf{e}(0) \in G_1$ and $\mathbf{e}(1) \in G_2$, and let $\Delta \in (G_1)_{\mathbf{e}(0)}$ and $\Delta' \in (G_2)_{\mathbf{e}(1)}$. Then we can let $\mathbf{\Delta}_{\mathbf{e},\mathbf{e}}^{\times} = \Delta' \cdot \mathbf{e}^{-1} \cdot \Delta$ and $\mathbf{\Delta}_{\mathbf{e},\mathbf{e}^{-1}}^{\times} = \Delta'$.

Now suppose $\{\mathbf{e}, \mathbf{e}^{-1}\} \neq \{\mathbf{e}', \mathbf{e}'^{-1}\}.$

- (a) If $G'' = G' \setminus \{e, e'\}$ is connected, let T be a spanning tree of G'' and $\Delta_{\mathbf{e},\mathbf{e}'}^{\times} = T(\mathbf{e}(1),\mathbf{e}'(0)).$
- (b) If e is a cut-edge of G' such that $G' \setminus e = G_1 \sqcup G_2$ and e' is an edge of G_1 , suppose e' is not a cut-edge of G_1 and let T be a spanning tree of $G_1 \setminus e'$.
 - (i) If $\mathbf{e}(1)$ is a vertex of G_1 , then let $\mathbf{\Delta}_{\mathbf{e},\mathbf{e}'}^{\times} = T(\mathbf{e}(1),\mathbf{e}'(0))$.
 - (ii) If $\mathbf{e}(1)$ is a vertex of G_2 , then $\mathbf{e}(0)$ is a vertex of G_1 and let $\mathbf{\Delta}_{\mathbf{e},\mathbf{e}'}^{\times} = \mathbf{\Delta} \cdot \mathbf{e}^{-1} \cdot T(\mathbf{e}(0),\mathbf{e}'(0))$ for some $\mathbf{\Delta} \in (G_2)_{\mathbf{e}(1)}$.
- (c) If both e and e' are cut-edges of G', then $G' \setminus \{e, e'\} = G_1 \sqcup G_2 \sqcup G_3$ with G_1 , G_2 and G_3 as in Case (2) stated above. Without loss of generality, suppose e is in contact with G_1 and e' is in contact with G_3 . Let T be a spanning tree of G_2 .
 - (i) If $\mathbf{e}(1), \mathbf{e}'(0) \in V(G_2)$, let $\Delta_{\mathbf{e},\mathbf{e}'}^{\times} = T(\mathbf{e}(1), \mathbf{e}'(0))$.
 - (ii) If $\mathbf{e}(1), \mathbf{e}'(1) \in V(G_2)$, then $\mathbf{e}'(0)$ is a vertex of G_3 and let $\mathbf{\Delta}_{\mathbf{e},\mathbf{e}'}^{\times} = T(\mathbf{e}(1), \mathbf{e}'(1)) \cdot \mathbf{e}'^{-1} \cdot \Delta$ for some $\Delta \in (G_3)_{\mathbf{e}'(0)}$.
- (iii) If $\mathbf{e}(0), \mathbf{e}'(0) \in V(G_2)$, then $\mathbf{e}(1)$ is a vertex of G_1 and let $\mathbf{\Delta}_{\mathbf{e},\mathbf{e}'}^{\times} = \mathbf{\Delta} \cdot \mathbf{e}^{-1} \cdot T(\mathbf{e}(0), \mathbf{e}'(0))$ for some $\mathbf{\Delta} \in (G_1)_{\mathbf{e}(1)}$.
- (iv) If $\mathbf{e}(0), \mathbf{e}'(1) \in V(G_2)$, then $\mathbf{e}(1)$ is a vertex of G_1 and $\mathbf{e}'(0)$ is a vertex of G_3 . Let $\mathbf{\Delta}_{\mathbf{e},\mathbf{e}'}^{\times} = \Delta \cdot \mathbf{e}^{-1} \cdot T(\mathbf{e}(0), \mathbf{e}'(1)) \cdot \mathbf{e}'^{-1} \cdot \Delta'$ for some $\Delta \in (G_1)_{\mathbf{e}(1)}$ and $\Delta' \in (G_3)_{\mathbf{e}'(0)}$.
- (d) If $\{e, e'\}$ is a cut of G' while neither e nor e' is a cut-edge of G', then $G' \setminus \{e, e'\}$ has exactly two connected components G_1 and G_2 . By Case (3) stated above, we may assume G_1 has genus at least 1. Let T_1 be a spanning tree of G_1 and T_2 be a spanning tree of G_2 .
 - (i) If $\mathbf{e}(1), \mathbf{e}'(0) \in V(G_1)$, let $\mathbf{\Delta}_{\mathbf{e},\mathbf{e}'}^{\times} = T_1(\mathbf{e}(1), \mathbf{e}'(0))$.
 - (ii) If $\mathbf{e}(1), \mathbf{e}'(1) \in V(G_1)$, let $\mathbf{\Delta}_{\mathbf{e},\mathbf{e}'}^{\times} = \mathbf{\Delta} \cdot \mathbf{e}^{-1} \cdot T_2(\mathbf{e}(0), \mathbf{e}'(0))$ for some $\mathbf{\Delta} \in (G_1)_{\mathbf{e}(1)}$.
 - (iii) If $\mathbf{e}(0), \mathbf{e}'(0) \in V(G_1)$, let $\mathbf{\Delta}_{\mathbf{e},\mathbf{e}'}^{\times} = T_2(\mathbf{e}(1), \mathbf{e}'(1)) \cdot \mathbf{e}'^{-1} \cdot \Delta$ for some $\Delta \in (G_1)_{\mathbf{e}'(0)}$.
 - (iv) If $\mathbf{e}(0), \mathbf{e}'(1) \in V(G_1)$, let $\mathbf{\Delta}_{\mathbf{e},\mathbf{e}'}^{\times} = T_2(\mathbf{e}(1), \mathbf{e}'(0))$.

We say two edge-walks Δ and Δ' of the same length with the same initial oriented edge and the same terminal oriented edge are *coherent* with respect to χ_{ω} if $\chi_{\omega}(\Delta) = \chi_{\omega}(\Delta')$, and *incoherent* with respect to χ_{ω} if $\chi_{\omega}(\Delta) \neq \chi_{\omega}(\Delta')$.

Lemma 4.15. $\rho(W_{1,\omega}) \leq \rho(W_1)$ for all $\omega \in \Omega$. In addition, $\rho(W_{1,\omega}) < \rho(W_1)$ if and only if there exist a pair of incoherent non-backtracking edge-walks with respect to χ_{ω} .

Proof. Analogous to the proof of Lemma 4.11, we will apply Gelfand's formula to $W_{1,\omega}$. Again, using the ∞ -norm of $W_{1,\omega}^k$,

$$\|W_{1,\omega}^k\| := \max_{\mathbf{e}\in\mathbf{E}(G)} \sum_{\mathbf{e}'\in\mathbf{E}(G)} |(W_{1,\omega}^k)_{\mathbf{e}\mathbf{e}'}|,$$

the spectral radius of $W_{1,\omega}$ can be expressed as follows:

$$\rho(W_{1,\omega}) = \lim_{k \to \infty} \|W_{1,\omega}^k\|^{1/k}$$
$$= \lim_{k \to \infty} \left(\max_{\mathbf{e} \in \mathbf{E}(G)} \sum_{\mathbf{e}' \in \mathbf{E}(G)} |(W_{1,\omega}^k)_{\mathbf{e}\mathbf{e}'}| \right)^{1/k}.$$

Let $\Gamma(\mathbf{e}, \mathbf{e}')$ and $\Gamma_k(\mathbf{e}, \mathbf{e}')$ be the set of all non-backtracking edge-walks and the set of all length-k non-backtracking edge-walks from \mathbf{e} to \mathbf{e}' respectively. Then

$$(W_{1,\omega}^k)_{\mathbf{e}\mathbf{e}'} = \sum_{\mathbf{\Delta}\in\Gamma_k(\mathbf{e},\mathbf{e}')} \chi_{\omega}(\mathbf{\Delta}).$$

Clearly $|(W_{1,\omega}^k)_{\mathbf{ee'}}| \leq \#\Gamma_k(\mathbf{e}, \mathbf{e'}) = (W_1^k)_{\mathbf{ee'}}$. Consequently $\rho(W_{1,\omega}) \leq \rho(W_1)$ for all $\omega \in \Omega$. Also, if all non-backtracking edge-walks in $\Gamma_k(\mathbf{e}, \mathbf{e'})$ are coherent with respect to χ_ω , then $|(W_{1,\omega}^k)_{\mathbf{ee'}}| = \#\Gamma_k(\mathbf{e}, \mathbf{e'}) = (W_1^k)_{\mathbf{ee'}}$. Again, by Gelfand's formula, this means that $\rho(W_{1,\omega}) = \rho(W_1)$ if there are no pairs of incoherent non-backtracking edge-walks with respect to χ_ω .

Conversely, suppose there are a pair of incoherent non-backtracking edge-walks $\Delta, \Delta' \in \Gamma_k(\mathbf{a}, \mathbf{b})$ for some $\mathbf{a}, \mathbf{b} \in \mathbf{E}(G)$. Then by Lemma 4.13, there exists $a \in \mathbb{N}$ such that for all $\mathbf{e}, \mathbf{e}' \in \mathbf{E}(G)$, we have either

- (i) $\Gamma_{ax}(\mathbf{e}, \mathbf{e}') = \emptyset$ for all $x \in \mathbb{N}$, or
- (ii) there exists a non-backtracking edge-walk $\Upsilon \in \Gamma_{ax}(\mathbf{e}, \mathbf{e}')$ containing Δ as a sub-edge-walk for all x large enough.

Analogous to the proof of Lemma 4.11, Case (i) means that $(W_{1,\omega}^{ax})_{\mathbf{ee}'} = (W_1^{ax})_{\mathbf{ee}'} = 0$ for all $x \in \mathbb{N}$, and Case (ii) means that for all x large enough, we can always get another non-backtracking edge-walk $\mathbf{\Upsilon}' \in \Gamma_{ax}(\mathbf{e}, \mathbf{e}')$ incoherent to $\mathbf{\Upsilon}$ by replacing a sub-edge-walk $\mathbf{\Delta}$ in $\mathbf{\Upsilon}$ by $\mathbf{\Delta}'$ and leaving the remaining part unchanged, which further implies that $|(W_{1,\omega}^{ax})_{\mathbf{ee}'}| < (W_1^{ax})_{\mathbf{ee}'}$.

Conclusively, there exists 0 < c < 1 and a large enough $y \in \mathbb{N}$ such that $|(W_{1,\omega}^y)_{ee'}| \leq c(W_1^y)_{ee'}$ for all $\mathbf{e}, \mathbf{e}' \in \mathbf{E}(G)$.

Therefore,
$$|(W_{1,\omega}^{ky})_{ee'}| \le c^k (W_1^{ky})_{ee'}$$
 and $||W_{1,\omega}^{ky}|| = \max_{e \in \mathbf{E}(G)} \sum_{e' \in \mathbf{E}(G)} |(W_{1,\omega}^{ky})_{ee'}| \le c^k ||W_1^{ky}||.$

Finally, we have

$$\rho(W_{1,\omega}) = \lim_{k \to \infty} \|W_{1,\omega}^{ky}\|^{1/(ky)} \le \lim_{k \to \infty} \left(c^k \|W_1^{ky}\|\right)^{1/(ky)} = c^{1/y} \rho(W_1) < \rho(W_1).$$

1//1

Lemma 4.16. If g = 1, then $\rho(W_{1,\omega}) = 1$ for all $\omega \in \Omega$.

Proof. If g = 1, then G must contain uniquely a subgraph G' which is a cycle graph with exactly two primes P and P^{-1} . Suppose G' has n' vertices. Then $\tau(P) = \tau(P^{-1}) = n'$, and for each $\omega \in \Omega$,

$$L_G(u, \chi_{\omega})^{-1} = L_{G'}(u, \chi_{\omega})^{-1}$$

= $\left(1 - \chi_{\omega}(P)u^{n'}\right) \left(1 - \chi_{\omega}(P^{-1})u^{n'}\right).$



FIGURE 5. Cases of two closed paths C_a and C_b , containing respectively the orientated edges \mathbf{e}_a and \mathbf{e}_b in $\mathbf{E}(G) \setminus T$: (a) C_a and C_b have an overlapping sub-path; (b) C_a and C_b are in contact at a single vertex; (c) C_a and C_b are disjoint.

By Theorem 2.26(a), the characteristic polynomial of $W_{1,\omega}$ is

$$p_{\omega}(\lambda) = \lambda^{2m-2n'} \left(\lambda^{n'} - \chi_{\omega}(P) \right) \left(\lambda^{n'} - \chi_{\omega}(P^{-1}) \right)$$

Then $\rho(W_{1,\omega}) = 1$ for all $\omega \in \Omega$, since $|\chi_{\omega}(P)| = |\chi_{\omega}(P^{-1})| = 1$.

Lemma 4.17. If $g \geq 2$, then

- (i) if G is non-bipartite, then $\rho(W_{1,\omega}) < \rho(W_1)$ for all $\omega \in \mathcal{X}_{(T,O)} \setminus \{0, \omega_{(T,O)}\}$; and
- (ii) if G is bipartite, then $\rho(W_{1,\omega}) < \rho(W_1)$ for all $\omega \in \mathcal{X}_{(T,O)}$ such that $\omega \notin \{\sum_{i=1}^g \omega_i d\mathbf{e}_i \mid \omega_i \in \{0, 1/2\}\}$ where $\{\mathbf{e}_1, \cdots, \mathbf{e}_q\} = \mathbf{E}_O(G) \setminus T$.

Proof. By Lemma 4.14, when $g \ge 2$, G must contain uniquely a subgraph G', also of genus g, which is edge-walk-connected. By definition, it is clear that G and G' have the same L-function for the same character χ_{ω} , i.e., $L_G(u, \chi_{\omega}) = L_{G'}(u, \chi_{\omega})$. Then by Theorem 2.26(a), the edge adjacency matrices twisted by χ_{ω} of G and G' must have the same spectral radius. Thus for further discussions here, we may assume G itself is edge-walk-connected.

As in the proof of Lemma 4.12, let T(v, v') denote the unique path from v to v' contained in T for $v, v' \in V(G)$, and we will inspect the closed paths $C_{\mathbf{e}}$ constituted of $\mathbf{e} \in \mathbf{E}(G) \setminus T$ and $T(\mathbf{e}(1), \mathbf{e}(0))$. Actually here we will consider pairs of $\mathbf{e}_a, \mathbf{e}_b \in \mathbf{E}(G) \setminus T$ such that $\{\mathbf{e}_a, \mathbf{e}_a^{-1}\} \neq \{\mathbf{e}_b, \mathbf{e}_b^{-1}\}$ (existence guaranteed by the assumption of $g \geq 2$). For simplicity of notation, we write $T_a = T(\mathbf{e}_a(1), \mathbf{e}_a(0))$, $T_b = T(\mathbf{e}_b(1), \mathbf{e}_b(0)), C_a = C_{\mathbf{e}_a}$ and $C_b = C_{\mathbf{e}_b}$. As shown in Figure 5, there are three possible cases based on the topological relations among C_a, C_b and T.

Case (a): There is a maximal overlapping path T(v, w) between C_a and C_b with vertices v and w in both T_a and T_b .

Case (b): C_a and C_b are in contact at exactly one vertex v.

Case (c): C_a and C_b are disjoint, and in this case there exists a minimal path T(v, w) from a vertex v in T_a to vertex w in T_b .

Note that here for convenience of discussion, switching \mathbf{e} and \mathbf{e}^{-1} if necessary for $\mathbf{e} = \mathbf{e}_a$ or \mathbf{e}_b , the orientations of \mathbf{e}_a , \mathbf{e}_b , C_a and C_b are chosen as in Figure 5.

Again, for V_1 and V_2 defined in Construction 4.2, we let $\mathbf{E}' = \{\mathbf{e} \in \mathbf{E}(G) \setminus T \mid \{\mathbf{e}(0), \mathbf{e}(1)\} \cap V_1 \neq \emptyset$ or $\{\mathbf{e}(0), \mathbf{e}(1)\} \cap V_2 \neq \emptyset\}$ and $\mathbf{E}'' = \{\mathbf{e} \in \mathbf{E}(G) \setminus T \mid \{\mathbf{e}(0), \mathbf{e}(1)\} \subseteq V_1$ or $\{\mathbf{e}(0), \mathbf{e}(1)\} \subseteq V_2\}$. Then

for $\omega \in \mathcal{X}_{(T,O)}$, we have $\omega \in \{0, \omega_{(T,O)}\}$ if and only if $\{\chi_{\omega}(\mathbf{e}) \mid \mathbf{e} \in \mathbf{E}'\} = \{1\}$ and $\{\chi_{\omega}(\mathbf{e}) \mid \mathbf{e} \in \mathbf{E}''\}$ is a singleton $\{1\}$ or $\{-1\}$. Now based on the parity of the length of C_a and C_b , we have four cases: *Case* (1): $\tau(C_a)$ is odd and $\tau(C_b)$ is odd. Equivalently, $\mathbf{e}_a, \mathbf{e}_b \in \mathbf{E}''$.

- Case (2): $\tau(C_a)$ is even and $\tau(C_b)$ is odd. Equivalently, $\mathbf{e}_a \in \mathbf{E}'$ and $\mathbf{e}_b \in \mathbf{E}''$.
- Case (3): $\tau(C_a)$ is odd and $\tau(C_b)$ is even. Equivalently, $\mathbf{e}_a \in \mathbf{E}''$ and $\mathbf{e}_b \in \mathbf{E}'$.
- Case (4): $\tau(C_a)$ is even and $\tau(C_b)$ is even. Equivalently, $\mathbf{e}_a, \mathbf{e}_b \in \mathbf{E}'$.

By Lemma 4.6, we see that $\rho(W_{1,\omega_{(T,O)}}) = \rho(W_1)$. Now consider $W_{1,\omega}$ for $\omega \in \mathcal{X}_{(T,O)}$ and assume that $\rho(W_{1,\omega}) = \rho(W_1)$, which means that any pair of non-backtracking edge-walks of the same length with the same initial edge and the same terminal edge are coherent by Lemma 4.15.

First, we will show that $\chi_{\omega}(\mathbf{e}) = \pm 1$ for all $\mathbf{e} \in \mathbf{E}(G) \setminus T$. Applied to all the three cases in Figure 5, consider two non-backtracking edge-walks $\boldsymbol{\Delta}$ and $\boldsymbol{\Delta}'$, both from \mathbf{e}_a to \mathbf{e}_a^{-1} , whose interior paths are the path $T(\mathbf{e}_a(1), \mathbf{e}_b(0)) \cdot \mathbf{e}_b \cdot T(\mathbf{e}_b(1), \mathbf{e}_a(0))$ and the path $T(\mathbf{e}_a(1), \mathbf{e}_b(1)) \cdot \mathbf{e}_b^{-1} \cdot T(\mathbf{e}_b(0), \mathbf{e}_a(0))$ respectively. Then it is clear that $\boldsymbol{\Delta}$ and $\boldsymbol{\Delta}'$ are of the same length. In addition, $\chi_{\omega}(\boldsymbol{\Delta}) = \chi_{\omega}(\mathbf{e}_b)$ and $\chi_{\omega}(\boldsymbol{\Delta}') = \chi_{\omega}(\mathbf{e}_b^{-1}) = \overline{\chi_{\omega}(\mathbf{e}_b)}$. Since $\boldsymbol{\Delta}$ and $\boldsymbol{\Delta}'$ are coherent by assumption, we must have $\chi_{\omega}(\boldsymbol{\Delta}) = \chi_{\omega}(\boldsymbol{\Delta}')$, which implies that $\chi_{\omega}(\mathbf{e}_b) = \overline{\chi_{\omega}(\mathbf{e}_b)} = \pm 1$. This argument applies to all $\mathbf{e} \in \mathbf{E}(G) \setminus T$. Therefore, we can conclude that $\omega \in \{\sum_{i=1}^g \omega_i d\mathbf{e}_i \mid \omega_i \in \{0, 1/2\}\}$ where $\{\mathbf{e}_1, \dots, \mathbf{e}_g\} = \mathbf{E}_O(G) \setminus T$, and we've proved (ii).

Now we will let G be a non-bipartite graph and further refine the conclusion as stated in (i). In particular, we will construct closed edge-walks from \mathbf{e}_a to \mathbf{e}_a , or equivalently circuits going through \mathbf{e}_a , using two strategies:

Strategy (I): Let $\Delta_k := (\mathbf{e}_a \cdot T(\mathbf{e}_a(1), \mathbf{e}_a(0)))^k$, i.e., a circuit constructed by winding k times around C_a . Then $\tau(\Delta_k) = k \cdot \tau(C_a)$.

Strategy (II): Let $\Delta'_{k'} := \left(\mathbf{e}_a \cdot T(\mathbf{e}_a(1), \mathbf{e}_b(1)) \cdot \mathbf{e}_b^{-1} \cdot T(\mathbf{e}_b(0), \mathbf{e}_a(0))\right)^{k'}$. Depending on the cases in Figure 5, we have $\tau(\Delta'_{k'}) = k' \tau_{ab}$ where

$$\tau_{ab} = \begin{cases} \tau(C_a) + \tau(C_b) - 2\tau(T(v, w)), & \text{for Case (a),} \\ \tau(C_a) + \tau(C_b), & \text{for Case (b),} \\ \tau(C_a) + \tau(C_b) + 2\tau(T(v, w)), & \text{for Case (c).} \end{cases}$$

Now we want to make Δ_k and $\Delta'_{k'}$ a coherent pair. Then we need to choose proper winding numbers k and k' such that $\tau(\Delta_k) = k\tau(C_a) = \tau(\Delta'_{k'}) = k'\tau_{ab}$. Thus we may let $k = \tau_{ab}/\gcd(\tau(C_a), \tau_{ab})$ and $k' = \tau(C_a)/\gcd(\tau(C_a), \tau_{ab})$. Moreover, since Δ_k and $\Delta'_{k'}$ are coherent, we must have $\chi_{\omega}(\Delta_k) = \chi_{\omega}(\Delta'_{k'})$. This actually means that $\chi_{\omega}(\mathbf{e}_a)^k = \chi_{\omega}(\mathbf{e}_a)^{k'}\chi_{\omega}(\mathbf{e}_b^{-1})^{k'} = \chi_{\omega}(\mathbf{e}_a)^{k'}\chi_{\omega}(\mathbf{e}_b)^{k'}$. Note that the second equality follows from $\chi_{\omega}(\mathbf{e}_a) = \chi_{\omega}(\mathbf{e}_a^{-1}) = \chi_{\omega}(\mathbf{e}_b) = \chi_{\omega}(\mathbf{e}_b^{-1}) = \pm 1$. Thus, based on the cases of parity of the length of C_a and C_b , we have further conclusions as follows.

Case (1): In this case, $\tau(C_a)$ is odd and $\tau(C_b)$ is odd. Then τ_{ab} is even (for all cases of (a), (b) and (c)), k is even and k' is odd, which means that $\chi_{\omega}(\mathbf{e}_a) = \chi_{\omega}(\mathbf{e}_b)$ (by the equality $\chi_{\omega}(\mathbf{e}_a)^k = \chi_{\omega}(\mathbf{e}_a)^{k'}\chi_{\omega}(\mathbf{e}_b)^{k'}$).

Case (2): In this case, $\tau(C_a)$ is even and $\tau(C_b)$ is odd. Then τ_{ab} is odd, k is odd and k' is even, which means $\chi_{\omega}(\mathbf{e}_a) = 1$.

Case (3): In this case, $\tau(C_a)$ is odd and $\tau(C_b)$ is even. Then τ_{ab} is odd, k is odd and k' is odd, which means $\chi_{\omega}(\mathbf{e}_b) = 1$.

Case (4): In this case, $\tau(C_a)$ is even and $\tau(C_b)$ is even. Then τ_{ab} is even, k and k' can be either even or odd.

Note that here by assumption, G is non-bipartite, which means \mathbf{E}'' is nonempty. Therefore, the above conclusion for Case (1) guarantees that $\{\chi_{\omega}(\mathbf{e}) \mid \mathbf{e} \in \mathbf{E}''\}$ is a singleton $\{1\}$ or $\{-1\}$, the conclusions for Case (2) and (3) guarantee that $\{\chi_{\omega}(\mathbf{e}) \mid \mathbf{e} \in \mathbf{E}'\} = \{1\}$. As a result, we must have $\omega \in \{0, \omega_{(T,O)}\}$ as claimed.

Now we can get a complete proof of Theorem 3.1.

39

Proof of Theorem 3.1. For (a), as in Lemma 4.4, we let the canonical character be $\theta = \underline{\omega_O} = \underline{\omega_{(T,O)}}$. Then by Lemma 2.20, Lemma 4.5, Lemma 4.6 and Lemma 4.7, a generalization can be made as for all $\omega, \omega' \in \Omega$ such that $\underline{\omega} + \underline{\omega'} = \theta$, $A_{\omega'}$ is similar to $-A_{\omega}$, and Spec $W_{1,\omega'} = -$ Spec $W_{1,\omega}$.

To show such a $\theta \in \mathcal{X}$ is unique, suppose that there exists $\theta' = \underline{\omega}' \in \mathcal{X}$ also satisfying the above property. Then Spec $W_{1,\omega'} = -$ Spec W_1 . Let $\omega = \omega_{(T,O)} - \omega'$ which means $\underline{\omega} = \theta - \theta'$. We must have Spec $W_{1,\omega} = -$ Spec $W_{1,\omega'} =$ Spec W_1 . Then we must have $\theta - \theta' = 0$ by Lemma 4.8.

For (b), by our construction, it is clear that θ is a 2-torsion in \mathcal{X} .

For (c), by the construction in Construction 4.2, we conclude that $\omega_{(T,O)} = 0$ if and only if G is bipartite. Therefore, $\theta = \omega_{(T,O)} = 0$ and by Lemma 2.20, Spec $A_{\omega} = -$ Spec $A_{-\omega} = -$ Spec A_{ω} and Spec $W_{1,\omega} = -$ Spec $W_{1,-\omega} = -$ Spec $W_{1,\omega}$.

Finally, (d)-(h) follow from (a), (b), Lemma 4.11, Lemma 4.12, Lemma 4.15, Lemma 4.16 and Lemma 4.17 respectively.

References

- [ABKS14] Yang An, Matthew Baker, Greg Kuperberg, and Farbod Shokrieh. Canonical representatives for divisor classes on tropical curves and the matrix-tree theorem. In *Forum of Mathematics, Sigma*, volume 2, page e24. Cambridge University Press, 2014.
- [AS87a] Toshiaki Adachi and Toshikazu Sunada. Homology of closed geodesics in a negatively curved manifold. J. Differ. Geom., 26:81–99, 1987.
- [AS87b] Toshiaki Adachi and Toshikazu Sunada. Twisted Perron-Frobenius theorem and L-functions. J. Funct. Anal., 71:1–46, 1987.
- [Bas92] Hyman Bass. The Ihara-Selberg zeta function of a tree lattice. Internat. J. Math., 3(6):717–797, 1992.
- [BF11] Matthew Baker and Xander Faber. Metric properties of the tropical Abel-Jacobi map. J. Algebr. Comb., 33(3):349–381, 2011.
- [BH12] Andries E. Brouwer and Willem H. Haemers. Spectra of graphs. Universitext. Berlin: Springer, 2012.
- [BN07] Matthew Baker and Serguei Norine. Riemann-roch and abel-jacobi theory on a finite graph. Advances in Mathematics, 215(2):766–788, 2007.
- [Chu97] Fan R. K. Chung. Spectral graph theory, volume 92 of Reg. Conf. Ser. Math. Providence, RI: AMS, American Mathematical Society, 1997.
- [CRS10] Dragoš Cvetković, Peter Rowlinson, and Slobodan Simić. An introduction to the theory of graph spectra, volume 75 of Lond. Math. Soc. Stud. Texts. Cambridge: Cambridge University Press, 2010.
- [Del42] Jean Delsarte. Sur le gitter fuchsien. C. R. Acad. Sci., Paris, 214:147–149, 1942.
- [Has89] Ki-ichiro Hashimoto. Zeta functions of finite graphs and representations of p-adic groups. In Automorphic forms and geometry of arithmetic varieties, volume 15 of Adv. Stud. Pure Math., pages 211–280. Academic Press, Boston, MA, 1989.
- [Has92] Ki-ichiro Hashimoto. Artin type *L*-functions and the density theorem for prime cycles on finite graphs. Internat. J. Math., 3(6):809–826, 1992.
- [Hej76] Dennis A. Hejhal. The Selberg trace formula and the Riemann zeta-function. Duke Math. J., 43:441–482, 1976.
- [HFGO99] Dennis A. Hejhal, Joel Friedman, Martin C. Gutzwiller, and Andrew M. Odlyzko, editors. Emerging applications of number theory, volume 109 of The IMA Volumes in Mathematics and its Applications. Springer-Verlag, New York, 1999. Papers from the IMA Summer Program held at the University of Minnesota, Minneapolis, MN, July 15–26, 1996.
- [HS15] Takehiro Hasegawa and Seiken Saito. A generalization of the graph theory prime-number theorem of a finite graph. *Internat. J. Math.*, 26(9):1550071, 19, 2015.
- [HST06] Matthew D. Horton, H. M. Stark, and Audrey A. Terras. What are zeta functions of graphs and what are they good for? In *Quantum graphs and their applications*, volume 415 of *Contemp. Math.*, pages 173–189. Amer. Math. Soc., Providence, RI, 2006.
- [Hub61] Heinz Huber. Zur analytischen Theorie hyperbolischer Raumformen und Bewegungsgruppen. II. Math. Ann., 142:385–398, 1961.
- [Iha66] Yasutaka Ihara. On discrete subgroups of the two by two projective linear group over p-adic fields. J. Math. Soc. Japan, 18:219–235, 1966.
- [Iwa84] Henryk Iwaniec. Prime geodesic theorem. J. Reine Angew. Math., 349:136–159, 1984.
- [KS87] Atsushi Katsuda and Toshikazu Sunada. Homology and closed geodesics in a compact Riemann surface. Am. J. Math., 110(1):145–155, 1987.

SPECTRAL	ANTISYMMETRY	OF TWISTED	GRAPH	ADJACENCY
----------	--------------	------------	-------	-----------

- [KS00] Motoko Kotani and Toshikazu Sunada. Zeta functions of finite graphs. J. Math. Sci., Tokyo, 7(1):7–25, 2000.
- [Lal89] Steven P. Lalley. Closed geodesics in homology classes on surfaces of variable negative curvature. Duke Math. J., 58(3):795–821, 1989.
- [LS95] Wenzhi Luo and Peter Sarnak. Quantum ergodicity of eigenfunctions on $PSL_2(\mathbb{Z})\backslash H^2$. Publ. Math., Inst. Hautes Étud. Sci., 81:207–237, 1995.
- [MKS22] Ranjit Mehatari, M. Rajesh Kannan, and Aniruddha Samanta. On the adjacency matrix of a complex unit gain graph. *Linear Multilinear Algebra*, 70(9):1798–1813, 2022.
- [Pol91] Mark Pollicott. Homology and closed geodesics in a compact negatively curved surface. Am. J. Math., 113(3):379–385, 1991.
- [PS87] Ralph Phillips and Peter Sarnak. Geodesics in homology classes. Duke Math. J., 55:287–297, 1987.
- [Ref12] Nathan Reff. Spectral properties of complex unit gain graphs. *Linear Algebra Appl.*, 436(9):3165–3176, 2012.
- [Sar82] Peter Sarnak. Class numbers of indefinite binary quadratic forms. Journal of Number Theory, 15(2):229– 247, 1982.
- [Ser80] Jean-Pierre Serre. Trees. Springer Berlin Heidelberg, 1980.
- [ST96] H. M. Stark and A. A. Terras. Zeta functions of finite graphs and coverings. Adv. Math., 121(1):124–165, 1996.
- [ST00] H. M. Stark and A. A. Terras. Zeta functions of finite graphs and coverings. II. Adv. Math., 154(1):132–195, 2000.
- [SY13] K. Soundararajan and Matthew P. Young. The prime geodesic theorem. J. Reine Angew. Math., 676:105– 120, 2013.
- [Ter11] Audrey Terras. Zeta functions of graphs, volume 128 of Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 2011. A stroll through the garden.
- [TS07] A. A. Terras and H. M. Stark. Zeta functions of finite graphs and coverings. III. Adv. Math., 208(1):467– 489, 2007.
- [Zas89] Thomas Zaslavsky. Biased graphs. I: Bias, balance, and gains. J. Comb. Theory, Ser. B, 47(1):32–52, 1989.

40

APPENDIX A. NOTATION GUIDE

Section 1.	
G	a connected finite graph without multiple edges or loops
g	the genus (first Bettie number) of G
V(G)	the vertex set of G
E(G)	the edge set of G
$\mathbf{E}(G)$	the set of all oriented edges of G
$\mathbf{E}_O(G)$	the set of all positively oriented edges with respect to an orientation Q of G
$\mathbf{a}(0)$ and $\mathbf{a}(1)$	the initial and terminal vertices of an oriented edge a
$\Delta(0)$ and $\Delta(1)$	the initial and terminal vertices of a walk Λ
$\Delta(0)$ and $\Delta(1)$ $\Delta(0)$ and $\Delta(1)$	the initial and terminal oriented edges of an edge-walk Λ
$\tau(\Delta)$ and $\pi(1)$	the length of Λ
$z(u)$ or $z_{\alpha}(u)$	the Ibara zeta function of G
$\mathcal{T}(G)$ or \mathcal{T}	the tangent space of G (elements being vector fields)
$\Omega(G)$ or Ω	the cotangent space of G (elements being 1-forms)
Λ^{ab}	the abelianization of a walk Λ
	the unit character associated to a 1-form ω
$L(u, \chi)$ or $L_{\alpha}(u, \chi)$) the L-function with respect to a character χ
$L(u, \chi_{\omega})$ of $LG(u, \chi_{\omega})$	ω) the n-random with respect to a character $\chi\omega$
Section 2. $\mathcal{U}^{1}(\mathcal{C})$	the space of harmonic 1 forms on C
π (G) Im(d)	the space of marmonic 1-forms on C
$\lim_{a \to a} (a) + \phi(a)$	the space of exact 1-forms of G a 1 form (1) with $\phi(x) \in \mathcal{U}^{1}(C)$
$\omega = \phi_1(\omega) + \phi_2(\omega)$	orthogonal decomposition of a 1-iorin ω with $\varphi_1(\omega) \in \pi$ (G)
$U(C \mathbb{D})$	and $\omega_2 \in \operatorname{III}(a)$
$\Pi_1(G,\mathbb{K})$ $\Pi_1(G,\mathbb{Z})$	the 1st real homomory group of G
$\pi_1(G,\mathbb{Z})$	the 1st integral homomoly group of G, which is a full-rank lattice in $H_1(G, \mathbb{R})$
$H_1(G,\mathbb{Z})^{\vee}$	the dual of $H_1(G,\mathbb{Z})$ which is a full rank lattice in $\mathcal{H}^1(G)$
$e(\cdot)$	$e(x) := \exp\left(2\pi\sqrt{-1}x\right)$
$\mathcal{X}(G)$ or \mathcal{X}	$\mathcal{H}^1(G)/H_1(G,\mathbb{Z})^{\vee}$, the character group of G
ω	$\phi_1(\omega) + H_1(G,\mathbb{Z})^{\vee}$, the projection of a 1-form ω to \mathcal{X}
$\overline{\Lambda}$	a full-rank sublattice of $H_1(G,\mathbb{Z})$
Λ^{ee}	the dual lattice of Λ
\mathcal{X}_{Λ}	$\mathcal{H}^1(G)/\Lambda^{ee}$
$\underline{\omega}$	$\phi_1(\omega) + \Lambda^{\vee}$, the projection of a 1-form ω to \mathcal{X}_{Λ}
\overline{Q}_{Λ}	$H_1(G,\mathbb{Z})/\Lambda$
$\underline{\alpha}$	$\alpha + \Lambda \in Q_{\Lambda}$ for $\alpha \in H_1(G, \mathbb{Z})$
$\widehat{Q_{\Lambda}}$	$\Lambda^{\vee}/H_1(G,\mathbb{Z})^{\vee}$
Ă	the adjacency matrix of G
W_1	the edge adjacency matrix of G
A_{ω}	the adjacency matrix of G twisted by χ_{ω}
$\tilde{W_{1,\omega}}$	the edge adjacency matrix of G twisted by χ_{ω}
$W'_{1,i}$ and $W''_{1,i}$	variants of $W_{1,\omega}$
$\operatorname{Spec}^{1,\omega} M$	the spectrum of a square matrix M
*	

Section 3.

- $\rho(M)$ the spectral radius of a square matrix M
- θ the canonical character of G
- N(l) the number of circuits of length l
- $N(\alpha, l)$ the number of circuits C of length l such that $C^{ab} = \alpha \in H_1(G, \mathbb{Z})$
- $N(\underline{\alpha}, l)$ the number of circuits C of length l such that $\underline{C}^{ab} = \underline{\alpha} \in Q_{\Lambda}$
- $\mathcal{K}(\omega, l)$ the trace distribution function of order l
- $\pi(l)$ the number of prime cycles of length l
- $\pi_c(l)$ the number of cycles of length l
- $\pi(\alpha, l)$ the number of prime cycles P of length l such that $P^{ab} = \alpha \in H_1(G, \mathbb{Z})$
- $\pi_c(\alpha, l)$ the number of cycles C of length l such that $C^{ab} = \alpha \in H_1(G, \mathbb{Z})$
- $\pi(\underline{\alpha}, l)$ the number of prime cycles P of length l such that $\underline{P^{ab}} = \underline{\alpha} \in Q_{\Lambda}$
- $\pi_c(\underline{\alpha}, l)$ the number of cycles C of length l such that $\underline{C^{ab}} = \underline{\alpha} \in Q_{\Lambda}$

Section 4.

- ω_O a canonical 1-form with respect to an orientation O, which is the output of Construction 4.1
- $\omega_{(T,O)}$ a canonical 1-form with respect to an orientation O and a spanning tree T, which is the output of Construction 4.2
- $\mathcal{X}_{(T,O)} \quad \{\sum_{i=1}^{g} \omega_i d\mathbf{e}_i \mid 0 \le \omega_i < 1\} \text{ where } \{\mathbf{e}_1, \cdots, \mathbf{e}_g\} = \mathbf{E}_O(G) \setminus T$

APPENDIX B. SOME TRANSFORM FORMULAS

Based on technical considerations with respect to sublattices of $H_1(G,\mathbb{Z})$, we introduce new notions of zeta functions and L-functions, and show a theorem about some transform formulas on these functions (Theorem B.2), and provide an alternative proof of Theorem 3.12.

Again, let Λ be a full rank sublattice of $H_1(G, \mathbb{Z})$ and use the following notations: for $\omega \in \Omega$ and $\alpha \in H_1(G, \mathbb{Z})$, let $\underline{\omega} = \phi_1(\omega) + H_1(G, \mathbb{Z})^{\vee} \in \mathcal{X}$, $\underline{\underline{\omega}} = \phi(\omega) + \Lambda^{\vee} \in \mathcal{X}_{\Lambda}$, and $\underline{\alpha} = \alpha + \Lambda \in Q_{\Lambda}$. We introduce the following notions of zeta functions and L-functions:

(i)
$$z_{\alpha}(u) = \exp\left(\sum_{[C]\in\overline{\mathcal{C}},C^{\mathrm{ab}}=\alpha}\frac{1}{r(C)}u^{\tau(C)}\right)$$

(ii) $z_{\underline{\alpha}}(u) = \exp\left(\sum_{[C]\in\overline{\mathcal{C}},\underline{C^{\mathrm{ab}}}=\underline{\alpha}}\frac{1}{r(C)}u^{\tau(C)}\right)$

(iii) For $\alpha \in H_1(G, \mathbb{Z})$ and $\underline{\underline{\omega}} \in \mathcal{X}_{\Lambda}$, let

$$L_{(\Lambda,\alpha)}(u,\chi_{\omega}) = \exp\left(\sum_{[C]\in\overline{\mathcal{C}},C^{\mathrm{ab}}-\alpha\in\Lambda}\frac{\chi_{\omega}(C^{\mathrm{ab}}-\alpha)}{r(C)}u^{\tau(C)}\right).$$

Note that if $\alpha - \alpha' \in \Lambda$, then $\log L_{(\Lambda,\alpha)}(u,\chi) = \chi(\alpha' - \alpha) \cdot \log L_{(\Lambda,\alpha')}(u,\chi)$. (iv) $z_{\Lambda}(u) := z_{\underline{0}}(u)$ and $L_{\Lambda}(u,\chi) := L_{(\Lambda,0)}(u,\chi)$.

Remark B.1. Using the counting functions N(l), $N(\alpha, l)$ and $N(\underline{\alpha}, l)$ defined in 3.6, a classical result is that the Ihara zeta function has an alternative expression as $z(u) = \exp\left(\sum_{l=1}^{\infty} \frac{N(l)}{l}u^l\right)$, and it is can also be easily verified by definition that similar expressions hold for $z_{\alpha}(u)$ and $z_{\underline{\alpha}}(u)$ as $z_{\alpha}(u) = \exp\left(\sum_{l=1}^{\infty} \frac{N(\alpha, l)}{l}u^l\right)$ and $z_{\underline{\alpha}}(u) = \exp\left(\sum_{l=1}^{\infty} \frac{N(\alpha, l)}{l}u^l\right)$ respectively.

Proposition B.2. We have the following identities:

- (a) Transforms between $L_{\Lambda}(u, \chi_{\omega})$ and $z_{\alpha}(u)$ for $\underline{\omega} \in \mathcal{X}_{\Lambda}$ and $\alpha \in \Lambda$:
 - (i) For $\underline{\underline{\omega}} \in \mathcal{X}_{\Lambda}$, $\log L_{\Lambda}(u, \chi_{\omega}) = \sum_{\alpha \in \Lambda} \chi_{\omega}(\alpha) \log(z_{\alpha}(u))$. In particular, $\log z_{\Lambda}(u) = \sum_{\alpha \in \Lambda} \log(z_{\alpha}(u))$.

- (*ii*) For $\alpha \in \Lambda$, $\frac{1}{\operatorname{vol}(\mathcal{X}_{\Lambda})} \int_{\mathcal{X}_{\Lambda}} \chi_{-\omega}(\alpha) \log L_{\Lambda}(u, \chi_{\omega}) dV_{\omega} = \log z_{\alpha}(u).$
- (b) Transforms between $L(u, \chi_{\omega})$ and $z_{\alpha}(u)$:
- (i) For $\underline{\omega} \in \mathcal{X}$, $\log L(u, \chi_{\omega}) = \sum_{\alpha \in H_1(G,\mathbb{Z})} \chi_{\omega}(\alpha) \log(z_{\alpha}(u))$. In particular, $\log z(u) = \sum_{\alpha \in H_1(G,\mathbb{Z})} \log(z_{\alpha}(u))$. (ii) For $\alpha \in H_1(G,\mathbb{Z})$, $\frac{1}{\operatorname{vol}(\mathcal{X})} \int_{\mathcal{X}} \chi_{-\omega}(\alpha) \log L(u, \chi_{\omega}) dV_{\omega} = \log z_{\alpha}(u)$.
- (c) Transforms between $L(u, \chi_{\omega+\omega'})$ and $L_{(\Lambda,\alpha)}(u, \chi_{\omega})$ for $\underline{\underline{\omega}} \in \mathcal{X}_{\Lambda}$, $\underline{\omega'} \in \widehat{Q_{\Lambda}}$ and $\alpha \in H_1(G, \mathbb{Z})$:
 - (i) For $\underline{\omega} \in \mathcal{X}_{\Lambda}$ and $\underline{\omega}' \in \widehat{Q_{\Lambda}}$,

$$\log L(u, \chi_{\omega+\omega'}) = \sum_{\underline{\alpha} \in Q_{\Lambda}} \chi_{\omega+\omega'}(\alpha) \log L_{(\Lambda,\alpha)}(u, \chi_{\omega}).$$

(*ii*) For $\underline{\underline{\omega}} \in \mathcal{X}_{\Lambda}$ and $\alpha \in H_1(G, \mathbb{Z})$, $\frac{1}{|Q_{\Lambda}|} \sum_{\underline{\omega'} \in \widehat{Q_{\Lambda}}} \chi_{-(\omega+\omega')}(\alpha) \log L(u, \chi_{\omega+\omega'}) = \log L_{(\Lambda,\alpha)}(u, \chi_{\omega}).$

(d) Transforms between $L(u, \chi_{\omega})$ and $z_{\underline{\alpha}}(u)$ for $\underline{\omega} \in \widehat{Q_{\Lambda}}$ and $\underline{\alpha} \in Q_{\Lambda}$:

- (i) For $\underline{\omega} \in \widehat{Q_{\Lambda}}$, $\log L(u, \chi_{\omega}) = \sum_{\underline{\alpha} \in Q_{\Lambda}} \chi_{\underline{\omega}}(\underline{\alpha}) \log z_{\underline{\alpha}}(u)$. (ii) For $\underline{\alpha} \in Q_{\Lambda}$, $\frac{1}{|Q_{\Lambda}|} \sum_{\omega \in \widehat{Q_{\Lambda}}} \chi_{-\underline{\omega}}(\underline{\alpha}) \log L(u, \chi_{\omega}) = \log z_{\underline{\alpha}}(u)$.
- (e) For $\underline{\omega} \in \mathcal{X}_{\Lambda}$, we have
 - (i) $\log L(u, \chi_{\omega}) = \sum_{\underline{\alpha} \in Q_{\Lambda}} \chi_{\omega}(\alpha) \log L_{(\Lambda, \alpha)}(u, \chi_{\omega}), and$ (ii) $\frac{1}{|Q_{\Lambda}|} \sum_{\underline{\omega'} \in \widehat{Q_{\Lambda}}} \log L(u, \chi_{\omega+\omega'}) = \log L_{\Lambda}(u, \chi_{\omega}).$
- (f) We have
- (i) $\log z(u) = \sum_{\underline{\alpha} \in Q_{\Lambda}} \log z_{\underline{\alpha}}(u)$, and (ii) $\frac{1}{|Q_{\Lambda}|} \sum_{\underline{\omega} \in \widehat{Q_{\Lambda}}} \log L(u, \chi_{\omega}) = \log z_{\Lambda}(u)$.

Proof. (a)(i) follows from the definitions directly. (a)(ii) follows from the orthogonality relation in Proposition 2.13(a) as follows:

$$\frac{1}{\operatorname{vol}(\mathcal{X}_{\Lambda})} \int_{\mathcal{X}_{\Lambda}} \chi_{-\omega}(\alpha) \log L_{\Lambda}(u, \chi_{\omega}) dV_{\omega}$$

$$= \frac{1}{\operatorname{vol}(\mathcal{X}_{\Lambda})} \int_{\mathcal{X}_{\Lambda}} \chi_{-\omega}(\alpha) \left(\sum_{[C] \in \overline{\mathcal{C}}, C^{\operatorname{ab}} \in \Lambda} \frac{\chi_{\omega}(C^{\operatorname{ab}})}{r(C)} u^{\tau(C)} \right) dV_{\omega}$$

$$= \sum_{[C] \in \overline{\mathcal{C}}, C^{\operatorname{ab}} \in \Lambda} \left(\frac{1}{r(C) \operatorname{vol}(\mathcal{X}_{\Lambda})} \int_{\mathcal{X}_{\Lambda}} \chi_{-\omega}(\alpha) \chi_{\omega}(C^{\operatorname{ab}}) dV_{\omega} \right) u^{\tau(C)}$$

$$= \sum_{[C] \in \overline{\mathcal{C}}, C^{\operatorname{ab}} \in \Lambda} \left(\frac{1}{r(C) \operatorname{vol}(\mathcal{X}_{\Lambda})} \int_{\mathcal{X}_{\Lambda}} \overline{\chi_{\omega}(\alpha)} \chi_{\omega}(C^{\operatorname{ab}}) dV_{\omega} \right) u^{\tau(C)}$$

$$= \sum_{[C] \in \overline{\mathcal{C}}, C^{\operatorname{ab}} \in \Lambda} \frac{\delta_{C^{\operatorname{ab}}\alpha}}{r(C)} u^{\tau(C)} = \sum_{[C] \in \overline{\mathcal{C}}, C^{\operatorname{ab}} = \alpha} \frac{1}{r(C)} u^{\tau(C)} = \log z_{\alpha}(u).$$

(b) is a special case of (a) when $\Lambda = H_1(G, \mathbb{Z})$.

(c)(i) follows from the definitions. For (c)(ii), we will use the orthogonality relation in Proposition 2.13(b).

$$\begin{split} &\frac{1}{|Q_{\Lambda}|} \sum_{\underline{\omega'} \in \widehat{Q_{\Lambda}}} \chi_{-(\omega+\omega')}(\alpha) \log L(u, \chi_{\omega+\omega'}) \\ &= \frac{1}{|Q_{\Lambda}|} \sum_{\underline{\omega'} \in \widehat{Q_{\Lambda}}} \chi_{-(\omega+\omega')}(\alpha) \sum_{[C] \in \overline{\mathcal{C}}} \frac{\chi_{\omega+\omega'}(C^{\mathrm{ab}})}{r(C)} u^{\tau(C)} \\ &= \sum_{[C] \in \overline{\mathcal{C}}} \left(\frac{\chi_{-\omega}(\alpha)\chi_{\omega}(C^{\mathrm{ab}})}{r(C)|Q_{\Lambda}|} \sum_{\underline{\omega'} \in \widehat{Q_{\Lambda}}} \chi_{-\omega'}(\alpha)\chi_{\omega'}(C^{\mathrm{ab}}) \right) u^{\tau(C)} \\ &= \sum_{[C] \in \overline{\mathcal{C}}} \left(\frac{\chi_{\omega}(C^{\mathrm{ab}} - \alpha)}{r(C)|Q_{\Lambda}|} \sum_{\underline{\omega'} \in \widehat{Q_{\Lambda}}} \overline{\chi_{\omega'}(\alpha)}\chi_{\omega'}(C^{\mathrm{ab}}) \right) u^{\tau(C)} \\ &= \sum_{[C] \in \overline{\mathcal{C}}} \delta_{\underline{C^{\mathrm{ab}}}} \frac{\alpha}{\alpha} \frac{\chi_{\omega}(C^{\mathrm{ab}} - \alpha)}{r(C)} u^{\tau(C)} \\ &= \sum_{[C] \in \overline{\mathcal{C}}, C^{\mathrm{ab}} - \alpha \in \Lambda} \frac{\chi_{\omega}(C^{\mathrm{ab}} - \alpha)}{r(C)} u^{\tau(C)} \\ &= \log L_{(\Lambda,\alpha)}(u, \chi_{\omega}). \end{split}$$

(d) is a special case of (c) by restricting $\underline{\omega} + \underline{\omega}' = \underline{\omega}' \in \widehat{Q}_{\Lambda}$. (e)(i) is a special case of (c)(i) by setting $\overline{\omega}' = 0$. (e)(ii) is a special case of (c)(ii) by setting $\alpha = 0$. (f)(i) is a special case of (c)(i) by setting $\omega = \omega' = 0$. (f)(ii) is a special case of (c)(ii) by setting $\omega = 0$ and $\alpha = 0$.

As an application of the transform formulas in Proposition B.2, here we provide another proof of Theorem 3.12 without referring to the interpretation of $\mathcal{K}(\omega, l)$ as $\sum_{C \in \mathcal{C}_l} \chi_{\omega}(C)$ (Lemma 3.8(a)).

An alternative proof of Theorem 3.12. For (a)(i), note that $z_{\alpha}(u) = \exp\left(\sum_{l=1}^{\infty} \frac{N(\alpha,l)}{l}u^l\right)$. Therefore, $u\frac{d}{du}\log z_{\alpha}(u) = \sum_{l\geq 1} N(\alpha,l)u^l$. Using the relation $\frac{1}{\operatorname{vol}(\mathcal{X})} \int_{\mathcal{X}} \chi_{-\omega}(\alpha)\log L(u,\chi_{\omega})dV_{\omega} = \log z_{\alpha}(u)$ (Proposition B.2(b)(ii)), we have

$$\sum_{l\geq 1} N(\alpha, l)u^l = \frac{1}{\operatorname{vol}(\mathcal{X})} \int_{\mathcal{X}} \chi_{-\omega}(\alpha) \left(u \frac{d}{du} \log L(u, \chi_{\omega}) \right) dV_{\omega}.$$

On the other hand,

$$u\frac{d}{du}\log L(u,\chi_{\omega}) = u\frac{d}{du}\sum_{\lambda\in\operatorname{Spec}W_{1,\omega}}\log(1-\lambda u)$$
$$= \sum_{\lambda\in\operatorname{Spec}W_{1,\omega}}\sum_{l\geq 1}(\lambda u)^{l}$$
$$= \sum_{l\geq 1}\left(\sum_{\lambda\in\operatorname{Spec}W_{1,\omega}}\lambda^{l}\right)u^{l}$$
$$= \sum_{l\geq 1}\mathcal{K}(\omega,l)u^{l}.$$

Therefore,

$$\sum_{l\geq 1} N(\alpha, l) u^{l} = \frac{1}{\operatorname{vol}(\mathcal{X})} \int_{\mathcal{X}} \chi_{-\omega}(\alpha) \left(\sum_{l\geq 1} \mathcal{K}(\omega, l) u^{l} \right) dV_{\omega}$$
$$= \sum_{l\geq 1} \left(\frac{1}{\operatorname{vol}(\mathcal{X})} \int_{\mathcal{X}} \chi_{-\omega}(\alpha) \mathcal{K}(\omega, l) dV_{\omega} \right) u^{l},$$

which means that

$$N(\alpha, l) = \frac{1}{\operatorname{vol}(\mathcal{X})} \int_{\mathcal{X}} \chi_{-\omega}(\alpha) \mathcal{K}(\omega, l) dV_{\omega}$$

for all $l \ge 1$. For (a)(ii),

$$\sum_{\alpha \in H_1(G,\mathbb{Z})} \chi_{\omega}(\alpha) N(\alpha, l) = \sum_{\alpha \in H_1(G,\mathbb{Z})} \chi_{\omega}(\alpha) \left(\frac{1}{\operatorname{vol}(\mathcal{X})} \int_{\mathcal{X}} \chi_{-\omega'}(\alpha) \mathcal{K}(\omega', l) dV_{\omega'} \right)$$
$$= \int_{\mathcal{X}} \left(\frac{1}{\operatorname{vol}(\mathcal{X})} \sum_{\alpha \in H_1(G,\mathbb{Z})} \chi_{\omega-\omega'}(\alpha) \right) \mathcal{K}(\omega', l) dV_{\omega'}$$
$$= \int_{\mathcal{X}} \left(\sum_{\gamma \in H_1(G,\mathbb{Z})^{\vee}} \delta(\omega - \omega' - \gamma) \right) \mathcal{K}(\omega', l) dV_{\omega'}$$
$$= \mathcal{K}(\omega, l).$$

Here we note that the last identity follows from the fact that \mathcal{X} is the fundamental domain of the lattice $H_1(G,\mathbb{Z})^{\vee}$ in $\mathcal{H}^1(G)$.

For (b), we may use an analogous argument as in the proof of (a). More specifically, (b)(i) follows from

$$\begin{split} \sum_{l\geq 1} N(\underline{\alpha}, l) u^l &= u \frac{d}{du} \log z_{\underline{\alpha}}(u) \\ (\text{Proposition B.2(d)(ii)}) &= \frac{1}{|Q_\Lambda|} \sum_{\underline{\omega} \in \widehat{Q_\Lambda}} \chi_{-\underline{\omega}}(\underline{\alpha}) \left(u \frac{d}{du} \log L(u, \chi_{\omega}) \right) \\ &= \frac{1}{|Q_\Lambda|} \sum_{\underline{\omega} \in \widehat{Q_\Lambda}} \chi_{-\underline{\omega}}(\underline{\alpha}) \left(\sum_{l\geq 1} \mathcal{K}(\underline{\omega}, l) u^l \right) \\ &= \sum_{l\geq 1} \left(\frac{1}{|Q_\Lambda|} \sum_{\underline{\omega} \in \widehat{Q_\Lambda}} \chi_{-\underline{\omega}}(\underline{\alpha}) \mathcal{K}(\underline{\omega}, l) \right) u^l, \end{split}$$

and (b)(ii) follows from

$$\begin{split} \sum_{\underline{\alpha}\in Q_{\Lambda}}\chi_{\underline{\omega}}(\underline{\alpha})N(\underline{\alpha},l) &= \sum_{\underline{\alpha}\in Q_{\Lambda}}\chi_{\underline{\omega}}(\underline{\alpha})\left(\frac{1}{|Q_{\Lambda}|}\sum_{\underline{\omega'}\in\widehat{Q_{\Lambda}}}\chi_{-\underline{\omega'}}(\underline{\alpha})\mathcal{K}(\underline{\omega'},l)\right) \\ &= \sum_{\underline{\omega'}\in\widehat{Q_{\Lambda}}}\left(\frac{1}{|Q_{\Lambda}|}\sum_{\underline{\alpha}\in Q_{\Lambda}}\chi_{\underline{\omega}}(\underline{\alpha})\chi_{-\underline{\omega'}}(\underline{\alpha})\right)\mathcal{K}(\underline{\omega'},l) \\ &= \sum_{\underline{\omega'}\in\widehat{Q_{\Lambda}}}\delta_{\underline{\omega}\ \underline{\omega'}}\mathcal{K}(\underline{\omega'},l) = \mathcal{K}(\underline{\omega},l). \end{split}$$

	٦
	1
	1
_	 _