

Dynamical Dark Energy Implies a Coupled Dark Sector: Insights from DESI DR2 via a Data-Driven Approach

CHANGYU YOU,¹ DAN WANG,¹ AND TAO YANG ¹

¹*School of Physics and Technology, Wuhan University, Wuhan 430072, China*

ABSTRACT

Recent observations from the Dark Energy Spectroscopic Instrument (DESI) Data Release 2 (DR2) have revealed compelling evidence for dynamical dark energy, challenging the Λ CDM paradigm. In this work, we adopt a data-driven, model-independent approach to reconstruct the dark energy equation of state (EoS) and its potential interaction with dark matter using combined background cosmological datasets, including DESI DR2, cosmic chronometers, observational Hubble data, and Type Ia supernovae. Using Gaussian Process regression and a non-parametric formalism, we first confirm a $\sim 2\sigma$ indication of dynamical dark energy, featuring a phantom crossing around redshift $z \sim 0.4$, consistent with DESI results. We then explore the implications of dynamical EoS from DESI DR2 for dark sector coupling. Incorporating priors on the EoS from DESI DR2, we find a 2.2σ signal for non-zero interactions between dark energy and dark matter at low redshift. Our results suggest that if DESI's evidence for time-varying dark energy is confirmed, a coupled dark sector may be a necessary extension beyond Λ CDM.

1. INTRODUCTION

The Λ CDM cosmology, established as the standard paradigm in modern cosmology, has been highly successful in explaining a wide range of astrophysical phenomena – from cosmic microwave background (CMB) anisotropies to large-scale structure formation (A. G. Riess et al. 1998; S. Perlmutter et al. 1999; G. Hinshaw et al. 2013; N. Aghanim et al. 2020; M. Tegmark et al. 2004; D. J. Eisenstein et al. 2005). However, growing discrepancies between early- and late-universe observations – most notably the Hubble tension ($a \sim 5\sigma$ discrepancy in H_0 (A. G. Riess et al. 2022; N. Aghanim et al. 2020)), the S_8 tension (N. Aghanim et al. 2020; T. M. C. Abbott et al. 2025), and cosmic shear anomalies (I. G. McCarthy et al. 2018; H. Hildebrandt et al. 2020; M. Asgari et al. 2020) – challenge the model’s validity. Proposed resolutions include modifications to gravity (M. Kunz & D. Sapone 2007; T. Clifton et al. 2012; W. Hu & I. Sawicki 2007) and dynamical dark energy scenarios (E. J. Copeland et al. 2006; S. Tsujikawa 2013; C. Armendariz-Picon et al. 2001).

The first data release (DR1) of the Dark Energy Spectroscopic Instrument (DESI) (A. G. Adame et al. 2025) suggested evidence for dynamical dark energy, inconsistent with the Λ CDM framework. This result has spurred significant interest in constraining the dark en-

ergy equation of state (EoS) through various data analysis approaches (K. Lodha et al. 2025; D. Shlivko & P. J. Steinhardt 2024; Y. Tada & T. Terada 2024; W. Giarè et al. 2024; I. D. Gialamas et al. 2025). More recently, the second data release (DR2) from DESI (M. Abdul Karim et al. 2025), based on three years of observations, has reinforced these findings. Using parameterized dark energy models, such as the Chevallier-Polarski-Linder (CPL) parameterization (E. V. Linder 2003; N. Aghanim et al. 2020; P. A. R. Ade et al. 2016; D. M. Scolnic et al. 2018), DESI DR2 reports a 2.8σ to 4.2σ statistical preference for time-varying dark energy over the standard Λ CDM model.

The study of dynamical dark energy models has intensified in recent decades, with each model offering distinct advantages and limitations (E. J. Copeland et al. 2006; C. Armendariz-Picon et al. 2001; E. V. Linder 2008; D. M. Scolnic et al. 2018). Quintessence – a widely studied dynamical model in which dark energy is represented by a scalar field – has received significant attention (S. Tsujikawa 2013; E. V. Linder 2008; E. N. Saridakis & S. V. Sushkov 2010). However, recent DESI DR2 results challenge this framework by revealing a phantom crossing at $z \simeq 0.4$, with $w > -1$ at $z < 0.4$ and $w < -1$ at $z > 0.4$, a behavior not permitted in standard quintessence models. Among the various parameterizations designed to capture such dynamics, the CPL form, $w(z) = w_0 + w_a z / (1 + z)$, stands out as the most widely used phenomenological approach and has been extensively constrained with DESI BAO data

Corresponding author: Tao Yang
 Email: yangtao@whu.edu.cn

recently (A. G. Adame et al. 2025; M. Abdul Karim et al. 2025; D. Wang 2024; C.-G. Park et al. 2024). In this work, we model dark energy as a phenomenological fluid, reconstruct its EoS using a model-independent (non-parametric) approach, and reassess its evolution in light of the DESI DR2 results.

Another scenario beyond the Λ CDM model involves considering interactions between the dark sectors. Given the unknown nature of both dark matter and dark energy, it is plausible that they may interact through non-zero couplings. This possibility has motivated the development of interacting dark energy (IDE) models (L. Amendola 2000; B. Wang et al. 2016; R.-G. Cai & A. Wang 2005; C. G. Boehmer et al. 2008; L. Amendola & C. Quercellini 2003; E. Di Valentino et al. 2020; W. Yang et al. 2018), which may help alleviate the aforementioned cosmological tensions and the coincidence problem (S. Weinberg 2000). The interaction between dark energy and dark matter has been investigated using recent DESI data, assuming a specific form of the coupling function (A. Chakraborty et al. 2025). However, in the absence of a theoretically preferred interaction model, it is preferable to constrain the coupling of the dark sector using a model-independent approach.

In this paper, we adopt the non-parametric, data-driven method proposed by T. Yang et al. (2015); T. Yang (2020) to reconstruct the interaction between dark energy and dark matter using background cosmological data sets. We highlight the degeneracy between dark sector coupling and the EoS of dark energy in the context of this model-independent framework. By incorporating priors on the dynamical EoS from the DESI DR2 results, we identify evidence for non-zero dark sector interactions at low redshift. Our findings indicate that the dynamical dark energy inferred from DESI DR2 inevitably implies a coupled dark sector when analyzed from a data-driven perspective using background cosmological data sets.

2. METHOD

In interacting dark sector scenarios, the dark matter density evolves as $\rho_c \sim f/a^3$, rather than following the standard a^{-3} scaling, where f is an arbitrary function representing the coupling between dark energy and dark matter (S. Das et al. 2006). To remain consistent with the equivalence principle and Solar System tests of gravity (B. Bertotti et al. 2003; C. M. Will 2014), we do not couple dark energy to baryons². Within the flat

Friedmann-Lemaître-Robertson-Walker(FLRW) framework, the Friedmann equations and continuity equations for every species take the form:

$$3H^2 = \rho_{de} + \rho_c + \rho_b, \quad (1)$$

$$\dot{\rho}_b + 3H\rho_b = 0, \quad (2)$$

$$\dot{\rho}_c + 3H\rho_c = 3H_0^2\Omega_c \frac{\dot{f}}{a^3 f_0}, \quad (3)$$

$$\dot{\rho}_{de} + 3H(1+w_{de})\rho_{de} = -3H_0^2\Omega_c \frac{\dot{f}}{a^3 f_0} \quad (4)$$

Here, $a = 1/(1+z)$ is the scale factor, $H = \dot{a}/a$ is the Hubble parameter, f_0 denotes the present-day value of the coupling function f , and dots represent derivatives with respect to time. The density parameter for a component X is given by $\Omega_X = \rho_{X0}/(3H_0^2)$, where ρ_{X0} is its present energy density. Since we focus on the late-time Universe, where dark energy and matter dominate, the radiation contributions are ignored. For simplicity, we adopt natural units with $M_{\text{Pl}} = 1$ throughout our analysis.

To avoid specifying prior information on the three parameters f_0 , Ω_c , and Ω_b during the reconstruction of the coupling, we use $\Omega_f = \Omega_c \frac{f}{f_0} + \Omega_b$ to investigate the interaction between dark energy and dark matter. A constant Ω_f (i.e., $\dot{\Omega}_f = 0$) indicates the absence of coupling between the dark sectors. Using Eqs. (1-2) we get the EoS of dark energy:

$$w_{de} = \frac{(-2\dot{H} - 3H^2)}{(3H^2 - 3H_0^2 \frac{\Omega_f}{a^3})} \quad (5)$$

In the uncoupled limit ($f \equiv f_0$), Ω_f reduces to $\Omega_c + \Omega_b = \Omega_m$. The equation above enables the reconstruction of the EoS of dark energy directly from cosmological expansion data.

For coupled dark energy, the coupling and the EoS of dark energy are related through Eq. (5),

$$\Omega_f = \frac{a^3}{3H_0^2} \left(3H^2 + \frac{2\dot{H} + 3H^2}{w_{de}} \right), \quad (6)$$

$$\dot{\Omega}_f = \frac{a^3}{3H_0^2} \left((9H^3 + 6H\dot{H}) + \frac{2\ddot{H} + 12H\dot{H} + 9H^3}{w_{de}} - \frac{(2\dot{H} + 3H^2)\dot{w}_{de}}{w_{de}^2} \right). \quad (7)$$

The dark energy EoS w_{de} exhibits degeneracy with dark sector interactions in background expansion observables (e.g., Hubble parameter, luminosity distances). For a fixed expansion history, reconstructing either EoS (w_{de}) or coupling (Ω_f or $\dot{\Omega}_f$) requires prior

² Introducing a universal coupling between dark energy and the total non-relativistic matter (including subdominant baryons) does not alter our conclusions.

knowledge of the other component. This degeneracy implies that dynamical dark energy and dark sector couplings can mutually compensate for each other, yielding identical expansion signatures in observational data.

To reconstruct these parameters from observational data, we convert all time derivatives to redshift derivatives using the transformation $\frac{d}{dt} = -H(1+z)\frac{d}{dz}$. Given a prior on w_{de} , our framework can simultaneously reconstruct the coupling parameters: Ω_f and its evolution $\dot{\Omega}_f$. A non-zero $\dot{\Omega}_f$ signals the presence of interactions between the dark sectors.

3. DATA

3.1. Baryonic acoustic oscillations

- 6 $H(z)$ data from DESI DR2 (M. Abdul Karim et al. 2025). The DESI-BAO measurements are expressed by three ratios: the three-dimensional BAO mode \tilde{D}_Y , the transverse mode \tilde{D}_M , and the radial mode \tilde{D}_H . In this work, we focus on the radial mode.

Along the line of sight we measure:

$$\tilde{D}_H = \frac{D_H}{r_d} = \frac{c}{r_d H(z)} \quad (8)$$

So we get $H(z)$:

$$H(z) = \frac{c}{r_d \frac{D_H}{r_d}} = \frac{c}{r_d \tilde{D}_H} \quad (9)$$

- 18 $H(z)$ data from the homogenized model-independent OHD (BAO features) compiled in Table 2 of J. Magana et al. (2018). We replace the three highest-redshift Lyman- α (Ly α) BAO measurements (A. Font-Ribera et al. 2014; T. Delubac et al. 2015; J. E. Bautista et al. 2017) at $z = 2.33, 2.34$ and 2.36 with the updated $z = 2.34$ eBOSS DR14 result (M. Blomqvist et al. 2019; V. de Sainte Agathe et al. 2019). The $H(z)$ value at this redshift is derived using the same methodology applied to DESI BAO data.

Note that we adopt the same sound horizon $r_d = 147.33$ Mpc at the drag epoch using in (J. Magana et al. 2018) from Planck measurements to ensure that all $H(z)$ from BAO have the same prior on r_d .

3.2. Cosmic chronometers

- 31 $H(z)$ data from cosmic chronometers (CC) obtained using the differential-age technique in Table 1 of A. Gómez-Valent & L. Amendola (2018).

3.3. Pantheon+MCT SNe Ia

- 6 $E(z)$ data from Pantheon+MCT SNe Ia Measurements given by A. G. Riess et al. (2018). Here $E(z) = H(z)/H_0$ denotes the dimensionless Hubble parameter. For the $z = 1.5$ point, we employ a Gaussian likelihood approximation following A. Gómez-Valent (2019), incorporating the full covariance matrix of uncertainties.

4. RECONSTRUCTIONS AND RESULTS

To ensure self-consistency across the four observational datasets employed in our reconstructions, we standardize all $H(z)$ data measurements to the dimensionless Hubble parameter $E(z) \equiv H(z)/H_0$, adopting a fiducial H_0 value. We use Gaussian Process (GP) regression³, a machine learning method notable for its ability to predict the evolution of derived cosmological parameters in a model-independent manner (M. Seikel et al. 2012; A. Shafieloo et al. 2012). The fiducial value of H_0 is determined using the combined DESI DR2+OHD+CC data. We then reconstruct $E(z)$, $E'(z)$, $E''(z)$, and $E'''(z)$ based on the combination of all four datasets. We obtain a mean value of $H_0 = 68.87 \text{ km s}^{-1} \text{ Mpc}^{-1}$, which lies between the estimates derived from Planck (N. Aghanim et al. 2020) and SNe Ia (A. G. Riess et al. 2022). We rewrite Eqs. (5-7) to substitute $H(z)$ with $E(z)$ by dividing $(H_0)^n$ on both sides of the equations. Here, n corresponds to the dimensional order of each equation with respect to H_0 . Then the dimensionless parameters we can reconstruct are actually w_{de} , Ω_f , $\frac{\dot{\Omega}_f}{H_0}$, etc.

For the uncoupled case, we adopt the value $\Omega_f = \Omega_m = 0.31$ from Planck (N. Aghanim et al. 2020). The equation of state for uncoupled dark energy, reconstructed using different combinations of data sets, is shown in Fig. 1. Note that for the reconstruction of $w(z)$ using the CC+OHD+SNIa data combination, the fiducial value of H_0 is $68.06 \text{ km s}^{-1} \text{ Mpc}^{-1}$, derived from the same data set with GP. As shown in Fig. 1, without DESI BAO data, the reconstruction of w_{de} is fully consistent with the ΛCDM model ($w = -1$). However, incorporating DESI DR2 data reveals a 2σ indication of dynamical dark energy evolution at $z < 1$, featuring a phantom crossing at $z \simeq 0.4$, in agreement with the DESI DR2 findings – although the deviation from $w = -1$ at lower redshift is less pronounced than that reported by DESI. The overall trend of the reconstructed $w(z)$ shows a decrease with increasing redshift. However, the confidence level diminishes at higher redshifts

³ <https://github.com/JCGoran/GaPP/tree/feature/python3>




Figure 1. Reconstructions of the dark energy equation of state in the uncoupled case using different datasets. We present the 68% and 95% confidence levels (CL) as dark and light shaded bands, respectively. Dashed lines represent the Λ CDM model ($w = -1$).

due to larger uncertainties, stemming from the limited availability of observational data in that regime.

To reconstruct the interaction between dark energy and dark matter, as demonstrated in Sec. 2, the equation of state of dark energy must be provided *a priori* to break the degeneracy. Instead of directly reconstructing f and \dot{f} , as done in T. Yang (2020), we reconstruct $\Omega_f = \Omega_c \frac{f}{f_0} + \Omega_b$ and its derivative, thereby avoiding potential issues related to the priors of f , Ω_c , and Ω_b at $z = 0$. Our focus is on whether $\dot{\Omega}_f = 0$, which is independent of the specific values of f_0 , Ω_c , and Ω_b .

In this paper, we adopt two priors for the equation of state by extracting constraints on the CPL form of $w(z)$ from both Planck (N. Aghanim et al. 2020) and DESI DR2 (M. Abdul Karim et al. 2025) (Planck: $w_0 = -0.957 \pm 0.080$, $w_a = -0.29^{+0.32}_{-0.26}$; DESI DR2: $w_0 = -0.838 \pm 0.055$, $w_a = -0.62^{+0.22}_{-0.19}$), with the former being consistent with the Λ CDM model and the latter exhibiting a significant signal of dynamical dark energy.

As shown in the upper panel of Fig. 2, the reconstruction of dark sector interactions – based on the combined datasets from DESI DR2, OHD, CC, and SNIa using Planck priors – is consistent with a non-coupled dark energy scenario. The reconstructed Ω_f is also consistent with the value of Ω_m derived from Planck. Compared to previous work (T. Yang 2020), the previously reported $> 1\sigma$ coupling at high redshift is now reduced to within 1σ .

Interestingly, as demonstrated in the lower panel of Fig. 2, when the dynamical EoS prior from DESI is adopted, a 2.2σ signal of dark sector interactions emerges at $z < 0.5$. This behavior suggests that, from

a model-independent and data-driven perspective, dark sector interactions will inevitably emerge if the dynamical dark energy indicated by DESI is confirmed.

5. CONCLUSION AND DISCUSSIONS

In this paper, we present a data-driven, model-independent reconstruction of the dark energy equation of state and its interaction with dark matter using cosmic expansion data. By incorporating the new DESI DR2 BAO measurements into the previous CC+OHD+SNe Ia combination, we identify a 2σ dynamical signature in the uncoupled dark energy scenario, including a phantom crossing around $z \sim 0.4$, consistent with DESI's findings. Intriguingly, when the constraints on the CPL form of $w(z)$ from DESI are adopted as a prior to reconstruct the interaction between dark energy and dark matter, a 2.2σ signal of non-zero coupling at low redshift emerges. Our findings suggest that the significant dynamical signal of dark energy revealed by recent DESI data may also point to a notable interaction between the dark sectors. We need to investigate both the dynamics and interactions of dark energy as more data and improved methodologies become available in the future.

Employing a data-driven, model-independent GP framework, we determine the fiducial value of H_0 and reconstruct $E(z)$ along with its derivatives. While the reconstruction of H_0 is sensitive to both dataset selection and the choice of GP kernel (as discussed in J. P. Johnson & H. K. Jassal (2025)), we adopt the Matérn-9/2 kernel to mitigate potential overfitting artifacts associated with the widely used Radial Basis Function (RBF) kernel, which can amplify obser-




Figure 2. Reconstructions of the dark sector interactions from DESI DR2+CC+OHD+SNe Ia. We adopt two priors of $w(z) = w_0 + w_a \frac{z}{1+z}$ which are extracted from Planck (upper panel) and DESI DR2 (lower panel). The dashed lines denote $\Omega_m = 0.31$ from Planck (left panel) and zero coupling (right panel), respectively.

vational uncertainties (J. P. Johnson & H. K. Jassal 2025). The RBF kernel yields $h = 0.6932$ (where $H_0 = 100h \text{ km s}^{-1} \text{ Mpc}^{-1}$), slightly higher than the fiducial value $h = 0.6887$ used in our analysis. However, dependencies on H_0 cancel out in the subsequent reconstructions of w , Ω_f , and $\dot{\Omega}_f/H_0$, rendering our results robust to the choice of kernel.

Our methodology is inherently flexible within the data-driven paradigm, accommodating a wide range of machine learning (ML) techniques for regression and parameter reconstruction from large cosmological datasets. While GP is commonly used for non-parametric reconstructions, our framework is readily extendable to other ML approaches, including Neural Networks (NNs) (J. Fluri et al. 2018) and Deep Learning (D. George & E. A.

Huerta 2018). This versatility ensures that our approach remains adaptable to a wide range of problems and applicable to additional cosmological data sets (e.g., cosmological perturbation data). We leave such extensions for future work.

ACKNOWLEDGMENTS

This work is supported by “the Fundamental Research Funds for the Central Universities” under the reference No. 2042024FG0009. The numerical calculations in this paper have been done on the supercomputing system in the Supercomputing Center of Wuhan University.

REFERENCES

- Abbott, T. M. C., et al. 2025,
<https://arxiv.org/abs/2503.13632>
- Abdul Karim, M., et al. 2025,
<https://arxiv.org/abs/2503.14738>

- Adame, A. G., et al. 2025, JCAP, 02, 021, doi: [10.1088/1475-7516/2025/02/021](https://doi.org/10.1088/1475-7516/2025/02/021)
- Ade, P. A. R., et al. 2016, Astron. Astrophys., 594, A14, doi: [10.1051/0004-6361/201525814](https://doi.org/10.1051/0004-6361/201525814)
- Aghanim, N., et al. 2020, Astron. Astrophys., 641, A6, doi: [10.1051/0004-6361/201833910](https://doi.org/10.1051/0004-6361/201833910)
- Amendola, L. 2000, Phys. Rev. D, 62, 043511, doi: [10.1103/PhysRevD.62.043511](https://doi.org/10.1103/PhysRevD.62.043511)
- Amendola, L., & Quercellini, C. 2003, Phys. Rev. D, 68, 023514, doi: [10.1103/PhysRevD.68.023514](https://doi.org/10.1103/PhysRevD.68.023514)
- Armendariz-Picon, C., Mukhanov, V. F., & Steinhardt, P. J. 2001, Phys. Rev. D, 63, 103510, doi: [10.1103/PhysRevD.63.103510](https://doi.org/10.1103/PhysRevD.63.103510)
- Asgari, M., et al. 2020, Astron. Astrophys., 634, A127, doi: [10.1051/0004-6361/201936512](https://doi.org/10.1051/0004-6361/201936512)
- Bautista, J. E., et al. 2017, Astron. Astrophys., 603, A12, doi: [10.1051/0004-6361/201730533](https://doi.org/10.1051/0004-6361/201730533)
- Bertotti, B., Iess, L., & Tortora, P. 2003, Nature, 425, 374, doi: [10.1038/nature01997](https://doi.org/10.1038/nature01997)
- Blomqvist, M., et al. 2019, Astron. Astrophys., 629, A86, doi: [10.1051/0004-6361/201935641](https://doi.org/10.1051/0004-6361/201935641)
- Boehmer, C. G., Caldera-Cabral, G., Lazkoz, R., & Maartens, R. 2008, Phys. Rev. D, 78, 023505, doi: [10.1103/PhysRevD.78.023505](https://doi.org/10.1103/PhysRevD.78.023505)
- Cai, R.-G., & Wang, A. 2005, JCAP, 03, 002, doi: [10.1088/1475-7516/2005/03/002](https://doi.org/10.1088/1475-7516/2005/03/002)
- Chakraborty, A., Chanda, P. K., Das, S., & Dutta, K. 2025, <https://arxiv.org/abs/2503.10806>
- Clifton, T., Ferreira, P. G., Padilla, A., & Skordis, C. 2012, Phys. Rept., 513, 1, doi: [10.1016/j.physrep.2012.01.001](https://doi.org/10.1016/j.physrep.2012.01.001)
- Copeland, E. J., Sami, M., & Tsujikawa, S. 2006, Int. J. Mod. Phys. D, 15, 1753, doi: [10.1142/S021827180600942X](https://doi.org/10.1142/S021827180600942X)
- Das, S., Corasaniti, P. S., & Khoury, J. 2006, Phys. Rev. D, 73, 083509, doi: [10.1103/PhysRevD.73.083509](https://doi.org/10.1103/PhysRevD.73.083509)
- de Sainte Agathe, V., et al. 2019, Astron. Astrophys., 629, A85, doi: [10.1051/0004-6361/201935638](https://doi.org/10.1051/0004-6361/201935638)
- Delubac, T., et al. 2015, Astron. Astrophys., 574, A59, doi: [10.1051/0004-6361/201423969](https://doi.org/10.1051/0004-6361/201423969)
- Di Valentino, E., Melchiorri, A., Mena, O., & Vagnozzi, S. 2020, Phys. Dark Univ., 30, 100666, doi: [10.1016/j.dark.2020.100666](https://doi.org/10.1016/j.dark.2020.100666)
- Eisenstein, D. J., et al. 2005, Astrophys. J., 633, 560, doi: [10.1086/466512](https://doi.org/10.1086/466512)
- Fluri, J., Kacprzak, T., Refregier, A., et al. 2018, Phys. Rev. D, 98, 123518, doi: [10.1103/PhysRevD.98.123518](https://doi.org/10.1103/PhysRevD.98.123518)
- Font-Ribera, A., et al. 2014, JCAP, 05, 027, doi: [10.1088/1475-7516/2014/05/027](https://doi.org/10.1088/1475-7516/2014/05/027)
- George, D., & Huerta, E. A. 2018, Phys. Lett. B, 778, 64, doi: [10.1016/j.physletb.2017.12.053](https://doi.org/10.1016/j.physletb.2017.12.053)
- Gialamas, I. D., Hütsi, G., Kannike, K., et al. 2025, Phys. Rev. D, 111, 043540, doi: [10.1103/PhysRevD.111.043540](https://doi.org/10.1103/PhysRevD.111.043540)
- Giarè, W., Najafi, M., Pan, S., Di Valentino, E., & Firouzjaee, J. T. 2024, JCAP, 10, 035, doi: [10.1088/1475-7516/2024/10/035](https://doi.org/10.1088/1475-7516/2024/10/035)
- Gómez-Valent, A. 2019, JCAP, 05, 026, doi: [10.1088/1475-7516/2019/05/026](https://doi.org/10.1088/1475-7516/2019/05/026)
- Gómez-Valent, A., & Amendola, L. 2018, JCAP, 04, 051, doi: [10.1088/1475-7516/2018/04/051](https://doi.org/10.1088/1475-7516/2018/04/051)
- Hildebrandt, H., et al. 2020, Astron. Astrophys., 633, A69, doi: [10.1051/0004-6361/201834878](https://doi.org/10.1051/0004-6361/201834878)
- Hinshaw, G., Larson, D., Komatsu, E., et al. 2013, Astrophys. J. Supp. S., 208, 19, doi: [10.1088/0067-0049/208/2/19](https://doi.org/10.1088/0067-0049/208/2/19)
- Hu, W., & Sawicki, I. 2007, Phys. Rev. D, 76, 064004, doi: [10.1103/PhysRevD.76.064004](https://doi.org/10.1103/PhysRevD.76.064004)
- Johnson, J. P., & Jassal, H. K. 2025, <https://arxiv.org/abs/2503.04273>
- Kunz, M., & Sapone, D. 2007, Phys. Rev. Lett., 98, 121301, doi: [10.1103/PhysRevLett.98.121301](https://doi.org/10.1103/PhysRevLett.98.121301)
- Linder, E. V. 2003, Phys. Rev. Lett., 90, 091301, doi: [10.1103/PhysRevLett.90.091301](https://doi.org/10.1103/PhysRevLett.90.091301)
- Linder, E. V. 2008, Gen. Rel. Grav., 40, 329, doi: [10.1007/s10714-007-0550-z](https://doi.org/10.1007/s10714-007-0550-z)
- Lodha, K., et al. 2025, Phys. Rev. D, 111, 023532, doi: [10.1103/PhysRevD.111.023532](https://doi.org/10.1103/PhysRevD.111.023532)
- Magana, J., Amante, M. H., Garcia-Aspeitia, M. A., & Motta, V. 2018, Mon. Not. Roy. Astron. Soc., 476, 1036, doi: [10.1093/mnras/sty260](https://doi.org/10.1093/mnras/sty260)
- McCarthy, I. G., Bird, S., Schaye, J., et al. 2018, Mon. Not. Roy. Astron. Soc., 476, 2999, doi: [10.1093/mnras/sty377](https://doi.org/10.1093/mnras/sty377)
- Park, C.-G., de Cruz Perez, J., & Ratra, B. 2024, <https://arxiv.org/abs/2410.13627>
- Perlmutter, S., et al. 1999, Astrophys. J., 517, 565, doi: [10.1086/307221](https://doi.org/10.1086/307221)
- Riess, A. G., et al. 1998, Astron. J., 116, 1009, doi: [10.1086/300499](https://doi.org/10.1086/300499)
- Riess, A. G., et al. 2018, Astrophys. J., 853, 126, doi: [10.3847/1538-4357/aaa5a9](https://doi.org/10.3847/1538-4357/aaa5a9)
- Riess, A. G., et al. 2022, Astrophys. J. Lett., 934, L7, doi: [10.3847/2041-8213/ac5c5b](https://doi.org/10.3847/2041-8213/ac5c5b)
- Saridakis, E. N., & Sushkov, S. V. 2010, Phys. Rev. D, 81, 083510, doi: [10.1103/PhysRevD.81.083510](https://doi.org/10.1103/PhysRevD.81.083510)
- Scolnic, D. M., et al. 2018, Astrophys. J., 859, 101, doi: [10.3847/1538-4357/aab9bb](https://doi.org/10.3847/1538-4357/aab9bb)
- Seikel, M., Clarkson, C., & Smith, M. 2012, JCAP, 06, 036, doi: [10.1088/1475-7516/2012/06/036](https://doi.org/10.1088/1475-7516/2012/06/036)
- Shafieloo, A., Kim, A. G., & Linder, E. V. 2012, Phys. Rev. D, 85, 123530, doi: [10.1103/PhysRevD.85.123530](https://doi.org/10.1103/PhysRevD.85.123530)

- Shlivko, D., & Steinhardt, P. J. 2024, Phys. Lett. B, 855, 138826, doi: [10.1016/j.physletb.2024.138826](https://doi.org/10.1016/j.physletb.2024.138826)
- Tada, Y., & Terada, T. 2024, Phys. Rev. D, 109, L121305, doi: [10.1103/PhysRevD.109.L121305](https://doi.org/10.1103/PhysRevD.109.L121305)
- Tegmark, M., et al. 2004, Phys. Rev. D, 69, 103501, doi: [10.1103/PhysRevD.69.103501](https://doi.org/10.1103/PhysRevD.69.103501)
- Tsujikawa, S. 2013, Class. Quant. Grav., 30, 214003, doi: [10.1088/0264-9381/30/21/214003](https://doi.org/10.1088/0264-9381/30/21/214003)
- Wang, B., Abdalla, E., Atrio-Barandela, F., & Pavon, D. 2016, Rept. Prog. Phys., 79, 096901, doi: [10.1088/0034-4885/79/9/096901](https://doi.org/10.1088/0034-4885/79/9/096901)
- Wang, D. 2024, <https://arxiv.org/abs/2404.13833>
- Weinberg, S. 2000, in 4th International Symposium on Sources and Detection of Dark Matter in the Universe (DM 2000), 18–26, doi: [10.1007/978-3-662-04587-9_2](https://doi.org/10.1007/978-3-662-04587-9_2)
- Will, C. M. 2014, <https://arxiv.org/abs/1409.7871>
- Yang, T. 2020, Phys. Rev. D, 102, 083511, doi: [10.1103/PhysRevD.102.083511](https://doi.org/10.1103/PhysRevD.102.083511)
- Yang, T., Guo, Z.-K., & Cai, R.-G. 2015, Phys. Rev. D, 91, 123533, doi: [10.1103/PhysRevD.91.123533](https://doi.org/10.1103/PhysRevD.91.123533)
- Yang, W., Pan, S., Di Valentino, E., et al. 2018, JCAP, 09, 019, doi: [10.1088/1475-7516/2018/09/019](https://doi.org/10.1088/1475-7516/2018/09/019)