ANTI-COMMUTATIVE ANTI-ASSOCIATIVE ALGEBRAS. ACAA-ALGEBRAS

ELISABETH REMM

RÉSUMÉ. Let (A, μ) be a nonassociative algebra over a field of characteristic zero. The polarization process allows us to associate two other algebras, and this correspondence is one-one, one commutative, the other anti-commutative. Assume that μ satisfies a quadratic identity $\sum_{\sigma \in \Sigma_3} a_{\sigma} \mu(\mu(x_{\sigma(i)}, x_{\sigma(j)}), x_{\sigma(k)} - a_{\sigma} \mu(x_{\sigma(i)}, \mu(x_{\sigma(j)}, x_{\sigma(k)}) = 0$. Under certain conditions, the polarization of such a multiplication determines an anticommutative multiplication also verifying a quadratic identity. Now only two identities are possible, the first is the Jacobi identity which makes this anticommutative multiplication a Lie algebra and the multiplication μ is Lie admissible, the second, less classical is given by [[x, y], z] = [[y, z], x] = [[z, x], y]. Such a multiplication is here called Acaa for Anticommutative and Antiassociative. We establish some properties of this type of algebras.

TABLE DES MATIÈRES

1.	Definition and basic properties	2
1.1.	Definition	3
1.2.	A characterization of the Acaa-algebras	3
2.	Classification in small dimensions	4
2.1.	Dimension 2	4
2.2.	Dimension 3	4
2.3.	Dimension 4	4
2.4.	Dimension 5	5
2.5.	The free Acaa-algebras	5
3.	Representations of Acaa-algebras	6
3.1.	The operator $\mathrm{ad}X$	6
3.2.	Representations of Acaa algebras	7
3.3.	Acaa Admissible algebra	8
4.	Cohomology and operads	9
4.1.	Cohomology of Acaa algebras	9
4.2.	Quadratic operad associated with Acaa algebras	10

2	2	ς
ſ		
2	_	
С)
2		5
r		
	ę	
	È	
	C	٦.
	÷	<u> </u>
	1	
<	ч	
		۹.
	_	
L.	Г	2
_		
<		
		۹.
p		
6	Y	
<u> </u>		7
	_	
_		2
-	1	3
	-	5
	$\overline{2}$	5
	_	÷
	<u>_</u>	1
		2
	_	٦.
T		-
~	7	-
~		
-		-
-	Λ	

ELISABETH REMM

Références

Introduction The notion of polarization is naturally introduced in the elementary study of functions of a real variable. To such a function we uniquely associate a pair of functions formed by an even function and an odd function whose sum gives the initial function.

We easily extend this concept to the set of bilinear applications on a vector space with values in itself (or in an another space). If φ is such an application, the polarization process associates with it the pair φ_a, φ_c defined by the identities

$$\varphi_a(X,Y) = \varphi(X,Y) - \varphi(Y,X), \quad \varphi_c(X,Y) = \varphi(X,Y) + \varphi(Y,X).$$

The most classic example is to consider on the vector space A an algebra structure defined by a multiplication μ . If we assume that A is an associative algebra, thet is μ satisfies the relation

$$\mu(X, \mu(Y, Z)) = \mu(\mu(X, Y), Z)$$

then μ_c is a Lie bracket, that is satisfies the Jacobi identity

$$\mu(X, \mu(Y, Z)) + \mu(Y, \mu(Z, X)) + \mu(Z, \mu(X, Y)) = 0$$

This work is motivated by the following remark. Assume that φ is an algebra law on A (multiplication defining the structure of the algebra A) that is a nonassociative bilinear map satisfying an identity of the type

(1)
$$\begin{array}{l} a_1(x_1x_2)x_3 + a_2(x_2x_1)x_3 + a_3(x_3x_2)x_1 + a_4(x_1x_3)x_2 + a_5(x_2x_3)x_1 + a_6(x_3x_1)x_2 \\ + b_1x_1(x_2x_3) + b_2x_2(x_1x_3) + b_3x_3(x_2x_1) + b_4x_1(x_3x_2) + b_5x_2(x_3x_1) + b_6x_3(x_1x_2) \end{array} = 0. \end{array}$$

where we write xy in place of $\varphi(x, y)$ Suppose that this identity reduces to an identity relative to φ_a . For example

$$(x_1x_2)x_3 - (x_2x_1)x_3 - (x_3x_2)x_1 - (x_1x_3)x_2 + (x_2x_3)x_1 + (x_3x_1)x_2 - x_1(x_2x_3) + x_2(x_1x_3) + x_3(x_2x_1) + x_1(x_3x_2) + x_2(x_3x_1) + x_3(x_1x_2) = 0.$$

is equivalent to write the Jacobi identity for φ_a (We say in this case that φ or A is Lieadmissible). The question then arises whether there is any other type of relation than the Lie admissibility which is equivalent to a relation concerning only the bracket φ_a . This problem has been addressed in [2]. The following result appears : apart from the Lie admissibility there exists one (and only one) other relation, whose associated bracket verifies

(2)
$$\varphi_c(x_1, \varphi_c(x_2, x_3)) = \varphi_c(x_2, \varphi_c(x_3, x_1)) = \varphi_c(x_3, \varphi_c(x_1, x_2))$$

In this case we shall say that the algebra (A, φ_c) is an anticommutative anti-associative algebra. The antiassociativity [3] is satisfied because

$$\varphi_c(x_1,\varphi_c(x_2,x_3)+\varphi_c(\varphi_c(x_1,x_2),x_3)=\varphi_c(x_1,\varphi_c(x_2,x_3)-\varphi_c(x_3,\varphi_c(x_1,x_2))=0.$$

The goal is to study these algebras that we will call Acaa-algebras.

1. Definition and basic properties

Let \mathbb{K} be a field of characteristic 0. Recall that an algebra (A, μ) over \mathbb{K} (often simply called an algebra) is a \mathbb{K} -vector space A equipped with a bilinear product μ .

1.1. Definition.

Definition 1. We say that the algebra (A, μ) is an Acaa-algebra if the product μ is anticommutatif

$$\mu(x_1, x_2) = -\mu(x_2, x_1)$$

and satisfies the quadratic identity

$$\mu(x_1, \mu(x_2, x_3) = \mu(x_2, \mu(x_3, x_1))$$

for any $x_1, x_2, x_3 \in A$.

To clearly show the anticommutativity of this product we will denote $\mu(x, y)$ by [x, y]. It is clear that we have also

$$[x_1, [x_2, x_3)] = [x_2, [x_3, x_1]] = [x_3, [x_1, x_2]].$$

This product is also antiassociatif [3], because

$$[x_1, [x_2, x_3)] = [x_3, [x_1, x_2]] = -[[x_1, x_2], x_3,].$$

Examples

(3)

- (1) Any abelian algebra ([x, y] = 0 for any $x, y \in A$ is an Acaa-algebra.
- (2) A Lie algebra is an Acaa algebra if and only if the Lie bracket is 2-step nilpotent. In fact if [,] is an Acaa-bracket, the Jacobi condition gives

$$[x_1, [x_2, x_3)] + [x_2, [x_3, x_1]] + [x_3, [x_1, x_2]] = 3[x_1, [x_2, x_3)] = 0$$

This shows that the corresponding Lie algebra is 2-step-nilpotent.

(3) Let us denote by $(\mathcal{F}_{Acaa}(X_1, \dots, X_n), \cdot)$ the free Acaa-algebra with *n*-generators. It is a graded algebra. Let us examine the first particular cases

$$\mathcal{F}_{Acaa}(X_1) = \mathbb{K}\{X\}$$

(4) Let $\mathcal{F}_{Acaa}(X_1, X_2)$ be the free Acaa-algebra with 2 generators X_1, X_2 . Then

$$\mathcal{F}_{Acaa}(X_1, X_2) = \mathbb{K}\{X_1, X_2\} \oplus \mathbb{K}\{[X_1, X_2]\}.$$

(5) Let us consider the 7-dimensional anticommutative algebra whose bracket is given, in a basis $\{e_i, 1 \le i \le 7\}$ by

$$\begin{cases} [e_1, e_2] = e_4, [e_1, e_3] = e_5, [e_2, e_3] = e_6\\ [e_1, e_6] = -[e_2, e_5] = [e_3, e_4] = e_7. \end{cases}$$

It is a Acaa-algebra but not a Lie algebra.

1.2. A characterization of the Acaa-algebras. Assume that (A, [,]) is a Acaa-algebra. By definition

$$[x_1, [x_2, x_3]] = [x_2, [x_3, x_1]] = [x_3, [x_1, x_2]]$$

In particular, if $x_3 = x_1$, we obtain

$$[x_1, [x_2, x_1]] = [x_2, [x_1, x_1]] = 0$$

Conversely, assume that

$$[x_1, [x_2, x_1]] = 0$$

for any $x_1, x_2 \in A$. Then

$$[x_1 + x_3, [x_2, x_1 + x_3]] = 0 = [x_1, [x_2, x_3]] + [x_3, [x_2, x_1]]$$

We deduce

$$[x_1, [x_2, x_3]] = [x_3, [x_1, x_2]]$$

and (A, [,]) is an Acaa-algebra.

Proposition 2. Let (A, [,]) be an anticommutative algebra. Then A is an Acaa-algebra if and only if

$$[x_1, [x_2, x_1]] = 0$$

for any $x_1, x_2 \in A$.

2. Classification in small dimensions

In this section we denote by $\{e_1, \dots, e_n\}$ a basis of A when dim A = n.

2.1. **Dimension** 2. We put $[e_1, e_2] = ae_1 + be_2$. Then

$$[e_1, [e_1, e_2]] = 0 = b[e_1, e_2], \ [e_2, [e_1, e_2]] = 0 = a[e_2, e_1]$$

this implies $[e_1, e_2] = 0$. Any 2-dimensional Acaa-algebra is abelian.

2.2. Dimension 3.

Lemma 3. Let e and f be two independent vectors in A. Then, if [e, f] is not zero, then e, f, [e, f] are independent.

Proof. Assume that $[e, f] \neq 0$ and [e, f] = ae + bf. Then $\begin{bmatrix} e & f \end{bmatrix} = 0 - b[e & f] \quad [f & [e & f] \end{bmatrix} = 0$

$$[e, [e, f]] = 0 = b[e, f], \ [f, [e, f]] = 0 = -a[e, f]$$

this implies a = b = 0 which contradicts the hypothesis.

Assume that A is not abelian. There are two independent vectors, for example e_1 and e_2 such that $[e_1, e_2] \neq 0$. From the lemma, $e_3 = [e_1, e_2]$ is not in $\mathbb{K}\{e_1, e_2\}$ and $\{e_1, e_2, e_3\}$ is a basis of A. We have also

$$[e_1, e_3] = [e_1, [e_1, e_2]] = 0, \ [e_2, e_3] = [e_2, [e_1, e_2]] = 0.$$

In this case A is a Lie algebra isomorphic to the 3-dimensional Heisenberg algebra.

Proposition 4. Every 3-dimensional Acaa-algebra is isomorphic to one of the algebras below

- (1) the 3-dimensional abelian algebra
- (2) the 3-dimensional Heisenberg algebra \mathfrak{h}_3 given by $[e_1, e_2] = e_3$.

2.3. **Dimension** 4. We assume that A is not abelian. Thus we can find a basis $\{e_1, e_2, e_3, e_4\}$ of A such that $\{e_1, e_2, e_3\}$ generates a subalgebra isomorphic to the Heisenberg algebra that is $[e_1, e_2] = e_3$, $[e_i, e_3] = 0$ for i = 1, 2. It remains to compute $[e_i, e_4]$ for i = 1, 2, 3. Let us put $[e_1, e_4] = a_1e_1 + a_2e_2 + a_3e_3 + a_4e_4$. Then $[e_1, [e_1, e_4]] = 0$ gives $b_1e_3 + d_1[e_1, e_4] = 0$ If $d_1 \neq 0$, then $[e_1, e_4] = -(b_1/d_1)e_3$ which contradics $d_1 \neq 0$. Then $d_1 = b_1 = 0$ and $[e_1, e_4] = a_1e_1 + c_1e_2$. Now

$$[e_2, [e_1, e_4] = -a_1e_3 = [e_4, [e_2, e_1]] = [e_3, e_4].$$

We deduce $a_1 = 0$ and $[e_1, e_4] = c_1 e_3$, $[e_3, e_4] = 0$. A similar calculation gives $[e_2, e_4] = c_2 e_3$. In conclusion

$$[e_1, e_2] - e_3, \ [e_1, e_4] = c_1 e_3, \ [e_2, e_4] = c_2 e_3$$

and A is a Lie algebra isomorphic to the direct sum of \mathcal{H}_3 and the 1-dimensional Lie algeba.

Proposition 5. Every 4-dimensional Acaa-algebra is isomorphic to one of the algebras below

- (1) the 4-dimensional abelian algebra
- (2) the 4-dimensional 2-step-nilpotent Lie algebra $\mathcal{H}_3 \oplus \mathbb{K}$ given by $[e_1, e_2] = e_3$.

2.4. **Dimension 5.** As in dimension 4, we assume that A is not abelian. Thus we can find a basis $\{e_1, e_2, e_3, e_4, e_5\}$ of A such that $\{e_1, e_2, e_3\}$ generates a subalgebra isomorphic to the Heisenberg algebra that is $[e_1, e_2] = e_3$, $[e_i, e_3] = 0$ for i = 1, 2. In a first time we assume that $[e_1, e_4] = e_5$. This implies $[e_1, e_5] = [e_4, e_5] = 0$ and

$$[e_2, e_5] = [e_3, e_4] = [e_1, [e_4, e_2]] = ae_3 + be_5.$$

We deduce $[e_2, [e_3, e_4]] = b[e_2, e_5] = [e_4, [e_2, e_3]] = 0$ and b = 0 or $[e_2, e_5] = 0$. If b = 0, then $[e_3, e_4] = ae_3$ this giving a = 0. In any case, $[e_2, e_5] = [e_3, e_4] = 0$ and A is isomorphic to the 2-step nilpotent Lie algebra given by

$$[e_1, e_2] = e_3, [e_1, e_4] = e_5, [e_2, e_4] = ce_3 + de_5$$

by isomorphism we can take c = d = 0. If the hypothesis $[e_1, e_4] = e_5$ is not satisfied, that is $[e_1, e_4] \in \mathbb{K}\{e_1, e_2, e_3, e_4\}$, we can consider that $A_1 = \mathbb{K}\{e_1, e_2, e_3, e_4\}$ is subalgebra of A. Then it is isomorphic to $\mathcal{H}_3 \oplus \mathbb{K}$. In this case, we have to compute $[[e_i, e_5]$ and A, if it is not of the previous case, is isomorphic to \mathcal{H}_5 .

Proposition 6. Every 5-dimensional Acaa-algebra is isomorphic to one of the algebras below

- (1) the 5-dimensional abelian algebra
- (2) the 5-dimensional 2-step-nilpotent Lie algebra $\mathcal{H}_3 \oplus \mathbb{K}^2$ given by $[e_1, e_2] = e_3$.
- (3) the 5-dimensional 2-step-nilpotent Lie algebra defined by

$$[e_1, e_2] = e_3, \ [e_1, e_4] = e_5.$$

(4) the 5-dimensional 2-step-nilpotent Lie algebra \mathcal{H}_5 .

In this case also, any Acaa-algebra is a Lie algebra. In the next section, we shall see a 7-dimensional Acaa-algebra which is not a Lie algebra.

2.5. The free Acaa-algebras. In the first section we have given, as example, the structure of the free Acaa-algebras with 1 or 2-generators. Let us examine the general case. We consider n generators X_1, \dots, X_n and we denote by

$$\mathcal{F}_{Acaa}(X_1,\cdots,X_n)=\oplus_{k\geq 1}\mathcal{F}^k_{Acaa}(X_1,\cdots,X_n)$$

the free Acaa-algebras with n generators, where $\mathcal{F}_{Acaa}^k(X_1, \cdots, X_n)$ denotes the homogeneous component of degree k. We have

(1) $\mathcal{F}^1_{Acaa}(X_1, \cdots, X_n) = \mathbb{K}\{X_1, \cdots, X_n\}$

(2) $\mathcal{F}^{2}_{Acaa}(X_{1}, \cdots, X_{n}) = \mathbb{K}\{X_{ij}, 1 \le i < j \le n\}$ where $X_{ij} = [X_{i}, X_{j}],$

- (3) $\mathcal{F}^{3}_{Acaa}(X_1, \cdots, X_n) = \mathbb{K}\{X_{ijk}, 1 \le i < j < k \le n\}$ where $X_{ijk} = [X_i, [X_j, X_k]],$
- (4) $\mathcal{F}_{A_{caa}}^k(X_1, \cdots, X_n) = 0$ for any $k \ge 4$.

Proof. In fact since $[X_i, [X_j, X_k]] = [X_j, [X_k, X_i]] = [X_k, [X_i, X_j]]$ the components of degree 3 are linear combination of the vectors X_{ijk} with $1 \le i < j < k \le n$. Concerning the degree 4, from [3], any anti-associative algebra is 4-step nilpotent and any product of degree 4 is equal to 0.

Remark We have seen that $\mathcal{F}_{Acaa}(X_1, X_2)$ is a 3-dimensional Lie algebra isomorphic to \mathcal{H}_3 . For n = 3 the generators are $\{X_1, X_2, X_3, X_{12}, X_{13}, X_{23}, X_{123}\}$ and we have

$$\begin{cases} [X_1, X_2] = X_{12}, [X_1, X_3] = X_{13}, [X_2, X_3] = X_{23}, \\ [X_1, X_{23}] = -[X_2, X_{13}] = [X_3, X_{12}] = X_{123}. \end{cases}$$

This algebra is not 2-step nilpotent and it has no Lie algebra structure, for this bracket.

3. Representations of Acaa-Algebras

3.1. The operator adX. Let A be a Acaa-algebra. Let adX be the linear map on A defined by adX(Y) = [X, Y]. Since A is an Acaa-algebra, , for any $X, Y, Z \in A$, we have

[X, [Y, Z]] = [Y, [[Z, X]] = [Z, [X, Y]]

that is

$$[[X, Y], Z] = -[X, [Y, Z]] = [Y, [X, Z]]$$

This is equivalent to

$$\operatorname{ad}[X, Y] = -\operatorname{ad} X \circ \operatorname{ad} Y = \operatorname{ad} Y \circ \operatorname{ad} X.$$

We deduce

Lemma 7. The operators adX on A satisfy

(1) $\operatorname{ad} X \circ \operatorname{ad} Y + \operatorname{ad} Y \circ \operatorname{ad} X = 0$

(2) $2ad[X,Y] = -adX \circ adY + adY \circ adX = -[adX, adY]_{gl(n)}$

where $[,]_{ql(n)}$ denotes the classical bracket on the space of linear operators.

Applications The linear map

$$\operatorname{ad}: A \to End(A)$$

is a weighted linear representation of A. It satisfies

$$\operatorname{ad}[X,Y] = -\frac{1}{2}[\operatorname{ad}X,\operatorname{ad}Y].$$

Since A is also characterized by the relation

$$[X, [X, Y]] = 0$$

then $(adX)^2 = 0$ for all $X \in A$. If we consider ad as a matricial representation of A, the matrices which represent the elements of A are square null matrices.

Along the same lines, we have

$$\mathrm{ad}X[Y,Z] = -[Y,\mathrm{ad}X(Z)] = -[\mathrm{ad}X(Y),Z]$$

So,

$$2ad(X)[Y,Z] = -[Y,adX(Z)] - [adX(Y),Z]$$

In [3], we have introduced the notion of anti-derivation : Let A be an algebra. A linear map $f \in End(A)$ is an anti-derivation if this map satisfies

$$f(xy) = -xf(y) - f(x)y = 0$$

for any $x, y \in A$. As for the representations, we are here led to put coefficients on such a relation :

Definition 8. Let B be an algebra. A linear map $f \in End(B)$ is an weighted anti-derivation if there exists $k \in \mathbb{N}^*$ such that

$$kf(xy) = -xf(y) - f(x)y = 0$$

for any $x, y \in A$.

In these conditions

Lemma 9. Let A be an Acaa algebra. For any $X \in A$, the linear map adX is a weighted anti-derivation of weight 2.

Remark. In [3], we have seen that in any anti-associative algebra, the operator $L_X + R_X$ is an anti-derivation. But in case of anti-commutative anti-associative algebra we have $L_X + R_X = 0$.

3.2. Representations of Acaa algebras. Since an Acaa algebra is a nonassociative algebra, the notion of representation must be taken in the framework of the associative representations of non-associative algebras introduced in [4]. Take for example the map ad. It satisfies

$$\operatorname{ad}[X, Y] = -\operatorname{ad} X \circ \operatorname{ad} Y = \operatorname{ad} Y \circ \operatorname{ad} X.$$

So ad: $A \to End(A)$ is a linear mapping which gives a representation of A in the associative algebra End(A).

Definition 10. Let A be an Acaa algebra and let End(A) be the space of linear endomorphisms of the vector space A. A representation of A is a linear map

$$\rho: A \to End(A)$$

which satisfies the identity

$$\rho[X,Y] = -\rho(X) \circ \rho(Y) = \rho(Y) \circ \rho(X)$$

for any $X, Y \in A$.

The representation ρ is said to be faithful if it is injective. Of course, the linear map ad is an associative representation of A which is not faithful.

Let us note that, if ρ is a representation of A, then $\rho(X)^2 = 0$. Then, identifying End(A) with the space of square matrices of order $n = \dim A$, $\rho(X)$ is considered as a matrix \widetilde{X} satisfying $\widetilde{X}^2 = 0.$

Example : dim 3. The Heisenberg algebra We have seen that any 3-dimensional Acaa algebra is isomorphic to the Heisenberg Lie algebra \mathfrak{h}_3 . It is defined by $[e_1, e_2] = e_3$ and it is isomorphic to the free Acaa algebra with 2 generators. For any $X \in A$, the matrix $\widetilde{X} = \rho(X)$ is 2-nilpotent. Thee is a nonsingular matrix P such that

$$P^{-1}\widetilde{X}P = N_1 = \begin{pmatrix} 0 & 0 & 0\\ 1 & 0 & 0\\ 0 & 0 & 0 \end{pmatrix}$$

Let \widetilde{Y} be a matrix which satisfies $\widetilde{X}\widetilde{Y} = -Y\widetilde{X}$. Then $PN_1P^{-1}\widetilde{Y} = -\widetilde{Y}PN_1P^{-1}$

and

$$N_1 P^{-1} \widetilde{Y} P = -P^{-1} \widetilde{Y} P N_1$$

and this shows that $Y_1 = P^{-1} \widetilde{Y} P$ satisfies $N_1 Y_1 = -Y_1 N_1$. If we compute Y_1 we find

$$Y_1 = \begin{pmatrix} a_{11} & 0 & 0\\ a_{21} & -a_{11} & a_{23}\\ a_{31} & 0 & a_{33} \end{pmatrix}$$

Since $Y_1^2 = 0$, we obtain $a_{11} = a_{33} = a_{23}a_{31} = 0$ and

$$Y_1 = \begin{pmatrix} 0 & 0 & 0 \\ a_{21} & 0 & 0 \\ a_{31} & 0 & 0 \end{pmatrix} \quad \text{or} \quad \begin{pmatrix} 0 & 0 & 0 \\ a_{21} & 0 & a_{23} \\ 0 & 0 & 0 \end{pmatrix}$$

In both cases we have $[\tilde{X}, \tilde{Y}_1] = 0$ and this associative representation is not faithful. We deduce that any associative representation of the Acaa algebra \mathfrak{h}_3 is not faithful. This is not the case for the Lie representations of this Lie algebra.

Example : dim 7. The free Acaa algebra with 3 generators This Acaa algebra is defined in a graded basis $\{e_1, e_2, e_3, e_{12}, e_{13}, e_{23}, e_{123}\}$ by

$$\begin{cases} [e_1, e_2] = e_{12}, [e_1, e_3] = e_{13}, [e_2, e_3] = e_{23} \\ [e_1, e_{23}] = -[e_2, e_{13}] = [e_3, e_{12}] = e_{13} \end{cases}$$

If $X = \sum x_i e_i$, then X is represented by the matrix

3.3. Acaa Admissible algebra. A K-algebra (B, \cdot) is called Acaa-admissible if the bracket on B defined by $[b_1, b_2] = b_1 \cdot b_2 - b_2 \cdot b_1$ is an Acaa-bracket.

Proposition 11. The algebra (B, \cdot) is Acaa-admissible if and only if the product $b_1 \cdot b_2$ satisfies the identity

$$\rho_B(b_1, b_2, b_3) - \rho_B(b_2, b_1, b_3) + \rho_B(b_1, b_3, b_2) - \rho_B(b_3, b_1, b_2) = 0$$

where $\rho_B(b_1, b_2, b_3) = (b_1 \cdot b_2) \cdot b_3 - b_3 \cdot (b_1 \cdot b_2)$. for any $b_1, b_2, b_3 \in B$.

Proof. See [3].

In particular any algebra (B, \cdot) which satisfies the identity

$$\rho_B(b_1, b_2, b_3) = 0$$

is Acaa-admissible. We will call such an algebra a ρ -associative algebra.

Proposition 12. Let B be a ρ -associative algebra. Then the bracket on B given by [x, y] = xy - yx provides B with a Acaa-algebra structure which is also a 2-step nilpotent Lie algebra.

Proof. In fact

$$[[x, y], z] = (xy)z - (yx)z - z(xy) + z(yx) = \rho(x, y, z) - \rho(y, x, z).$$

Now, let $F = \{X_1, \dots, X_n\}$ be a finite set of cardinality n and let $\widetilde{\mathbb{K}}(F)$ the nonassociative free algebra generated by F. Let us consider now the quotient space

$$\widetilde{T}(F) = \widetilde{\mathbb{K}}(F) / \sim$$

where \sim is the equivalence relation given by $(X_iX_j)X_k - X_k(X_iX_j)$. Then $\widetilde{T}(F)$ is a ρ associative algebra. As consequence $\widetilde{T}(F)$ is also an infinite dimensional 2-step nilpotent Lie algebra. For example, if n = 2, this Lie algebra is isomorphic to $\mathfrak{h}_3 \oplus Ab_\infty$ where $Ab - \infty$ is an infinite dimensional abelian Lie algebra. If n = 3, let us consider the 6-dimensional 2-step nilpotent Lie algebra n_6 given by

$$[e_1, e_2] = e_4, \ [e_1, e_3] = e_5, \ [e_2, e_3] = e_6$$

then $\widetilde{T}(F) = n_6 \oplus Ab_{\infty}$. The non abelian part corresponds to the free 2-step nilpotent Lie algebra with 3 generators. This is also true for any $n \geq 3$.

Let A be an Acaa algebra and let us consider the ρ -associative algebra $\widetilde{T}(A)$ (here F means a given basis of A). Let $\widetilde{U}(A)$ be the quotient ring of $\widetilde{T}(A)$ by the ideal generated by elements of the form

$$[X_i, X_j] - X_i X_j + X_j X_i$$

where [,] is the anticommutative multiplication of A. Then $\widetilde{U}(A)$ is also a ρ -associative algebra. It can be also consider as an Acaa algebra which is also a 2-step nilpotent Lie algebra.

Examples

(1) dim A = 1. Then T̃(A) is the graded space given by
T̃¹(A) = K{X}, T̃²(A) = K{X²}, T̃³(A) = K{X³}, T̃⁴(A) = K{X⁴ = X³X, X²X²} and Ũ(F) is abelian.
(2) dim A = 2. Then T̃(A) = T̃(X₁, X₂) is the graded space given by

$$\widetilde{T}^{1}(A) = \mathbb{K}\{X_{1}, X_{2}\}, \ \widetilde{T}^{2}(A) = \mathbb{K}\{X_{1}, X_{2}, X_{1}^{2}, X_{2}^{2}, X_{1}X_{2}, X_{2}X_{1}\}, \cdots$$

and if in $\widetilde{U}(A)$ we put $[X_1, X_2] = X_1X_2 - X_2X_1 = X_{12}$, then $[X_i, X_{12}] = 0$ and $\widetilde{U}(A)$ is isomorphic to the free Acaa algebra with 2-generators.

4. Cohomology and operads

4.1. Cohomology of Acaa algebras. Let A be an Acaa algebra. We saw previously that all products of order 4 were zero. We have a natural filtration

$$\{0\} \subset A^3 \subset A^2 \subset A^1 = A$$

where A^i is the subspace of A generated by the product of degree greater than 3 of elements of A. Let us put $A_i = A^i - A^{i-1}$. This space contains only the product of degree *i*.

For any $X \in A_i$, we consider the linear map g_X defined by

$$g_X(Y) = (-1)^{i+j} ij[X,Y], \quad \forall Y \in A_i.$$

In particular $g_X(Y) = 0$ as soon as $i + j \ge 4$. Now, for every $f \in End(A)$, we put

$$\delta^{1}(f)(u,v) = f[u,v] - [u,f(v)] - [f(u),v]$$

for any $u, v \in A$. In particular we have, for any $X \in A_i, Y \in A_1$ and $Z \in A_1$:

$$\begin{split} \delta^{1}(g_{X})(Y,Z) &= g_{X}[Y,Z] - [Y,g_{X}(Z)] - [g_{X}(Y),Z] \\ &= -2i[X,[Y,Z]] - i[Y,[X,Z]] - i[[X,Y],Z] \\ &= -i(2[X,[Y,Z]] - [Y,[Z,X]] - [Z,[X,Y]] \\ &= -i([X,[Y,Z]] - [Y,[Z,X]]) - i([X,[Y,Z]] - [Z,[X,Y]]) = 0 \end{split}$$

Now, if φ is a skew-symmetric bilinear map on A with values in A, we consider the trilinear map

$$\delta^2(\varphi)(X,Y,Z) = \varphi(X,[Y,Z]) + [X,\varphi(Y,Z)] - \varphi(Y,[Z,X]) - [Y,\varphi(Z,X)].$$

In particular we have

$$\delta^2 \circ \delta^1 = 0.$$

This trilinear map also satisfies

$$\delta^2(\varphi)(X,Y,Z) + \delta^2(\varphi)(Y,Z,X) + \delta^2(\varphi)(Z,X,Y) = 0.$$

Thus, if we want to construct something analogous to Chevalley Eilenberg complex for Lie algebras, we can define the first cochain spaces as follows :

 $\mathcal{C}^{0}(A, A) = A, \ \mathcal{C}^{1}(A, A) = EndA, \ \mathcal{C}^{2}(A, A) = \{\varphi : A \times A \to A, \text{bilinear skew - symmetric}\}$ and $\mathcal{C}^{3}(A, A)$ the space of trilinear maps $\Psi : A \times A \times A \to A$ satisfying the following identity

$$\psi(X, Y, Z) = \psi(Y, X, Z).$$

with

$$\mathcal{C}^{0}(A,A) \to^{\delta^{0}} \mathcal{C}^{1}(A,A)$$
$$\xrightarrow{\delta^{0}} \mathcal{C}^{1}(A,A) \xrightarrow{\delta^{1}} \mathcal{C}^{2}(A,A) \xrightarrow{\delta^{2}} \mathcal{C}^{3}(A,A) \xrightarrow{\delta^{3}} \cdots$$

and

$$\begin{cases} \delta^{3}(\psi)(X_{1}, X_{2}, X_{3}) = & \psi(X_{1}, X_{2}, [X_{3}, X_{4}]) + \psi(X_{1}, [X_{3}, X_{4}], X_{2}) + \psi(X_{2}, [X_{3}, X_{4}], X_{1}) \\ & [X_{1}, \psi(X_{2}, X_{3}, X_{4})] + [X_{1}, (\psi(X_{2}, X_{4}, X_{3})] + [X_{1}, (\psi(X_{4}, X_{3}, X_{2})] \end{cases}$$

4.2. Quadratic operad associated with Acaa algebras. Let $\mathcal{A}caa = \bigoplus_{n \geq 1} \mathcal{A}caa(n)$ be the quadratic operad associated with Acaa algebras. We have

(1) $\mathcal{A}caa(1) = \mathbb{K}$

(2) $\mathcal{A}caa(2) = \mathbb{K}\{x_1x_2\}$ because we have the anticommutativity.

(3) $\mathcal{A}caa(3) = \mathbb{K}\{x_1(x_2x_3)\}.$ in fact

$$x_1(x_2x_3) = x_2(x_3x_1) = x_3(x_1x_2) = -x_1(x_3x_2) = -x_2(x_1x_3) = -x_3(x_2x_1) = -(x_2x_3)x_1 \cdots$$

(4) $\mathcal{A}caa(n) = \{0\}$ for $n \ge 4$ because all the products of degree 4 are null.

The Poincaré or generating series of $\mathcal{A}caa$ is the series

$$g_{\mathcal{A}caa}(t) = \sum \frac{(-1)^n}{n!} \dim(\mathcal{A}\mathcal{A}ss(n))t^n = -t + \frac{1}{2}t^2 - \frac{t^3}{6}$$

Since $\mathcal{A}caa(1) = \mathbb{K}$, $\mathcal{A}caa$ admits a minimal model, unique up to isomorphism. The generating series g_M associated with this minimal model is the formal inverse of $g_{\mathcal{A}caa(t)}$ taken with the opposite sign

$$g_{\mathcal{A}caa}(-g_M(t)) = t.$$

We deduce

$$g_M(t) = -t + \frac{t^2}{2} - \frac{t^3}{3} + \frac{5t^4}{24} - \frac{t^5}{12} - \frac{7t^6}{144} + \cdots$$

and this series cannot be the Poincare Series of the dual operad. In fact, if \mathcal{R} is the ideal of relations which defines $\mathcal{A}caa$ and which is generated as a Σ_3 -module by the relation $(x_1x_2)x_3 - (x_2x_3)x_1$, the ideal \mathcal{R}^{\perp} which generates the dual operad $\mathcal{A}caa^{!}$ is the orthogonal of \mathcal{R} for the classical inner product

(4)
$$\begin{cases} < (x_i \cdot x_j) \cdot x_k, (x_{i'} \cdot x_{j'}) \cdot x_{k'} >= 0, \text{ if } (i, j, k) \neq (i', j', k'), \\ < (x_i \cdot x_j) \cdot x_k, (x_i \cdot x_j) \cdot x_k >= \varepsilon(\sigma), \quad \text{with } \sigma = \begin{pmatrix} 1 & 2 & 3 \\ i & j & k \end{pmatrix} \\ < x_i \cdot (x_j \cdot x_k), x_{i'} \cdot (x_{j'} \cdot x_{k'}) >= 0, \text{ if } (i, j, k) \neq (i', j', k'), \\ < x_i \cdot (x_j \cdot x_k), x_i \cdot (x_j \cdot x_k) >= -\varepsilon(\sigma) \quad \text{with } \sigma = \begin{pmatrix} 1 & 2 & 3 \\ i & j & k \end{pmatrix}, \\ < (x_i \cdot x_j) \cdot x_k, x_{i'} \cdot (x_{j'} \cdot x_{k'}) >= 0, \end{cases}$$

where $\varepsilon(\sigma)$ is the signature of σ . Then $i\mathcal{R}^{\perp}$ is generated as a Σ_3 -module by the relation $(x_1x_2)x_3 + (x_2x_3)x_1$. But this relation is equivalent to $(x_1x_2)x_3 = 0$. In fact In fact any algebra associated with this operad $\mathcal{A}caa^{\dagger}$ have to satisfy

$$(x_1x_2)x_3 + (x_2x_3)x_1 = 0, \ (x_2x_3)x_1 + (x_3x_1)x_2 = 0, \ (x_3x_1)x_2 + (x_1x_2)x_3 = 0$$

and $(x_i x_j) x_k = 0$ for any i, j, k. Then any $\mathcal{A}caa^{!}$ -algebra is a 2-step nilpotent Lie algebra and the generating series of $\mathcal{A}caa^{!}$ is

$$-t+\frac{t^2}{2}.$$

We deduce

Theorem 13. The quadratic operad Acaa is not a Koszul operad.

Références

- [1] Goze M., Remm E., Lie-admissible algebras and operads. J. Algebra 273 (2004), no. 1, 129–152.
- [2] Remm E., Polarization and Depolarization. Preprint Mulhouse 2025.
- [3] Remm, E. Anti-associative algebras. Journal of Algebra and Its Applications. Online Ready (2025) https://doi.org/10.1142/S0219498826500805
- [4] Schafer R.D. Introduction to Nonassociative Algebras. Academic Press. 1966

UNIVERSITÉ DE HAUTE-ALSACE, IRIMAS UR 7499, F-68100 MULHOUSE, FRANCE. Email address: elisabeth.remm@uha.fr