

NOTE ON THE INDUCED METRIC ON POWER SETS OF GROUPS

TATSUHIKO YAGASAKI

ABSTRACT. In this note we clarify general properties of the Hausdorff-like metric on the power sets $\mathcal{S}(G)$ of a group G induced from word length norm and obtain some results on quasi-isometries between some subspaces of $\mathcal{S}(G)$ and $\mathcal{S}(H)$ for a group epimorphism $f : G \rightarrow H$, when $\mathcal{S}(\text{Ker } f)$ is metrically bounded.

CONTENTS

1. Introduction	1
2. Basics on asymmetric metrics and Hausdorff metrics	2
3. Metric on the power set of a group induced from the word length	5
4. The maps on the power sets induced from group homomorphisms	10
5. Word metrics in group actions	17
6. Word length on \ast -sets	20
References	23

1. INTRODUCTION

The study of boundedness of groups aims a quantitative understanding of various norms and metrics on groups or some spaces associated to them (cf. [1, 2, 6, 8, 9, 10, 11], etc). In [6, 7] J. Kędra, A. Libman and B. Martin studied strong and uniform boundedness of finitely normally generated groups based on the word lengths $\| \cdot \|_S$ with respect to finite normally generating sets S on a group. For a study of unbounded groups, in [8] they consider coarse type of Tsuboi metric spaces associated to relatively simple groups and also after [6, 7] Morimichi Kawasaki have studied the metric on the set of finite normally generating sets of a finitely normally generated group modulo symmetrization and conjugation, obtained by symmetrizing the word length norm in [6, 7].

Our goal in this subject is a comprehensive study of boundedness of bundle diffeomorphism groups and diffeomorphism groups of manifold pairs. In [4, 5] we have studied the cases of bundle diffeomorphism groups over a circle and diffeomorphism groups of manifold pairs, in which the submanifold is a finite disjoint union of circles, and are in preparation of successive papers in the higher dimensional cases. As a preliminary of these works, in this note we try to make a framework and a foundation under a general setting and clarify general properties on the extended metric on the power set $\mathcal{S}(G)$ of a group G , which is induced from the word lengths with respect to subsets of G . Here, the power set $\mathcal{S}(G)$ is the set of all subsets of G and we treat the wide class of extended (additive/multiplicative) asymmetric metrics with values in $\mathbb{R}_{\geq 0} \cup \{\infty\}$. Each $S \in \mathcal{S}(G)$ induces the word length ν_S on G and its Hausdorff version $(\nu_S)_H$ on $\mathcal{S}(G)$, from which obtained are the multiplicative/additive asymmetric metrics $\nu(S, T) = (\nu_S)_H(T)$ and $\varrho(S, T) = \log \nu_H(S, T)$ (cf. [6, Section 2]). The usual symmetric metrics $\widehat{\nu}$ and $\widehat{\varrho}$ are readily obtained by symmetrizing ν and ϱ . We list formal properties of these extended metrics. In this general setting,

2020 *Mathematics Subject Classification*. Primary 20F65; Secondary 51F30.

Key words and phrases. Word length, Hausdorff metric, Powerset of group.

one of main problems is to understand relations as metric spaces among (some subspaces of) the power sets $\mathcal{S}(G)$, $\mathcal{S}(H)$ and $\mathcal{S}(K)$ for a group epimorphism $f : G \rightarrow H$ with $K = \text{Ker } f$. We obtain some weak results on quasi-isometries between some subspaces of $\mathcal{S}(G)$ and $\mathcal{S}(H)$ when $\mathcal{S}(K)$ is metrically bounded.

Although the materials in this note are seemed to be essentially known for the authors in [6, 7], we expect that it is useful for our works to clarify general properties of the metrics in $\mathcal{S}(G)$ in this general setting. As an advantage of this framework we can treat various subsets of a group simultaneously within the ambient extended metric space. We also extend this framework to the cases of group actions and sets with binary operations.

2. BASICS ON ASYMMETRIC METRICS AND HAUSDORFF METRICS

This section is devoted to generalities on asymmetric metrics and Hausdorff metrics on powersets.

2.1. Asymmetric metrics.

Let X be a set.

Definition 2.1. For a function $d : X \times X \longrightarrow \mathbb{R}_{\geq 0} \cup \{\infty\}$ consider the following conditions :

- (i) $d(x, x) = 0 \quad (\forall x \in X)$ (i)' $[d(x, y) = d(y, x) = 0 \iff x = y] \quad (\forall x, y \in X)$
- (ii) $d(x, y) = d(y, x) \quad (\forall x, y \in X)$
- (iii) $d(x, z) \leq d(x, y) + d(y, z) \quad (\forall x, y, z \in X)$

We say that the function d is

- 1) an asymmetric metric on X if d satisfies (i) and (iii),
- 2) nondegenerate if it satisfies (i)',
- 3) a psuedo metric on X if it satisfies (i), (ii) and (iii),
- 4) a metric on X if it satisfies (i)', (ii) and (iii).

Definition 2.2. For a function $\varrho : X \times X \longrightarrow \mathbb{R}_{\geq 1} \cup \{\infty\}$ consider the following conditions :

- (i) $\varrho(x, x) = 1 \quad (\forall x \in X)$ (i)' $[\varrho(x, y) = \varrho(y, x) = 1 \iff x = y] \quad (\forall x, y \in X)$
- (ii) $\varrho(x, y) = \varrho(y, x) \quad (\forall x, y \in X)$
- (iii) $\varrho(x, z) \leq \varrho(x, y) \cdot \varrho(y, z) \quad (\forall x, y, z \in X)$

We say that the function ϱ is

- 1) a multiplicative asymmetric metric on X if ϱ satisfies (i) and (iii),
- 2) nondegenerate if it satisfies (i)',
- 3) a multiplicative psuedo metric on X if it satisfies (i), (ii) and (iii),
- 4) a multiplicative metric on X if it satisfies (i)', (ii) and (iii).

Briefly, “an asymmetric metric” is abbreviated as “an as metric”.

For notational simplicity we use the following notations. For any subset $J \subset \mathbb{R}$ and $r \in \mathbb{R}$ we set $J_{\geq r} := \{x \in J \mid x \geq r\}$ and $\tilde{J}_{\geq r} := J_{\geq r} \cup \{\infty\}$. For sets X and Y the symbol $F(X, Y)$ denotes the set of functions from X to Y . For $f \in F(X, \tilde{J}_{\geq r})$ we write $f < \infty$ if f takes its values in $J_{\geq r}$.

Notation 2.1. The notations $\mathcal{M}(X; J)$, $\mathcal{M}_p(X; J)$ and $\mathcal{M}_{as}(X; J)$ denote the sets of metrics, psuedo metrics and as metrics on a set X with values in $J \subset \tilde{\mathbb{R}}_{\geq 0}$, respectively. Similarly, $\mathcal{M}_m(X; I)$, $\mathcal{M}_{m,p}(X; I)$ and $\mathcal{M}_{m,as}(X; I)$ denote the sets of multiplicative metrics, psuedo metrics and as metrics a set X with

values in $J \subset \widetilde{\mathbb{R}}_{\geq 1}$, respectively. The symbols J and I are omitted when $J = \widetilde{\mathbb{R}}_{\geq 0}$ and $I = \widetilde{\mathbb{R}}_{\geq 1}$. For $d \in \mathcal{M}_{as}(X)$ its restriction to $A \subset X$, $d|_{A \times A} \in \mathcal{M}_{as}(A)$, is simply denoted by $d|_A$.

In this note we work in the set $\mathcal{M}_{as}(X)$. Since $d \in \mathcal{M}_{as}(X)$ may be degenerate or non-symmetric and may take the value ∞ , we list so basic properties in order to remind and confirm them, while their proofs are omitted since those follows directly from their definitions and routine arguments.

Notation 2.2. Let $d \in F(X^2, \widetilde{\mathbb{R}}_{\geq 0})$. The symmetrization of d is defined by

$$\widehat{d} : X \times X \longrightarrow \widetilde{\mathbb{R}}_{\geq 0} : \widehat{d}(x, y) = \max\{d(x, y), d(y, x)\}.$$

If $d \in \mathcal{M}_{as}(X)$, then (i) $\widehat{d} \in \mathcal{M}_p(X)$ and (ii) $\widehat{d} \in \mathcal{M}(X)$ if and only if d is nondegenerate.

We recall some notions on (quasi-)isometries.

Notation 2.3. Suppose $f : (X, d) \rightarrow (Y, \varrho)$ is a map between (additive/multiplicative) as metric spaces. We use the following terminologies.

- (1) (i) f is isometric if $\varrho(f(x), f(x')) = d(x, x')$ ($\forall x, x' \in X$).
If f is isometric and d is nondegenerate, then f is injective.
- (ii) f is an isometry if f is an isometric bijection.
- (2) f is a quasi-isometric map, if it satisfies the following condition :
 $(\dagger) \quad \varrho(f(x), f(x')) \leq \lambda d(x, x') + C \quad (\forall x, x' \in X) \quad \text{for some } \lambda \in \mathbb{R}_{\geq 1} \text{ and } C \in \mathbb{R}_{\geq 0}.$
 \circ When d and ϱ are multiplicative as metrics, the condition (\dagger) is replaced as follows :
 $(\ddagger) \quad \varrho(f(x), f(x')) \leq D d(x, x')^\lambda \quad (\forall x, x' \in X) \quad \text{for some } \lambda \in \mathbb{R}_{\geq 1} \text{ and } D \in \mathbb{R}_{\geq 1}.$
- (3) Let $r \in \mathbb{R}_{\geq 0}$. Two maps $h, h' : X \rightarrow (Y, \varrho)$ are r -close if $\widehat{\varrho}(h(x), h'(x)) \leq r$ ($\forall x \in X$).
- (4) f is a quasi-isometry if f is a quasi-isometric map and there exists a quasi-isometric map $g : (Y, \varrho) \rightarrow (X, d)$ such that gf is r -close to id_X and fg is r -close to id_Y for some $r \in \mathbb{R}_{\geq 0}$.

Notation 2.4. Let W be a set and $f, g \in F(W, \widetilde{\mathbb{R}}_{\geq 0})$.

- (1) $f \leq_q g \iff f \leq \lambda g + C$ for some $\lambda \in \mathbb{R}_{\geq 1}$ and $C \in \mathbb{R}_{\geq 0}$.
- (2) $f \sim_q g$ (quasi-equivalent) $\iff f \leq_q g$ and $g \leq_q f$

Note that for $d, d' \in \mathcal{M}(X)$, $d \sim_q d'$ if and only if $\text{id}_X : (X, d) \rightarrow (X, d')$ is a quasi-isometry.

We can introduce a natural metric to the set $\mathcal{M}_{as}(X) \subset F(X^2, \widetilde{\mathbb{R}}_{\geq 0})$.

Notation 2.5. Let W be any set.

- (1) The set $F(W, \widetilde{\mathbb{R}}_{\geq 0})$ has the following nondegenerate as metric.

$$\mu(f, f') := \begin{cases} \inf\{r \in \mathbb{R}_{\geq 0} \mid f' \leq f + r\} & (\exists r \in \mathbb{R}_{\geq 0} \text{ s.t. } f' \leq f + r) \\ \infty & (\text{otherwise}) \end{cases}$$

- (2) The multiplicative version of μ is defined by

$$\lambda : F(W, \widetilde{\mathbb{R}}_{\geq 1})^2 \longrightarrow \widetilde{\mathbb{R}}_{\geq 1} : \lambda(g, g') := \begin{cases} \inf\{s \in \mathbb{R}_{\geq 1} \mid g' \leq sg\} & (\exists s \in \mathbb{R}_{\geq 1} \text{ s.t. } g' \leq sg) \\ \infty & (\text{otherwise}) \end{cases}$$

- (3) The additive as metric μ and the multiplicative as metric λ correspond under the reciprocal bijections :

$$\begin{array}{ccc} \exp_* : F(W, \widetilde{\mathbb{R}}_{\geq 0}) & \longrightarrow & F(W, \widetilde{\mathbb{R}}_{\geq 1}) \\ f & \xrightarrow{\exp f} & \end{array} \quad \begin{array}{ccc} \log_* : F(W, \widetilde{\mathbb{R}}_{\geq 1}) & \longrightarrow & F(W, \widetilde{\mathbb{R}}_{\geq 0}) \\ g & \xrightarrow{\log g} & \end{array}$$

$$(i) \quad \mu = \log \lambda (\exp_* \times \exp_*)$$

$$(ii) \quad \text{For } f, f' \in F(W, \widetilde{\mathbb{R}}_{\geq 0}) \text{ and } r \in \mathbb{R}_{\geq 0}$$

$$\mu(f, f') \leq r \iff f' \leq f + r \iff e^{f'} \leq e^r e^f \iff \lambda(e^f, e^{f'}) \leq e^r$$

Definition 2.3. The set $\mathcal{M}_{as}(X)$ inherits the nondegenerate as metric $\mu|_{\mathcal{M}_{as}(X)}$ from $(F(X^2, \widetilde{\mathbb{R}}_{\geq 0}), \mu)$.

$$(i) \quad \mu(d, d') \leq r \iff d' \leq d + r \quad (d, d' \in \mathcal{M}_{as}(X), r \in \mathbb{R}_{\geq 0})$$

Notation 2.6. Suppose $(X, d), (Y, \varrho)$ are as metric spaces.

$$(1) \quad \text{We have an as metric } \mu \in \mathcal{M}_{as}(X \times Y) : \mu((x, y), (x', y')) = \max\{d(x, x'), \varrho(y, y')\}.$$

If d and ϱ are nondegenerate, then μ is also nondegenerate.

2.2. Hausdorff metrics on the power sets.

Next we recall basic notions on Hausdorff metrics. Let X be a set. The symbol $\mathcal{S}(X)$ denote the power set of X , that is, the set of all subsets of X . We also set $\mathcal{S}_f(X) := \{S \in \mathcal{S}(X) \mid |S| < \infty\}$. We regard as $X \subset \mathcal{S}(X)$ by identifying each point $x \in X$ with the singleton $\{x\} \in \mathcal{S}(X)$. Each $d \in \mathcal{M}_{as}(X)$ determines the Hausdorff metric $d_H \in \mathcal{M}_{as}(\mathcal{S}(X))$. The r balls and r neighborhoods with respect to d are denoted by the following notations.

Notation 2.7. Let $d \in \mathcal{M}_{as}(X)$, $x \in X$ and $A \subset X$.

$$(1) \quad B_d(x, r) := \{y \in X \mid d(x, y) \leq r\} \quad B_d(A, r) := \bigcup_{a \in A} B_d(a, r) \quad (r \in \mathbb{R}_{\geq 0})$$

$$(2) \quad B_d(x, \infty) := \bigcup_{r \in \mathbb{R}_{\geq 0}} B_d(x, r) \quad B_d(A, \infty) := \bigcup_{a \in A} B_d(a, \infty) = \bigcup_{r \in \mathbb{R}_{\geq 0}} B_d(A, r)$$

We omit the symbol d when it is understood from the context.

Fact 2.1.

$$(1) \quad x \in B(x, 0), \quad A \subset B(A, 0) \quad (2) \quad A \subset A' \implies B(A, r) \subset B(A', r) \quad (r \in \widetilde{\mathbb{R}}_{\geq 0})$$

$$(3) \quad B(B(A, r), s) \subset B(A, r + s) \quad (r, s \in \widetilde{\mathbb{R}}_{\geq 0})$$

Definition 2.4. Each $d \in \mathcal{M}_{as}(X)$ induces an as metric $d_H \in \mathcal{M}_{as}(\mathcal{S}(X))$ defined as follows.

$$(1) \quad d_H : \mathcal{S}(X) \times \mathcal{S}(X) \longrightarrow \widetilde{\mathbb{R}}_{\geq 0} :$$

$$d_H(A, A') := \begin{cases} \inf\{r \in \mathbb{R}_{\geq 0} \mid A' \subset B_d(A, r)\} & (\exists r \in \mathbb{R}_{\geq 0} \text{ s.t. } A' \subset B_d(A, r)) \\ \infty & (\text{otherwise}) \end{cases}$$

$$(2) \quad B_H(A, r) := \{x \in X \mid d_H(A, x) \leq r\} \quad B_H(A, \infty) := \bigcup_{r \in \mathbb{R}_{\geq 0}} B_H(A, r)$$

$$(3) \quad A \subset X : d\text{-closed} \iff B_H(A, 0) = A \quad \mathcal{F} \equiv \mathcal{F}(X, d) := \text{the set of } d\text{-closed subsets of } X$$

Note that $B_H(A, r) \subset B_{d_H}(A, r)$.

Fact 2.2. The function $d_H \in F(\mathcal{S}(X)^2, \widetilde{\mathbb{R}}_{\geq 0})$ has the following properties. Let $A, A', A'' \in \mathcal{S}(X)$.

$$(1) \quad (i) \quad A' \subset A'' \implies d_H(A, A') \leq d_H(A, A'') \quad (ii) \quad d_H(x, y) = d(x, y) \quad (\forall x, y \in X)$$

$$(3) \quad \text{For } r, s \in \mathbb{R}_{\geq 0} \text{ the following holds.}$$

$$(i) \quad d_H(A, A') \leq r \iff A' \subset B(A, s) \quad (\forall s \in \mathbb{R}_{>r})$$

$$(ii) \quad B(A, r) \subset B_H(A, r) \subset B(A, s) \quad (\forall s \in \mathbb{R}_{>r}) \quad B_H(A, \infty) = B(A, \infty)$$

$$(iii) \quad d_H(A, A') \leq r \iff d_H(A, a') \leq r \quad (\forall a' \in A') \iff A' \subset B_H(A, r)$$

$$(iv) \quad d_H(A, B_H(A, r)) \leq r \quad (v) \quad B_H(B_H(A, r), s) \subset B_H(A, r + s)$$

- (2) $d_H(A, A'') \leq d_H(A, A') + d_H(A', A'')$
- (3) (i) $d_H(A, A') = 0 \iff A' \subset B_H(A, 0) \iff B_H(A', 0) \subset B_H(A, 0)$
(ii) $d_H(A, A) = 0$ (iii) $d_H(A, A') = 0 = d_H(A', A) \iff B_H(A, 0) = B_H(A', 0)$
- (4) (i) $d_H \in \mathcal{M}_{as}(\mathcal{S}(X))$ and \widehat{d}_H is a pseudo-metric.
(ii) $d_H|_{\mathcal{F}} \in \mathcal{M}_{as}(\mathcal{F})$ is nondegenerate, so that $\widehat{d}_H|_{\mathcal{F}} = \widehat{d_H|_{\mathcal{F}}} \in \mathcal{M}(\mathcal{F})$, where $\mathcal{F} \equiv \mathcal{F}(X, d)$.

Fact 2.3. Let $d, d' \in \mathcal{M}_{as}(X)$. The symbols B, B_H and B', B'_H are determined by d and d' respectively.

- (1) If $d' \leq \lambda d + C$ for $\lambda \in \mathbb{R}_{\geq 1}$ and $C \in \mathbb{R}_{\geq 0}$, then
(i) $B_H(A, r) \subset B'_H(A, \lambda r + C)$ ($\forall r \in \widetilde{\mathbb{R}}_{\geq 0}$) (ii) $d'_H \leq \lambda d_H + C$
- (2) (i) $d \leq_q d' \implies d_H \leq_q d'_H$ (ii) $d \sim_q d' \implies d_H \sim_q d'_H \implies \widehat{d}_H \sim_q \widehat{d}'_H$

In the case where $d \in \mathcal{M}_{as}(X)$ takes discrete values, we do not need to distinguish between $B(A, r)$ and $B_H(A, r)$.

Fact 2.4. Suppose J is a discrete subset of $\mathbb{R}_{\geq 0}$ and $d \in \mathcal{M}_{as}(X, \widetilde{J})$. Then, the following holds.

- (1) (i) $0 \in J$ (if $X \neq \emptyset$), since $d(x, x) = 0$ for any $x \in X$.
(ii) For any $r \in \mathbb{R}_{\geq 0}$ there exists $\lfloor r \rfloor_J := \max J_{\leq r}$, since $J_{\leq r} = \{k \in J \mid k \leq r\}$ is a nonempty finite set of J .

Below, let $r \in \mathbb{R}_{\geq 0}$ and put $n := \lfloor r \rfloor_J$.

- (2) (i) $B(A, r) = B(A, n)$ (ii) $d_H(A, A') \leq r \iff d_H(A, A') \leq n$ ($A' \subset X$)
- (3) $d_H \in \mathcal{M}_{as}(\mathcal{S}(X), \widetilde{J})$: $d_H(A, A') := \begin{cases} \min\{k \in J \mid A' \subset B(A, k)\} & (\exists r \in \mathbb{R}_{\geq 0} \text{ s.t. } A' \subset B(A, r)) \\ \infty & (\text{otherwise}) \end{cases}$
- (4) (i) $B_H(A, r) = B_H(A, n)$ (ii) $B_H(A, r) = B(A, r)$

3. METRIC ON THE POWER SET OF A GROUP INDUCED FROM THE WORD LENGTH

3.1. Basics on operations for subsets of a group.

Suppose G is a group and $e \in G$ is the identity element. Let $G^\times := G - \{e\}$.

Definition 3.1.

- (1) A norm on G means a function $\nu : G \rightarrow \widetilde{\mathbb{R}}_{\geq 0}$ which satisfies the following conditions :
(i) $\nu(x) = 0 \iff x = e$
(ii) $\nu(x^{-1}) = \nu(x)$ ($\forall x \in G$)
(iii) $\nu(xy) \leq \nu(x) + \nu(y)$ ($\forall x, y \in G$)
- (2) A function $\nu : G \rightarrow \widetilde{\mathbb{R}}_{\geq 0}$ is conjugation-invariant if it satisfies the following condition :
(iv) $\nu(axa^{-1}) = \nu(x)$ ($\forall x, a \in G$)

Notation 3.1. For $S, T \subset G$ we use the following notations.

- (1) $ST := \{xy \mid x \in S, y \in T\} \subset G$
- (2) $S^n := \{x_1 \cdots x_n \mid x_i \in S \text{ (} i \in [n]\text{)}\}$ ($n \in \mathbb{Z}_{\geq 1}$) $S^0 := \{e\}$ $S^{-1} = \{x^{-1} \mid x \in S\}$
- (3) $S^{\leq n} = \bigcup_{k=0}^n S^k$ ($n \in \mathbb{Z}_{\geq 1}$) $S^\infty = \bigcup_{n=0}^\infty S^n$
- (4) (the conjugations)
(i) $x^a := a^{-1}xa$, ${}_ax := axa^{-1}$ ($x, a \in G$) (ii) $S^A := \{x^a \mid x \in S, a \in A\}$ ($A \subset G$)

Notation 3.2. For $S \subset G$ we use the following notations.

- (1) (i) $\langle S \rangle$: the subgroup of G generated by S
(ii) $\langle\langle S \rangle\rangle$: the normal subgroup of G normally generated by S
- (2) $S^\pm := S \cup S^{-1}$ (the symmetrization of S)
- (3) (i) $C(S) := S^G$ (the conjugation closure of S)
(ii) $C_S := (S^\pm)^G = (S^G)^\pm$ (the symmetrized conjugation closure of S)
- (4) (i) S is symmetric. $\stackrel{\text{def}}{\iff} S = S^{-1}$
(ii) S is conjugation-invariant. $\stackrel{\text{def}}{\iff} S^a \subset S \ (\forall a \in G) \stackrel{\text{equiv}}{\iff} S^a = S \ (\forall a \in G)$

Notation 3.3. For subsets of $\mathcal{S}(G)$ we use the following notations.

[I] For any condition P on subsets S of G , we set $\mathcal{S}_P(G) := \{S \in \mathcal{S}(G) \mid S : P\}$.

Examples of a condition P :

- (1) (i) $f : |S| < \infty$ (a finite set) (ii) $g : S^\infty = G$ (iii) $s : S^{-1} = S$ (symmetric)
- (2) (i) $c : S^G = S$ (conjugation-invariant) $\iff S = A^G \ (\exists A \in \mathcal{S}(G))$
(ii) $fc : S = A^G \ (\exists A \in \mathcal{S}_f(G))$ (iii) $s, c := s + c \iff S = C_A \ (\exists A \in \mathcal{S}(G))$
- (3) (i) $fg := g + s + f$ (ii) $ng := g + s + c$
(iii) $nfg := g + s + fc \iff S^\infty = G, S = C_A \ (\exists A \in \mathcal{S}_f(G))$

[II] To avoid the degeneracy of induced metrics on $\mathcal{S}(G)$ (cf. Fact 3.3(1)), we need some care on the identity element e . For this reason we introduce the following notations.

- (1) For $\mathcal{T} \subset \mathcal{S}(G)$ we set $\mathcal{T}^* := \mathcal{T} - \{\emptyset, \{e\}\}$.
- (2) $\mathcal{S}(G) = \mathcal{S}'(G) \amalg \mathcal{S}''(G) : \mathcal{S}'(G) := \{S \in \mathcal{S}(G) \mid e \notin S\}, \mathcal{S}''(G) := \{S \in \mathcal{S}(G) \mid e \in S\}$
(i) $\mathcal{S}'_P(G) := \mathcal{S}'(G) \cap \mathcal{S}_P(G), \mathcal{S}''_P(G) := \mathcal{S}''(G) \cap \mathcal{S}_P(G)$
- (3) $\tilde{\mathcal{S}}(G) := \mathcal{S}(G) / \sim :$
(i) \sim is the equivalence relation on $\mathcal{S}(G)$ generated by $S \sim S \cup \{e\} \ (S \in \mathcal{S}(G))$.
(ii) $S \sim T \iff S - \{e\} = T - \{e\} \ (S, T \in \mathcal{S}(G))$
(iii) The equivalence class of $S \in \mathcal{S}(G)$ is denoted by $[S]$.
(iv) $\tilde{\mathcal{S}}_P(G) := \{[S] \in \tilde{\mathcal{S}}(G) \mid S \in \mathcal{S}_P(G)\}$
- (4) There exist canonical bijections $\mathcal{S}'(G) \approx \mathcal{S}''(G) \approx \tilde{\mathcal{S}}(G)$

$$S \quad S \cup \{e\} \quad [S] = \{S, S \cup \{e\}\}.$$

Below we prefer the subset $\mathcal{S}''(G)$ since related expressions take a simple form (cf. Fact 3.1(4)(iv)). Next we list up basic properties of subsets of G under various operations.

Fact 3.1. Let $G \supset S, S_1, S_2, T, T_1, T_2, U, S_\lambda, T_\mu \ (\lambda \in \Lambda, \mu \in M)$ and $G \ni a, b$.

- (1) (i) $S_1 \subset S_2, T_1 \subset T_2 \implies S_1 T_1 \subset S_2 T_2, S_1^{T_1} \subset S_2^{T_2}$ (ii) $eS = S = Se$
(iii) $(ST)U = S(TU)$ (iv) $(S^T)^U = S^{TU}$
(v) $(S \cup T)^{-1} = S^{-1} \cup T^{-1} \quad (ST)^{-1} = T^{-1} S^{-1} \quad (S^T)^{-1} = (S^{-1})^T$
(vi) $TU \cup TU^{-1} \subset T \implies S^T U = U S^T$
 $(\because) S^T U \ni x^a y = y y^{-1} x^a y = y(x^a)^y = y x^{ay} \in U S^T$
 $U S^T \ni y x^a = y x^a y^{-1} y = (x^a)^{y^{-1}} y = x^{ay^{-1}} y \in S^T U$
- (2) $(S \cup T)^n = \bigcup_{I+J=[n]} S^I T^J \quad (S^I T^J = X_1 \cdots X_n : X_k = S \text{ if } k \in I, X_k = T \text{ if } k \in J)$
(i) $ST = TS \implies (S \cup T)^n = \bigcup \{S^k T^\ell \mid k, \ell \in \mathbb{Z}_{\geq 0}, k + \ell = n\} \quad (ST)^n = S^n T^n$

- (ii) $(S \cup T)^n = S^n \cup S^{\leq n-1}T$ if $T \triangleleft G$. $(S^{\leq -1} := \emptyset)$
- (3) (i) $S = \bigcup_{\lambda \in \Lambda} S_\lambda$, $T = \bigcup_{\mu \in M} T_\mu$
 $\implies ST = \bigcup_{(\lambda, \mu) \in \Lambda \times M} S_\lambda T_\mu$, $S^T = \bigcup_{(\lambda, \mu) \in \Lambda \times M} S_\lambda^{T_\mu}$, $S^{-1} = \bigcup_{\lambda \in \Lambda} S_\lambda^{-1}$
(ii) $S = \bigcap_{\lambda \in \Lambda} S_\lambda \implies S^{-1} = \bigcap_{\lambda \in \Lambda} S_\lambda^{-1}$, $S^T \subset \bigcap_{\lambda \in \Lambda} S_\lambda^T$
- (4) (i) $S^k S^\ell = S^{k+\ell}$ $(S^k)^\ell = S^{k\ell}$ $(k, \ell \in \mathbb{Z}_{\geq 0})$ $S^\infty S^\infty = S^\infty$
(ii) $(S^n)^{-1} = (S^{-1})^n$ $(S^{\leq n})^{-1} = (S^{-1})^{\leq n}$ $(n \in \tilde{\mathbb{Z}}_{\geq 0})$
(iii) $S^{\leq m} S^{\leq n} = S^{\leq m+n}$ $(S^{\leq m})^{\leq n} \subset S^{\leq mn}$ $(m, n \in \mathbb{Z}_{\geq 0})$
(iv) $e \in S \implies S^n = S^{\leq n}$ $(n \in \mathbb{Z}_{\geq 0})$ (v) $(S \cup \{e\})^n = (S \cup \{e\})^{\leq n} = S^{\leq n}$ $(n \in \mathbb{Z}_{\geq 0})$
(vi) $(x^a)^b = x^{ab}$ $(xy)^a = x^a y^a$ $(x^{-1})^a = (x^a)^{-1}$ $(S \cup T)^a = S^a \cup T^a$ $(S \cap T)^a = S^a \cap T^a$
 $(ST)^a = S^a T^a$ $(S^a)^b = S^{ab}$ $(S^{-1})^a = (S^a)^{-1}$
 $(S^n)^a = (S^a)^n$ $(S^{\leq n})^a = (S^a)^{\leq n}$ $(S^{\leq n})^T \subset (S^T)^{\leq n}$ $(n \in \tilde{\mathbb{Z}}_{\geq 0})$
- (5) (i) S^\pm is symmetric. $(S \cup T)^\pm = S^\pm \cup T^\pm$
(ii) S, T are symmetric. $\implies S^{-1}, S \cup T, S \cap T$ are symmetric. $(ST)^{-1} = TS$
(iii) S_λ is symmetric $(\forall \lambda \in \Lambda)$. $\implies \bigcap_{\lambda \in \Lambda} S_\lambda$ is symmetric.
- (6) (i) ${}^T S = S^{T^{-1}}$ (ii) $ST \subset TST$, $TS \subset {}^T S T = S^{T^{-1}} T$
(iii) S is conjugation-invariant. $\implies {}^T S = S^T = S$, $ST = TS$
(iv) S, T are conjugation-invariant. $\implies S^{-1}, S \cup T, S \cap T, ST$ are conjugation-invariant.
(v) S_λ is conjugation-invariant $(\forall \lambda \in \Lambda)$. $\implies S = \bigcap_{\lambda \in \Lambda} S_\lambda$ is conjugation-invariant.
(vi) S, T are symmetric, conjugation-invariant.
 $\implies S^{-1}, S \cup T, ST$ are symmetric, conjugation-invariant.
- (7) (i) $C(S)$ is conjugation-invariant. (ii) C_S is symmetric and conjugation-invariant.
(iii) $C_{S \cup T} = C_S \cup C_T$ $C_{S \cup \{e\}} = C_S \cup \{e\}$ (iv) $S \subset G^\times \implies C_S \subset G^\times$
- (8) (i) S^∞ is a submonoid of G generated by S .
(ii) $\langle S \rangle = \langle S^\pm \rangle = (S^\pm)^\infty$ (iii) $\langle \langle S \rangle \rangle = \langle C(S) \rangle = (C_S)^\infty$ $\circ \emptyset^0 = \emptyset^\infty = \langle \emptyset \rangle = \langle \langle \emptyset \rangle \rangle = \{e\}$

3.2. Metrics on the power set of a group.

Definition 3.2. Let $S \in \mathcal{S}(G)$.

- (1) the word metric on G with respect to S :

$$d_S : G \times G \longrightarrow \tilde{\mathbb{Z}}_{\geq 0} : d_S(x, y) := \begin{cases} \min\{n \in \mathbb{Z}_{\geq 0} \mid y \in xS^n\} & (y \in xS^\infty) \\ \infty & (x \in G - xS^\infty) \end{cases}$$

- (2) the Hausdorff metric induced from d_S :

$$(d_S)_H : \mathcal{S}(G) \times \mathcal{S}(G) \longrightarrow \tilde{\mathbb{Z}}_{\geq 0} :$$

$$(d_S)_H(A, A') := \begin{cases} \min\{n \in \mathbb{Z}_{\geq 0} \mid A' \subset AS^{\leq n}\} & (\exists n \in \mathbb{Z}_{\geq 0} \text{ s.t. } A' \subset AS^{\leq n}) \\ \infty & (\text{otherwise}) \end{cases}$$

- (i) For $n \in \mathbb{Z}_{\geq 0}$ $(d_S)_H(A, A') \leq n \iff A' \subset AS^{\leq n}$

Definition 3.3. Let $S \in \mathcal{S}(G)$.

- (1) the word length norm on G with respect to S :

$$\nu_S := d_S(e, *) : G \longrightarrow \tilde{\mathbb{Z}}_{\geq 0} : \nu_S(x) = d_S(e, x) = \begin{cases} \min\{n \in \mathbb{Z}_{\geq 0} \mid x \in S^n\} & (x \in S^\infty) \\ \infty & (x \in G - S^\infty) \end{cases}$$

- (i) For $n \in \mathbb{Z}_{\geq 0}$ $\nu_S(x) \leq n \iff x \in S^{\leq n}$
- (2) the Hausdorff-like extension of ν_S :
- $$(\nu_S)_H := (d_S)_H(e, *) : \mathcal{S}(G) \longrightarrow \tilde{\mathbb{Z}}_{\geq 0} : (\nu_S)_H(A) = (d_S)_H(e, A) = \sup \nu_S(A)$$
- (sup is taken in $\tilde{\mathbb{R}}_{\geq 0}$ so that $\sup \emptyset = 0$.)
- $$\varrho_S := \log (\nu_S)_H : \mathcal{S}(G)^* \longrightarrow \log (\tilde{\mathbb{Z}}_{\geq 1}) \subset \tilde{\mathbb{R}}_{\geq 0}$$
- (i) For $n \in \mathbb{Z}_{\geq 0}$ $(\nu_S)_H(A) \leq n \iff A \subset S^{\leq n}$
- (ii) $(\nu_S)_H(A) = 0 \iff A \subset \{e\} \iff A = \{e\}$ or \emptyset
- o If there is no ambiguity we omit the subscript H and write $(\nu_S)_H$ simply as ν_S .

Definition 3.4.

- (1) $\nu_H : \mathcal{S}(G) \times \mathcal{S}(G) \longrightarrow \tilde{\mathbb{Z}}_{\geq 0} : \nu_H(S, T) := (\nu_S)_H(T)$
- $$\varrho := \log \nu_H : \mathcal{S}(G)^* \times \mathcal{S}(G)^* \longrightarrow \tilde{J} \subset \tilde{\mathbb{R}}_{\geq 0} : \varrho(S, T) = \varrho_S(T)$$
- (2) $\widehat{\varrho} := \log \widehat{\nu}_H : \mathcal{S}(G)^* \times \mathcal{S}(G)^* \longrightarrow \tilde{J} \subset \tilde{\mathbb{R}}_{\geq 0}$
- $$\widehat{\varrho}(S, T) = \max \{ \varrho_S(T), \varrho_T(S) \} = \log \max \{ (\nu_S)_H(T), (\nu_T)_H(S) \} = \log \sup (\nu_S(T) \cup \nu_T(S))$$

Fact 3.2. Let $S \in \mathcal{S}(G)$.

- (1) $d_S \in \mathcal{M}_{as}(G, \tilde{\mathbb{Z}}_{\geq 0})$
- (i) $d_S(x, y) = 0 \iff x = y$ (ii) $d_S(x, y) = d_{S^{-1}}(y, x)$ ($\forall x, y \in G$)
- (iii) $d_S(ax, ay) = d_S(x, y)$ ($\forall a, x, y \in G$) (iv) $d_{S^a}(x^a, y^a) = d_S(x, y)$ ($\forall a, x, y \in G$)
- (2) ν_S satisfies the conditions (i), (iii) in Definition 3.1
- (i) $d_S(x, y) = \nu_S(x^{-1}y)$ ($\forall x, y \in G$) (ii) $\nu_{S^a}(x^a) = \nu_S(x)$ ($\forall a, x \in G$)
- (iii) $\nu_S^{-1}(0) = \{e\}$, $\nu_S^{-1}(1) = S - \{e\}$
- (3) If $S = S^{-1}$, then (i) $d_S \in \mathcal{M}(G, \tilde{\mathbb{Z}}_{\geq 0})$ (ii) ν_S is a norm on G .
- (4) If S is conjugation invariant, then d_S and ν_S are conjugate-invariant.

Fact 3.3.

- (1) Let $S, T, A \subset G$.
- (i) $(\nu_S)_H(A) = 1 \iff A \not\subset \{e\}$ and $A \subset S \cup \{e\}$
- (ii) $(\nu_S)_H(T) = (\nu_T)_H(S) = 1 \iff S - \{e\} = T - \{e\} \neq \emptyset$
- (iii) $(\nu_S)_H(A \cup \{e\}) = (\nu_S)_H(A)$, $(\nu_{S \cup \{e\}})_H(A) = (\nu_S)_H(A)$
- (2) $\nu_H \in \mathcal{M}_{m,as}(\mathcal{S}(G)^*)$ and $\varrho \in \mathcal{M}_{as}(\mathcal{S}(G)^*)$.
- (3) $\widehat{\varrho} \in \mathcal{M}(\mathcal{S}''(G)^*)$
- (4) ν_H induces the function : $\tilde{\nu}_H : \tilde{\mathcal{S}}(G) \times \tilde{\mathcal{S}}(G) \longrightarrow \tilde{\mathbb{Z}}_{\geq 0} : \tilde{\nu}_H([S], [T]) = \nu_H(S, T)$.
- $$\tilde{\varrho} := \log \tilde{\nu}_H : \tilde{\mathcal{S}}(G) \times \tilde{\mathcal{S}}(G) \longrightarrow \log \tilde{\mathbb{Z}}_{\geq 0}$$
- (i) We have natural isometric bijections
- $$\begin{array}{ccc} \mathcal{S}'(G) & \nu_H & \approx \\ S & & S \cup \{e\} \end{array} \approx \begin{array}{ccc} \mathcal{S}''(G) & \nu_H & \approx \\ S \cup \{e\} & & [S] \end{array} \approx (\tilde{\mathcal{S}}(G), \tilde{\nu}_H).$$
- (ii) ν_H is nondegenerate on $\mathcal{S}'(G)^*$ and $\mathcal{S}''(G)^*$ and $\tilde{\nu}_H$ is nondegenerate on $\tilde{\mathcal{S}}(G)^*$.
- (5) For $S, T \in \mathcal{S}''(G)$ (i) $S \cup T \subset ST \subset (S \cup T)^2$ (ii) $\nu_H(S \cup T, ST) \leq 2$, $\nu_H(ST, S \cup T) \leq 1$

Fact 3.4. There exists an isometric bijection

$$\varphi : (F(G - A, J), \mu) \longrightarrow (F_L(G^2 - \tilde{A}, J), \mu) : f \longmapsto \tilde{f} := f\eta|.$$

- Here, (i) $\eta : G^2 \twoheadrightarrow G : \eta(x, y) = x^{-1}y, \quad J \subset \mathbb{R}_{\geq 0}$
(ii) $A \subset G, \quad \tilde{A} := \eta^{-1}(A), \quad \eta| : G^2 - \tilde{A} \twoheadrightarrow G - A$
(iii) $F_L(G^2 - \tilde{A}, J) := \{g \in F(G^2 - \tilde{A}, J) \mid g(ax, ay) = g(x, y) \quad (\forall a \in G, (x, y) \in G^2 - \tilde{A})\}$
(iv) The metric μ is defined in Notation 2.5 (1).
(v) $\tilde{f}(x, y) = f(x^{-1}y), \quad f(x) = \tilde{f}(e, x) \quad (x, y \in G)$

Proposition 3.1. There exists a natural isometric injection

$$\zeta : \underset{S}{(\mathcal{S}'(G)^*, \widehat{\varrho})} \hookrightarrow \underset{\log \nu_S}{(F(G^\times, \widetilde{\mathbb{Z}}_{\geq 0}), \widehat{\mu})} \overset{\varphi}{\approx} \underset{\log d_S}{(F_L(G^2 - \Delta(G), \widetilde{\mathbb{Z}}_{\geq 0}), \widehat{\mu})}$$

$\circ \zeta$ is anti-isometric with respect to ϱ and μ (i.e., $\varrho(S, T) = \mu(\log \nu_T, \log \nu_S)$ ($\forall S, T \in \mathcal{S}'(G)^*$)).

Proof.

- (1) Consider the map $\psi : F(G, \widetilde{\mathbb{Z}}_{\geq 0}) \longrightarrow \mathcal{S}'(G) : \psi(f) = f^{-1}(0)$. Since $\psi\zeta = \text{id}$ by Fact 3.2 (2)(iii), it follows that ζ is injective.
- (2) $\nu_H(S, T) \equiv (\nu_S)_H(T) = \lambda(\nu_T, \nu_S)$ ($\forall S, T \in \mathcal{S}(G)$).

Hence, the map $\underset{S}{(\mathcal{S}'(G), \nu_H)} \longrightarrow \underset{\nu_S}{(F(G^\times, \widetilde{\mathbb{Z}}_{\geq 1}), \lambda)}$ is an anti-isometric injection.

This follows from the following observation :

$$\text{For } n \in \mathbb{Z}_{\geq 1} \quad (\nu_S)_H(T) \leq n \iff T \subset S^{\leq n} \underset{(*)}{\iff} \nu_S \leq n \nu_T \iff \lambda(\nu_T, \nu_S) \leq n$$

(*) (\implies) For any $x \in G$ we have to show that $\nu_S(x) \leq n \nu_T(x)$. Let $m := \nu_T(x)$.

If $m = \infty$, then the assertion is trivial.

If $m < \infty$, then $x \in T^m \subset (S^{\leq n})^m \subset S^{\leq mn}$. This implies that $\nu_S(x) \leq mn = n \nu_T(x)$.

(\impliedby) For any $x \in T$ it follows that $\nu_T(x) \leq 1$ and $\nu_S(x) \leq n \nu_T(x) \leq n$, so that $x \in S^{\leq n}$.

- (3) For any $S, T \in \mathcal{S}(G)^*$ it follows that

$$\varrho(S, T) \equiv \varrho_S(T) = \log (\nu_S)_H(T) = \log \lambda(\nu_T, \nu_S) = \mu(\log \nu_T, \log \nu_S) \quad \text{and}$$

$$\widehat{\varrho}(S, T) = \widehat{\mu}(\log \nu_S, \log \nu_T).$$

□

Here we have some examples of subspaces $\mathcal{X} \equiv (\mathcal{X}, \widehat{\varrho}|_{\mathcal{X}})$ of $(\mathcal{S}'(G), \widehat{\varrho})$ with $\widehat{\varrho}|_{\mathcal{X}} < \infty$

Example 3.1.

- (1) Suppose G is finitely generated and $n := \text{rank } G \geq 1$.

$$\mathcal{S}'_{fg}(G) := \{S \in \mathcal{S}'(G) \mid S = S^{-1}, \quad G = S^\infty, \quad |S| < \infty\}$$

$$\mathcal{S}'_{fg}(G)_n := \{S \in \mathcal{S}'_{fg}(G) \mid |S| = n\}$$

- (2) Suppose G is finitely normally generated and $n := \text{n-rank } G \geq 1$.

$$\mathcal{S}'_{nfg}(G)^* := \{S \in \mathcal{S}'(G)^* \mid G = S^\infty, \quad S = C_A \text{ for some } A \subset G^\times \text{ with } |A| < \infty\}$$

$$\mathcal{S}'_{nfg}(G)_n^* := \{S \in \mathcal{S}'(G)^* \mid G = S^\infty, \quad S = C_A \text{ for some } A \subset G^\times \text{ with } |A| = n\}$$

$\circ (\nu_S)_H(T) \in \mathbb{R}_{\geq 1}$ ($\forall S, T \in \mathcal{S}'_{nfg}(G)^*$)

(Proof)

- (i) Since $S = C_A$ for some $A \subset G^\times$, it follows that S is symmetric and conjugation invariant. Hence, ν_S is a conjugation invariant norm on G (cf. Fact 3.2 (3), (4)).
- (ii) Since $G = S^\infty$ and $T = C_B$ for some $B \subset G^\times$ with $|B| < \infty$, it follows that
- $$\nu_S(T) = \nu_S((B^\pm)^G) = \nu_S(B^\pm) = \nu_S(B) \quad \text{and} \quad (\nu_S)_H(T) = \sup \nu_S(T) = \max \nu_S(B) < \infty.$$

4. THE MAPS ON THE POWER SETS INDUCED FROM GROUP HOMOMORPHISMS

Suppose $f : G \rightarrow H$ is a group homomorphism between groups. Let $K := \text{Ker } f$. We are interested in the following general problem.

Problem. Describe the relation among the metric spaces $(\mathcal{S}''(G)^*, \nu_H^G)$, $(\mathcal{S}''(H)^*, \nu_H^H)$ and $(\mathcal{S}''(K)^*, \nu_H^K)$.

4.1. Generalities.

First we list some general properties.

Fact 4.1.

[I] Let $G \supset S, A, B, A_1, \dots, A_n$.

- (1) (i) $f(AB) = f(A)f(B)$ (ii) $f(A_1 \cdots A_n) = f(A_1) \cdots f(A_n)$ (iii) $f(A^B) = f(A)^{f(B)}$
 (iv) $f(A^{-1}) = f(A)^{-1}$
- (2) (i) $f(S^n) = f(S)^n$, $f(S^{\leq n}) = f(S)^{\leq n}$ ($\forall n \in \mathbb{Z}_{\geq 0}$) (ii) $f(S^\infty) = f(S)^\infty$
- (3) Suppose f is surjective.
- (i) $f(C_A) = C_{f(A)}$
- (ii) The following implication holds for the conditions $P = f, g, s, c, fc, fg, ng, nfg$:

$$S \in \mathcal{S}_P''(G) \implies f(S) \in \mathcal{S}_P''(H)$$

[II] Suppose f is surjective. Let $H \supset S, A, B, A_1, \dots, A_n$

- (1) (i) $f^{-1}(AB) = f^{-1}(A)f^{-1}(B)$ (ii) $f^{-1}(A_1 \cdots A_n) = f^{-1}(A_1) \cdots f^{-1}(A_n)$
 $\circ f^{-1}(A)K = f^{-1}(A)$
- (2) (i) $f^{-1}(S^n) = (f^{-1}(S))^n$ ($\forall n \in \mathbb{Z}_{\geq 1}$)
- (ii) $f^{-1}(S^{\leq n}) = (f^{-1}(S))^{\leq n} \cup K = \begin{cases} K & (n = 0) \\ (f^{-1}(S) \cup K)^{\leq n} & (n \in \mathbb{Z}_{\geq 1}) \end{cases}$
- (iii) $f^{-1}(AS^{\leq n}) = f^{-1}(A)(f^{-1}(S))^{\leq n}$ ($\forall n \in \mathbb{Z}_{\geq 0}$)
- (iv) $f^{-1}(S^{-1}) = (f^{-1}(S))^{-1}$ (v) $f^{-1}(S^\infty) = (f^{-1}(S))^\infty \cup K$

Notation 4.1. The group homomorphism $f : G \rightarrow H$ induces the following maps.

$$\begin{aligned} f_* : \mathcal{S}(G) &\longrightarrow \mathcal{S}(H) : f_*(S) = f(S) & \tilde{f}_* : \tilde{\mathcal{S}}(G) &\longrightarrow \tilde{\mathcal{S}}(H) : \tilde{f}_*([S]) = [f(S)] \\ f^* : \mathcal{S}(H) &\longrightarrow \mathcal{S}(G) : f^*(S) = f^{-1}(S) \end{aligned}$$

Fact 4.1 indicates that these maps are compatible with various operations in $\mathcal{S}(G)$ and $\mathcal{S}(H)$.

Fact 4.2. Suppose $f : G \rightarrow H$ is a group homomorphism.

- (1) Let $S \subset G$.
- (i) $d_{f(S)}(f(x), f(y)) \leq d_S(x, y)$ $\nu_{f(S)}(f(x)) \leq \nu_S(x)$ ($\forall x, y \in G$)
- (ii) $(d_{f(S)})_H(f(A), f(B)) \leq (d_S)_H(A, B)$ ($\forall A, B \subset G$)
 $(\nu_{f(S)})_H(f(A)) \leq (\nu_S)_H(A)$ $\nu_H(f(S), f(T)) \leq \nu_H(S, T)$

- (iii) $f_* : (\mathcal{S}(G), (d_S)_H) \longrightarrow (\mathcal{S}(H), (d_{f(S)})_H)$ is a 1-Lipschitz map.
- (2) $f_* : (\mathcal{S}(G)^* - \mathcal{S}(K), \nu_H) \longrightarrow (\mathcal{S}(H)^*, \nu_H)$ is a 1-Lipschitz map.
 $f_* : (\mathcal{S}(G)^* - \mathcal{S}(K), \widehat{\varrho}) \longrightarrow (\mathcal{S}(H)^*, \widehat{\varrho})$ is a 1-Lipschitz map.
- (3) $\widetilde{f}_* : (\widetilde{\mathcal{S}}(G)^*, \widehat{\varrho}) \longrightarrow (\widetilde{\mathcal{S}}(H)^*, \widehat{\varrho})$ is a 1-Lipschitz map.

Suppose f is surjective.

- (4) Let $S \subset H$.
 - (i) $(d_{f^{-1}(S)})_H(f^{-1}(A), f^{-1}(B)) = (d_S)_H(A, B) \quad (\forall A, B \subset H)$
 $(\nu_{f^{-1}(S)})_H(f^{-1}(A)) = (\nu_S)_H(A) \quad (\forall A \in \mathcal{S}(H)^*)$
 - (ii) $f^* : (\mathcal{S}(H), (d_S)_H) \longrightarrow (\mathcal{S}(G), (d_{f^{-1}(S)})_H)$ is an isometric injection. $f_* f^* = \text{id}$
- (5) $f^* : (\mathcal{S}''(H)^*, \nu_H) \longrightarrow (\mathcal{S}''(G)^*, \nu_H)$ is an isometric injection.
 $f^* : (\mathcal{S}''(H)^*, \widehat{\varrho}) \longrightarrow (\mathcal{S}''(G)^*, \widehat{\varrho})$ is an isometric injection.

Proposition 4.1. Suppose $f : G \rightarrow H$ is a group epimorphism and $K := \text{Ker } f$.

- (1) $f_* : (\mathcal{S}''(G)^* - \mathcal{S}(K), \nu_H) \longrightarrow (\mathcal{S}''(H)^*, \nu_H)$ is a 1-Lipschitz map.
- (2) $f^* : (\mathcal{S}''(H)^*, \nu_H) \longrightarrow (\mathcal{S}''(G)^* - \mathcal{S}(K), \nu_H)$ is an isometric section of f_* . $(f_* f^* = \text{id})$
- (3) $\widehat{\nu}_H(T, f_* f^*(T)) \leq \nu_H(T, K) + 1 \quad (\forall T \in \mathcal{S}''(G)^* - \mathcal{S}(K))$
 - (i) $T \subset f^{-1}(f(T)) = TK \quad (\forall T \in \mathcal{S}(G))$
 - (ii) $\nu_H(T, f^{-1}(f(T))) \leq \nu_H(T, K) + 1 \quad \nu_H(f^{-1}(f(T)), T) = 1$

Notation 4.2. Suppose $f : G \rightarrow H$ is a group epimorphism and $K := \text{Ker } f$. Consider the following conditions for $T \in \mathcal{S}(G)$:

- (i) $Q_m : K \subset T^{\leq m} \quad (m \in \mathbb{Z}_{\geq 0})$
- (ii) $R : T \cap K \in \mathcal{S}_g(K)$

We have the subsets $\mathcal{S}_{Q_m}(G), \mathcal{S}_R(G) \subset \mathcal{S}(G)$.

Corollary 4.1. Suppose $f : G \rightarrow H$ is a group epimorphism and $K := \text{Ker } f$.

[I] If f is an isomorphism, then

$$f_* : (\mathcal{S}''(G)^*, \widehat{\nu}_H) \longrightarrow (\mathcal{S}''(H)^*, \widehat{\nu}_H) \text{ is an isometric bijection and } f^* = f_*^{-1}.$$

[II] In each of the following cases, f_* is a 1-Lipschitz surjection and f^* is an isometric section of f_* .
Moreover, f_* is a quasi-isometric map and f^* is a quasi-isometric inverse of f_*

$$(1) (\mathcal{S}_{Q_m}''(G)^* - \mathcal{S}(K), \widehat{\nu}_H) \xrightleftharpoons[f^*]{f_*} (\mathcal{S}''(H)^*, \widehat{\nu}_H) \quad \circ \quad \widehat{\nu}_H(T, f_* f^*(T)) \leq \nu_H(T, K) + 1 \leq m + 1$$

$$(\forall T \in \mathcal{S}_{Q_m}''(G)^* - \mathcal{S}(K))$$

(2) Suppose K is a finite group.

$$(i) (\mathcal{S}_R''(G)^* - \mathcal{S}(K), \widehat{\nu}_H) \xrightleftharpoons[f^*]{f_*} (\mathcal{S}''(H)^*, \widehat{\nu}_H)$$

(a) Since $\mathcal{S}(K)$ is a finite set, there exists $m \in \mathbb{Z}_{\geq 0}$ with $K = U^{\leq m} \quad (\forall U \in \mathcal{S}_g(K))$.

(b) $\forall T \in \mathcal{S}_R(G) \quad T \cap K \in \mathcal{S}_g(K) \quad \therefore K = (T \cap K)^{\leq m} \subset T^{\leq m}$

$$\therefore \mathcal{S}_R''(G)^* \subset \mathcal{S}_{Q_m}''(G)^*.$$

$$(ii) \text{ Suppose } G \text{ is finitely generated.} \quad (\mathcal{S}_{fg,R}''(G)^*, \widehat{\nu}_H) \xrightleftharpoons[f^*]{f_*} (\mathcal{S}_{fg}''(H)^*, \widehat{\nu}_H)$$

(a) If $S \in \mathcal{S}_{fg}''(H)^*$, then $(f^{-1}(S))^{\infty} = f^{-1}(S^{\infty}) = f^{-1}(H) = G. \quad (\text{cf. Fact 4.1 [II]}).$

◦ In the statements [I] and [II] (1), (2) we can replace $\widehat{\nu}_H$ by $\widehat{\varrho}$.

4.2. The case of $\mathcal{S}_{nfg}(G)$.

We extend the arguments in the previous subsection to the case of the map $f_* : (\mathcal{S}_{nfg}''(G)^*, \widehat{\varrho}) \longrightarrow (\mathcal{S}_{nfg}''(H)^*, \widehat{\varrho})$. Throughout this subsection we assume that $f : G \rightarrow H$ is a group epimorphism between nontrivial groups and $K \equiv \text{Ker } f$.

Notation 4.3. For $S \in \mathcal{S}(H)$ we say that $R \in \mathcal{S}(G)$ is a lift of S with respect to f if

$$f(R) = S \quad \text{and} \quad |R \cap f^{-1}(x)| = 1 \quad (\forall x \in S).$$

We use the following notations.

- (i) $\mathcal{S}(G; S, f) := \{R \in \mathcal{S}(G) \mid R \text{ is a lift of } S \text{ with respect to } f.\}$
- (ii) $R_x := R \cap f^{-1}(x) \in G \quad (R \in \mathcal{S}(G; S, f), x \in S)$

Notation 4.4.

- (1) For $S \in \mathcal{S}(H)$ and $R \in \mathcal{S}(G; S, f)$ we have the map
$$h = h_S^R : f^{-1}(S) \longrightarrow K : h(x) = (R_{f(x)})^{-1}x$$
 - (i) $x = R_{f(x)}h(x) \quad (\forall x \in f^{-1}(S))$
 - (ii) $h(xa) = h(x)a \quad (\forall x \in f^{-1}(S), a \in K)$
 - (iii) $e \in S, R_e = e \implies h|_K = \text{id}, h^{-1}(e) = R$
- (2) There exists $\theta : \mathcal{S}(H) \longrightarrow \mathcal{S}(G)$ such that $\theta(S) \in \mathcal{S}(G; S, f)$ for each $S \in \mathcal{S}(H)$.
 - (i) If $g : H \rightarrow G$ is a (non-homomorphic) section of f , then we have the map
$$\theta_g : \mathcal{S}(H) \longrightarrow \mathcal{S}(G) : \theta_g(S) := g(S) \in \mathcal{S}(G; S, f)$$
- (3) Consider the following maps :
$$\chi : \mathcal{S}(G) \longrightarrow \mathcal{S}(H \times K) : \chi(T) := \{(f(x), h_S(x)) \mid x \in T\} \subset f(T) \times K,$$

$$\omega : \mathcal{S}(H \times K) \longrightarrow \mathcal{S}(G) : \omega(W) = \{\theta(p(W))_u v \mid (u, v) \in W\},$$

where $S := f(T) \in \mathcal{S}(H)$ and $p : H \times K \longrightarrow H$ is the projection.

 - (i) $T = \{\theta(S)_u v \mid (u, v) \in \chi(T)\} \subset f^{-1}(S) \quad (T \in \mathcal{S}(G)).$
 - (ii) $\omega\chi = \text{id}, \chi\omega = \text{id} \quad (\omega\chi(T) = T, \chi\omega(W) = W)$

Fact 4.3.

- (1) Let $S \in \mathcal{S}''(H)$, $R \in \mathcal{S}''(G; S, f)$ and $L, U \in \mathcal{S}''(K)$. Then, the following holds.
 - (i) $f(R^L \cup U) = f(R^L U) = S$
 - (ii) $(R^L \cup U) \cap K = (R^L U) \cap K = U$
- (2) There exists a map $\theta : \mathcal{S}(H) \rightarrow \mathcal{S}(G)$ such that
$$(*) \quad \theta(S) \in \mathcal{S}(G; S, f) \quad (S \in \mathcal{S}(H)) \quad \text{and} \quad \theta(\mathcal{S}''(H)) \subset \mathcal{S}''(G).$$

The following holds.

- (i) $\theta(\mathcal{S}''(H)^*) \subset \mathcal{S}''(G)^* - \mathcal{S}(K)$
- (ii) For each $S \in \mathcal{S}(H)$ we have $\theta(S) \in \mathcal{S}(G; S, f)$ and the map
$$h_S := h_S^{\theta(S)} : f^{-1}(S) \rightarrow K : h_S(x) = (\theta(S)_{f(x)})^{-1}x.$$
- (iii) $x = \theta(S)_{f(x)}h_S(x) \quad (\forall x \in f^{-1}(S))$
- (iv) There exists a (non-homomorphic) section $g : H \rightarrow G$ of f with $g(x^{-1}) = g(x)^{-1} \quad (x \in H)$ and $g(e) = e$. Then, the map θ_g satisfies the above conditions $(*)$ and has the following property
$$\text{For } S \in \mathcal{S}(H) \quad (a) \quad g(S^{-1}) = g(S)^{-1} \quad (b) \quad S \in \mathcal{S}_s(H) \implies g(S) \in \mathcal{S}_s(G)$$
- (3) Consider the following maps
$$\varphi : \mathcal{S}''(G) \longrightarrow \mathcal{S}''(H) \times \mathcal{S}''(K) : \varphi(T) = (f(T), T \cap K)$$

$$\psi_L, \psi'_L : \mathcal{S}''(H)^* \times \mathcal{S}''(K)^* \longrightarrow \mathcal{S}''(G)^* - \mathcal{S}(K) : \quad (L \in \mathcal{S}''(K))$$

$$\psi_L(S, U) = \theta(S)^L \cup U, \quad \psi'_L(S, U) = \theta(S)^L U$$

- (i) $\varphi\psi_L = \varphi\psi'_L = \text{id} \quad \circ \quad f(R^L \cup U) = f(R^L U) = S \quad (R^L \cup U) \cap K = (R^L U) \cap K = U$
- (ii) $\widehat{\nu}_H(\psi_L(S, U), \psi'_L(S, U)) \leq 2 \quad \circ \quad \theta(S)^L \cup U \subset \theta(S)^L U \subset (\theta(S)^L \cup U)^2$

(4) We have the following restrictions of χ and ω .

$$\begin{array}{ccccccc} \mathcal{S}(\underline{G}) & \supset & \mathcal{S}''(\underline{G}) & \supset & \mathcal{S}''(G) & \xrightarrow{\chi} \mathcal{S}(K) & \\ \chi & \omega & \chi & \omega & \chi & \omega & \omega = \chi^{-1} \\ \mathcal{S}(H \times K) & \supset & \mathcal{S}''(H \times K) & \supset & \mathcal{S}''(H \times K) - \mathcal{S}(e \times K) & & \end{array}$$

Notation 4.5. For a group G we use the following notations and terminologies :

- (1) (i) $\text{rank}_n G := \min\{n \in \widetilde{\mathbb{Z}}_{\geq 0} \mid \exists A \subset G, |A| = n, G = \langle\langle A \rangle\rangle = (C_A)^\infty\} \in \widetilde{\mathbb{Z}}_{\geq 0}$
- (ii) $\text{diam}_{nfg} G := \min\{\nu_H(S, G) \mid S \in \mathcal{S}_{nfg}(G)\} \in \widetilde{\mathbb{Z}}_{\geq 0}$
- \circ If G is a nontrivial group, then these values are in $\widetilde{\mathbb{Z}}_{\geq 1}$.
- (2) G is finitely normally generated $\iff \mathcal{S}_{nfg}(G) \neq \emptyset \iff \text{rank}_n G < \infty$
 \iff There exists $A \in \mathcal{S}_f(G)$ with $G = \langle\langle A \rangle\rangle = (C_A)^\infty$.
- (3) G is weak-uniformly finitely normally generated
 $\iff \nu_H(S, G) < \infty$ for some/any $S \in \mathcal{S}_{nfg}(G) \iff \text{diam}_{nfg} G < \infty$
 \iff There exists $A \in \mathcal{S}_f(G)$ with $\nu_H(C_A, G) < \infty$ (i.e., $G = (C_A)^{\leq n}$ for some $n \in \mathbb{Z}_{\geq 0}$).
 \circ If $S, T \in \mathcal{S}_{nfg}''(G)^*$, then $\nu_H(T, G) < \infty$ and $\nu_H(T, G) \leq \nu_H(T, S) \cdot \nu_H(S, G)$
(cf. Example 3.1 (2)).
- (4) In [6, Section 2] the following notations are introduced.
- (i) $\Delta(G) := \sup\{\nu_H(S, G) \mid S \in \mathcal{S}_{nfg}(G)\} \in \widetilde{\mathbb{Z}}_{\geq 0}$
- (ii) G is uniformly bounded. $\iff \Delta(G) < \infty$

Theorem 4.1. Suppose $f : G \rightarrow H$ is a group epimorphism between nontrivial groups and $K \equiv \text{Ker } f$ is normally finitely generated.

- [I] The map $f_* : (\mathcal{S}_{nfg}''(G)^*, \widehat{\varrho}) \longrightarrow (\mathcal{S}_{nfg}''(H)^*, \widehat{\varrho})$ is a 1-Lipschitz surjection with a section.
- [II] If K is weak-uniformly normally finitely generated and $m_0 := \text{diam}_{nfg} K \in \mathbb{Z}_{\geq 0}$, then the following holds.

- 1) The map $f_* : \mathcal{S}_{nfg}''(G)^* \longrightarrow \mathcal{S}_{nfg}''(H)^*$ has a section $\eta : \mathcal{S}_{nfg}''(H)^* \longrightarrow \mathcal{S}_{nfg, Q_{m_0}}''(G)^*$.
- 2) For any $m \in \mathbb{Z}_{\geq m_0}$ the map $f_* : (\mathcal{S}_{nfg, Q_m}''(G)^*, \widehat{\varrho}) \longrightarrow (\mathcal{S}_{nfg}''(H)^*, \widehat{\varrho})$ is a quasi-isometry and η is a quasi-isometric inverse of f_* .
- (i) $\widehat{\varrho}(\eta \times \eta) \leq \widehat{\varrho} + \log(m_0 + 1)$ (ii) $\widehat{\varrho}(\eta f_*, \text{id}) \leq \log(m + 1)$

Proof. The assertions follow from the following observations.

- [I] (1) (i) $f_*(\mathcal{S}_{nfg}''(G)^*) \subset \mathcal{S}_{nfg}''(H)^*$ (cf. Fact 4.1 [I](3)) (ii) f_* is 1-Lipschitz (cf. Proposition 4.1 (1)).
- (2) Since K is normally finitely generated, there exists $B \in \mathcal{S}_{f,s}''(K)$ with $K = (C_B)^\infty = (B^K)^\infty$ in K .
- (i) Let $U := C_B = B^G$ in G . Then $U \in \mathcal{S}_{s,fc}''(G) \cap \mathcal{S}(K)$ and $K = U^\infty$.
- (3) We define a map $\eta : \mathcal{S}_{nfg}''(H)^* \longrightarrow \mathcal{S}_{nfg}''(G)^*$ by the following rule :
 (\sharp) For each $S \in \mathcal{S}_{nfg}''(H)^*$ (a) choose $A \in \mathcal{S}_{f,s}''(H)$ with $S = C_A = A^H$ and
(b) with the lift $\theta(A) \in \mathcal{S}_f''(G, A, f)$ we set $\eta(S) := C_{\theta(A)} \cup U$.

- (i) $f(C_{\theta(A)}) = C_{f(\theta(A))} = C_A = S$. (cf. Fact 4.1 [I])
- (4) $T := \eta(S) \in \mathcal{S}_{nfg}''(G)^*$
- (i) $T \in \mathcal{S}_g(G) : f((C_{\theta(A)})^\infty) = f(C_{\theta(A)})^\infty = S^\infty = H$. (cf. Fact 4.1 [I])
 $\therefore G = (C_{\theta(A)})^\infty K \subset (C_{\theta(A)})^\infty U^\infty \subset (C_{\theta(A)} \cup U)^\infty = T^\infty$.
- (ii) $T \in \mathcal{S}_{s,fc}(G) : T = C_{\theta(A)} \cup U = C_{\theta(A)} \cup C_B = C_{\theta(A) \cup B}$ in G $\theta(A) \cup B \in \mathcal{S}_f(G)$
- (5) $f_*\eta = \text{id} : f(\eta(S)) = f(C_{\theta(A)} \cup U) = f(C_{\theta(A)}) \cup f(U) = S \cup \{e\} = S$.
- [II] (1) By the assumption there exists $\exists B \in \mathcal{S}_{f,s}''(K)$ with $K = (C_B)^{\leq m_0} = (B^K)^{\leq m_0}$ in K .
- (i) Let $U := C_B = B^G$ in G . Then, $U \in \mathcal{S}_{s,fc}''(G)^* \cap \mathcal{S}(K)$ and $K = U^{\leq m_0}$.
- (2) $T, T' \in \mathcal{S}_{Q_n}''(G)^* - \mathcal{S}(K)$ ($n \in \mathbb{Z}_{\geq 0}$), $S = f(T), S' = f(T') \implies \nu_H(T, T') \leq (n+1)\nu_H(S, S')$.
- (i) $K \subset T^{\leq n} \quad T \subset f^{-1}(S) = TK \subset T^{\leq n+1}$
 $\therefore \nu_H(T, f^{-1}(S)) \leq n+1, \nu_H(f^{-1}(S), T) = 1 \quad \therefore \widehat{\nu}_H(T, f^{-1}(S)) \leq n+1$
 \circ Similar for T' and S' .
- (ii) $\nu_H(f^{-1}(S), f^{-1}(S')) = \nu_H(S, S')$ (cf. Proposition 4.1 (2))
- (iii) $\nu_H(T, T') \leq \nu_H(T, f^{-1}(S)) \cdot \nu_H(f^{-1}(S), f^{-1}(S')) \cdot \nu_H(f^{-1}(S'), T') \leq (n+1)\nu_H(S, S')$.
- (3) We define the map $\eta : \mathcal{S}_{nfg}''(H)^* \longrightarrow \mathcal{S}_{nfg}''(G)^*$ as in [I].
- (i) $\eta(S) \in \mathcal{S}_{nfg, Q_{m_0}}''(G)^*$, since $K = U^{\leq m_0} \subset \eta(S)^{\leq m_0}$. (ii) $f(\eta(S)) = S$.
- (iii) $\nu_H(\eta(S), \eta(S')) \leq (m_0+1)\nu_H(S, S')$ ($S, S' \in \mathcal{S}_{nfg}''(H)^*$) (by (2))
- (iv) $\widehat{\nu}_H(\eta \times \eta) \leq (m_0+1)\widehat{\nu}_H \quad \therefore \widehat{\varrho}(\eta \times \eta) \leq \widehat{\varrho} + \log(m_0+1)$
- (4) $f_* : (\mathcal{S}_{nfg, Q_m}''(G)^*, \widehat{\nu}_H) \longrightarrow (\mathcal{S}_{nfg}''(H)^*, \widehat{\nu}_H)$ is a 1-Lipschitz surjection with a section η .
- (i) $\eta : (\mathcal{S}_{nfg}''(H)^*, \widehat{\nu}_H) \longrightarrow (\mathcal{S}_{nfg, Q_m}''(G)^*, \widehat{\nu}_H)$ is a quasi-isometric map. (ii) $f_*\eta = \text{id}$
- (iii) For $T \in \mathcal{S}_{nfg, Q_m}''(G)^*$ and $S := f(T)$
 $\widehat{\nu}_H(T, \eta(S)) \leq m+1$ by (2) $\therefore \widehat{\varrho}_H(T, \eta(S)) \leq \log(m+1)$. □

4.3. Semidirect products.

This subsection is devoted to the case of semidirect products (or the case of group epimorphisms with homomorphic sections). Suppose a group G is the semidirect product of a subgroup H and a normal subgroup K ($G = H \ltimes K$). Let $p : G \rightarrow H$ and $q : G \rightarrow K$ denote the projections onto the factors H and K . The map p is a group epimorphism with $p|_H = \text{id}_H$ and $\text{Ker } p = K$. Any $S \in \mathcal{S}(H)$ is its canonical lift with respect to p , to which we can apply the arguments in the previous subsection. We set

$$\mathcal{S}_p''(K)^G := \mathcal{S}_p''(K) \cap \mathcal{S}_c(G) \subset \mathcal{S}_p''(K) \quad (\text{cf. Notation 3.3}).$$

Fact 4.4.

- (1) Let $S \in \mathcal{S}''(H)$, $U \in \mathcal{S}''(K)$ and $L \in \mathcal{S}''(K)$.
- (i) $S \subset S^L \subset S^L \cup U \subset S^L U$
- (ii) $p(S^L) = p(S^L \cup U) = p(S^L U) = S \quad (S^L \cup U) \cap K = (S^L U) \cap K = U$
 $S^L \cap H = S \quad S^L U \cap H = S$
- (iii) $S^K U = U S^K$ ($\because S^K : K$ -conjugate-invariant $U \subset K$)
 $\circ p(S^L) = p(S)^{p(L)} = S^e = S, \quad p(S^L U) = p(S^L)p(U) = S^e = S$
 $\circ S^K \cap K = \{e\},$
 $\circ S^K = ((S - \{e\}) \cup \{e\})^K = (S - \{e\})^K \cup \{e\}^K = (S - \{e\})^K \cup \{e\} \subset (S - \{e\})K \cup \{e\}$

- $S^K U \subset ((S - \{e\})K \cup \{e\})U \subset (S - \{e\})K \cup U$
- (iii) $AK \cap BK = (A \cap B)K \quad (A, B \subset H), \quad (H^\times K) \cap K = \emptyset, \quad A^K \subset A^K K \subset AK \quad (A \subset G)$
- (2) Let $S \in \mathcal{S}_{s,c}''(H)$ and $U \in \mathcal{S}_{s,c}''(K)^G := \mathcal{S}_{s,c}''(K) \cap \mathcal{S}_c(G)$
 - (i) $S^K \in \mathcal{S}_{s,c}''(G)$ (ii) $S^K \cup U, S^K U \in \mathcal{S}_{s,c}''(G)$ (cf. Fact 3.1 (6)(vi))
 - $(S^K)^G = S^{KG} = S^G = S^{HK} = (S^H)^K = S^K \quad (S^K)^{-1} = (S^{-1})^K = S^K$
- (3) Let $S \in \mathcal{S}_g''(H)$ and $U \in \mathcal{S}_g''(K)$.
 - (i) $S \cup U, SU \in \mathcal{S}_g''(G)$
 - $(S \cup U)^\infty \supset S^\infty \cup U^\infty = H \cup K, \quad (S \cup U)^\infty = ((S \cup U)^\infty)^2 \supset (H \cup K)^2 \supset HK = G$
 - (ii) $S \in \mathcal{S}_{fg}''(H), \quad U \in \mathcal{S}_{fg}''(K) \implies S \cup U \in \mathcal{S}_{fg}''(G), \quad SU \in \mathcal{S}_{fg}''(G)$
 - (iii) $S \in \mathcal{S}_{nfg}''(H), \quad U \in \mathcal{S}_g''(K), \quad U = B^G \quad (\exists B \in \mathcal{S}_{f,s}(K)) \implies S^K \cup U \in \mathcal{S}_{nfg}''(G)$
 - $S = A^H \quad (\exists A \in \mathcal{S}_{f,s}(H)) \quad U = B^G \quad (\exists B \in \mathcal{S}_{f,s}(K)) \quad e \in S, e \in U \quad \therefore e \in A, e \in B$
 - $S^K = (A^H)^K = A^{HK} = A^G \quad S^K \cup U = A^G \cup B^G = (A \cup B)^G = C_{A \cup B} \quad A \cup B \in \mathcal{S}_{f,s}(G)$
 - $S^K U \supset S^K \cup U \supset S \cup U \in \mathcal{S}_g''(G)$
 - $S^K U = A^G B^G = B^G A^G \supset (AB \cup BA)^G \supset A^G \cup B^G = S^K \cup U$

Fact 4.5. Let $T \in \mathcal{S}''(G)$. We have $p(T) \in \mathcal{S}''(H)$ and $T \cap K \in \mathcal{S}''(K)$.

- (1) The following implication holds for the conditions $P = f, g, s, c, fc, fg, ng, nfg$:

$$T \in \mathcal{S}_P''(G) \implies p(T) \in \mathcal{S}_P''(H) \quad (\text{cf. Fact 4.1 [I]})$$
- (2) The following implication holds for the conditions $P = f, s, c$:

$$T \in \mathcal{S}_P''(G) \implies T \cap K \in \mathcal{S}_P''(K) \quad (\text{cf. Fact 3.1 (5), (6)})$$

Notation 4.6. Consider the following maps.

- (1) $\varphi : \mathcal{S}''(G) \longrightarrow \mathcal{S}''(H) \times \mathcal{S}''(K) : \varphi(T) = (p(T), T \cap K)$
- (2) $\psi = \psi_L : \mathcal{S}''(H)^* \times \mathcal{S}''(K)^* \longrightarrow \mathcal{S}''(G)^* : \psi(S, U) = S^L \cup U \quad (L \in \mathcal{S}''(K))$
 $\psi' = \psi'_L : \mathcal{S}''(H)^* \times \mathcal{S}''(K)^* \longrightarrow \mathcal{S}''(G)^* : \psi'(S, U) = S^L U$
 - (i) $\varphi\psi = \varphi\psi' = \text{id} \quad (\text{cf. Fact 4.4 (1)})$
 - (ii) $\widehat{\nu}_H(\psi(S, U), \psi'(S, U)) \leq 2$
 - $S^L \cup U \subset S^L U \subset (S^L \cup U)^2 \quad \therefore \nu_H(S^L \cup U, S^L U) \leq 2, \quad \nu_H(S^L U, S^L \cup U) \leq 1$
- (iii) When G is the product of H and K , we have $S^L = S$.
- (3) We use the following notations :
 - (i) $\nu_H^G, \nu_H^H, \nu_H^K$: the multiplicative as metric ν_H on $\mathcal{S}''(G)^*, \mathcal{S}''(H)^*, \mathcal{S}''(K)^*$ respectively.
 - (ii) $\nu_H^{H,K} \in \mathcal{M}_{m,as}(\mathcal{S}''(H)^* \times \mathcal{S}''(K)^*)$: the product metric of ν_H^H and ν_H^K

$$\nu_H^{H,K}((S, U), (S', U')) = \max\{\nu_H^H(S, S'), \nu_H^K(U, U')\}$$
 - (iii) $\mu, \mu' \in \mathcal{M}_{m,as}(\mathcal{S}''(H)^* \times \mathcal{S}''(K)^*)$: the metrics induced from ν_H^G by the injections ψ and ψ'

$$\mu((S, U), (S', U')) = \nu_H^G(\psi(S, U), \psi(S', U')) \quad ((S, U), (S', U') \in \mathcal{S}''(H)^* \times \mathcal{S}''(K)^*)$$

$$\mu'((S, U), (S', U')) = \nu_H^G(\psi'(S, U), \psi'(S', U'))$$

We are interested in the description of $\text{Im } \psi$ and a deviation of μ and μ' from $\nu_H^{H,K}$.

Proposition 4.2. Let $(S, U), (S', U') \in \mathcal{S}''(H)^* \times \mathcal{S}''(K)^*$ and $L \in \mathcal{S}''(K)$.

[I] the case of ψ : Let $T = S^L \cup U, T' = (S')^L \cup U'$.

- (1) For $X \in \mathcal{S}''(G) \quad X \in \text{Im } \psi \iff X = (X \cap H)^L \cup (X \cap K), \quad X \cap H \in \mathcal{S}''(H)^*, \quad X \cap K \in \mathcal{S}''(K)^*$
 - $T \cap H = p(T) = S, \quad T \cap K = U$

- (2) (i) $\psi : \mathcal{S}_P''(H)^* \times \mathcal{S}_P''(K)^* \longrightarrow \mathcal{S}_P''(G)^*$ in the following cases :
- (a) $P = g$ (b) $P = s$ (c) $P = f$ and $L \in \mathcal{S}_f''(K)$ (d) $P = fg$ and $L \in \mathcal{S}_f''(K)$

When $L = K$:

- (ii) $\psi : \mathcal{S}_P''(H)^* \times (\mathcal{S}_P''(K)^* \cap \mathcal{S}_c(G)) \longrightarrow \mathcal{S}_P''(G)^*$ for (a) $P = c$, (b) $P = ng$
 ◦ If $S \in \mathcal{S}_c''(H)$, then $(S^K)^G = S^{KG} = S^G = S^{HK} = (S^H)^K = S^K$.
- (iii) $\psi : \mathcal{S}_{fc}''(H) \times (\mathcal{S}''(K) \cap \mathcal{S}_{fc}''(G)) \longrightarrow \mathcal{S}_{fc}''(G)$
 ◦ If $S = A^H$ ($A \in \mathcal{S}_f(H)$), $U = B^G$ ($B \in \mathcal{S}_f(K)$), then
 $S^K = (A^H)^K = A^{HK} = A^G$ $S^K \cup U = A^G \cup B^G = (A \cup B)^G$
- (iii) $\psi : \mathcal{S}_{nfg}''(H) \times (\mathcal{S}_{s,g}''(K) \cap \mathcal{S}_{fc}''(G)) \longrightarrow \mathcal{S}_{nfg}''(G)$
 ◦ $\text{Im } \psi \subset \{T \in \mathcal{S}_{nfg}''(G) \mid T \cap K \in \mathcal{S}_{s,g}''(K) \cap \mathcal{S}_{fc}''(G)\}$
- (3) $\mu((S, U), (S', U')) = \nu_H^G(T, T') = \max\{\nu_H^H(S, S'), \nu_H^G(T, U')\}$
 (i) For $n \in \mathbb{Z}_{\geq 0}$ $\nu_H^G(T, T') \leq n \iff T' \subset T^{\leq n} \iff [S' \subset S^{\leq n}, U' \subset T^{\leq n}]$
 $\iff [\nu_H^H(S, S') \leq n, \nu_H^G(T, U') \leq n]$
 ◦ $T' \subset T^{\leq n} \iff [S' \subset S^{\leq n}, U' \subset T^{\leq n}]$
 $(\implies) S' = p(T') \subset p(T^{\leq n}) = p(T)^{\leq n} = S^{\leq n}$
 $(\impliedby) (S')^L \subset (S^{\leq n})^L \subset (S^L)^{\leq n} \subset T^{\leq n}$

[II] the case of $\psi' : \text{Let } T = S^L U, T' = (S')^L U'.$

- (1) For $X \in \mathcal{S}''(G)$ $X \in \text{Im } \psi' \iff X = (X \cap H)^L (X \cap K)$, $X \cap H \in \mathcal{S}''(H)^*$, $X \cap K \in \mathcal{S}''(K)^*$
 ◦ $T \cap H = p(T) = S$, $T \cap K = U$
- (2) (i) $\psi' : \mathcal{S}_P''(H)^* \times \mathcal{S}_P''(K)^* \longrightarrow \mathcal{S}_P''(G)^*$ in the following cases :
 (a) $P = g$ (b) $P = s$ (c) $P = f$ and $L \in \mathcal{S}_f''(K)$ (d) $P = fg$ and $L \in \mathcal{S}_f''(K)$
 (\therefore) (b) If $S^{-1} = S$ and $U^{-1} = U$, then
 $(S^K U)^{-1} = U^{-1} (S^K)^{-1} = U (S^{-1})^K = U S^K = S^K U$ (cf. Fact 3.1 (1)(vi))

When $L = K$:

- (ii) $\psi' : \mathcal{S}_P''(H)^* \times (\mathcal{S}_P''(K)^* \cap \mathcal{S}_c(G)) \longrightarrow \mathcal{S}_P''(G)^*$ for (a) $P = c$, (b) $P = ng$
 (\therefore) (a) If $S^H = S$ and $U^G = U$, then $(S^K)^G = S^K$ and $(S^K U)^G \subset (S^K)^G U^G = S^K U$.
- (3) $\mu'((S, U), (S', U')) = \nu_H^G(T, T') :$
 $\max\{\nu_H^H(S, S'), \nu_H^G(T, U')\} \leq \nu_H^G(T, T') \leq 2 \max\{\nu_H^H(S, S'), \nu_H^G(T, U')\}$
 (i) For $n \in \mathbb{Z}_{\geq 0}$ $T' \subset T^{\leq n} \xRightarrow{(\#)} [S' \subset S^{\leq n}, U' \subset T^{\leq n}] \xRightarrow{(b)} T' \subset T^{\leq 2n}$
 ($\#$) $S' = p(T') \subset p(T^{\leq n}) = p(T)^{\leq n} = S^{\leq n}$ (b) $(S')^L \subset (S^{\leq n})^L \subset (S^L)^{\leq n} \subset T^{\leq n}$

As expected, we can see that $\mu = \mu' = \nu_H^{H,K}$, when G is the direct product of the subgroups H and K .

Fact 4.6. In the case $G = H \times K$, the following holds.

- (1) $[H, K] = \{e\}$
 (2) The map $q : G \rightarrow K$ is also a group epimorphism with $q|_K = \text{id}$ and $\text{Ker } q = H$.
 (3) Let $S \in \mathcal{S}''(H)$ and $U \in \mathcal{S}''(K)$.
 (i) $q(S \cup U) = q(SU) = U$
 (ii) $S^L = S$ ($\forall L \in \mathcal{S}''(K)$) $\therefore \psi, \psi'$ are independent of L .
 (iii) $SU = US$ $(SU)^n = S^n U^n$, $(S \cup U)^n \supset S^n \cup U^n$ ($n \in \mathbb{Z}_{\geq 0}$)

- (iv) $S^K = S \quad S^G = S^{KH} = (S^K)^H = S^H$
- (v) $U^H = U \quad U^G = U^{HK} = (U^H)^K = U^K$
- (vi) $(SU)^G = S^H U^K \quad \circ \quad (xy)^{ab} = x^{ab} y^{ab} = (x^b)^a (y^a)^b = x^a y^b \quad ((x, a, y, b) \in S \times H \times U \times K)$
- (4) (i) $\mathcal{S}_c''(H) = \mathcal{S}''(H) \cap \mathcal{S}_c(G)$
- (ii) $\mathcal{S}_c''(K) = \mathcal{S}''(K) \cap \mathcal{S}_c(G) \quad \mathcal{S}_P''(K) \cap \mathcal{S}_c(G) = \mathcal{S}_{P,c}''(K)$
- (iii) $\mathcal{S}''(K) \cap \mathcal{S}_{fc}''(G) = \mathcal{S}_{fc}''(K)$
- (iv) $\mathcal{S}_{s,g}''(K) \cap \mathcal{S}_{fc}(G) = \mathcal{S}_{nfg}''(K)$

Proposition 4.3. Suppose $G = H \times K$. For $(S, U), (S', U') \in \mathcal{S}''(H)^* \times \mathcal{S}''(K)^*$ we set

$T = S \cup U, T' = S' \cup U'$ in the case of ψ and $T = SU, T' = S'U'$ in the case of ψ' .

- (1) $\mu = \mu' = \nu_H^{H,K}$ (i.e., $\mu((S, U), (S', U')) = \nu_H^G(T, T') = \max\{\nu_H^H(S, S'), \nu_H^K(U, U')\}$)
- (i) For $n \in \mathbb{Z}_{\geq 0} \quad T' \subset T^n \iff S' \subset S^n, U' \subset U^n$
- ($\cdot \cdot$) (\implies) $S' = q(T') \subset q(T^n) = q(T)^n = S^n, \quad U' = q(T') \subset q(T^n) = q(T)^n = U^n.$
- (\iff) $(S \cup U)^n \supset S^n \cup U^n \supset S' \cup U', \quad (SU)^n = S^n U^n \supset S'U'$
- \circ The following maps are isometric injections :

$$\psi, \psi' : (\mathcal{S}''(H)^* \times \mathcal{S}''(K)^*, \hat{\varrho}^{H,K}) \longrightarrow (\mathcal{S}''(G)^*, \hat{\varrho}^G) : \quad \psi(S, U) = S \cup U, \quad \psi'(S, U) = SU$$

$$\text{Here, } \hat{\varrho}^{H,K} = \log \hat{\nu}_H^{H,K} = \max\{\hat{\varrho}^H, \hat{\varrho}^K\}$$

- (2) $\psi, \psi' : \mathcal{S}_P''(H)^* \times \mathcal{S}_P''(K)^* \longrightarrow \mathcal{S}_P''(G)^* \quad \text{for } P = g, s, f, c, fc, fg, ng, nfg.$

- (i) The assertions follow from Fact 4.6 (4) and Proposition 4.2 [I](2), [II](2) except the cases for ψ' and $P = fc, nfg (= g + s + fc)$.
- (ii) the case of ψ' and $P = fc$: If $S = A^H \quad (A \in \mathcal{S}_f(H)), \quad U = B^K \quad (B \in \mathcal{S}_f(K))$, then $SU = A^H B^K = (AB)^G$ (cf. Fact 4.6 (3)(iv)).

5. WORD METRICS IN GROUP ACTIONS

5.1. Word metrics in group actions.

Suppose X is a set, G is a group, e is the identity element of G and $\eta : X \times G \rightarrow X$ is a right group action.

Notation 5.1. We use the following notations. Let $x \in X, A \subset X$ and $a \in G, S \subset G$.

- (1) We have the maps $\eta_x : G \rightarrow xG \subset X : \eta_x(a) = xa$ and $\eta^a : X \approx X : \eta^a(x) = xa.$
- \circ The map η_x is injective if and only if the action of G on the orbit xG is free.
- (2) $AS := \{xs \mid x \in A, s \in S\} \subset X$

Fact 5.1. Let $S, T, U, S_\lambda \subset G \quad (\lambda \in \Lambda)$ and $x, y \in X, A, B \subset X$.

- (1) (i) $y \in xS \iff x \in yS^{-1}$ (ii) $y \in xS^n \iff x \in y(S^{-1})^n = y(S^n)^{-1} \quad (n \in \tilde{\mathbb{Z}}_{\geq 0})$
- (2) (i) $A \subset B, S \subset T \implies AS \subset BT$ (ii) $(AS)T = A(ST)$ (iii) $A(\bigcup_{\lambda \in \Lambda} S_\lambda) = \bigcup_{\lambda \in \Lambda} AS_\lambda$
- (3) (i) $AS^0 = A$ (ii) $AS^\infty = \bigcup_{n=0}^\infty AS^n$

Definition 5.1. Let $S \subset G$.

- (1) the word metric on X with respect to S :

$$d_S : X \times X \longrightarrow \tilde{\mathbb{Z}}_{\geq 0} : \quad d_S(x, y) = \begin{cases} \min\{n \in \mathbb{Z}_{\geq 0} \mid y \in xS^n\} & (y \in xS^\infty) \\ \infty & (y \in X - xS^\infty) \end{cases}$$

(2) the Hausdorff metric on $\mathcal{S}(X)$ induced from d_S :

$$(d_S)_H : \mathcal{S}(X) \times \mathcal{S}(X) \longrightarrow \widetilde{\mathbb{Z}}_{\geq 0} :$$

$$(d_S)_H(A, A') := \begin{cases} \min\{n \in \mathbb{Z}_{\geq 0} \mid A' \subset AS^{\leq n}\} & (\exists n \in \mathbb{Z}_{\geq 0} \text{ s.t. } A' \subset AS^{\leq n}) \\ \infty & (\text{otherwise}) \end{cases}$$

$$(i) \text{ For } n \in \mathbb{Z}_{\geq 0} \quad (d_S)_H(A, A') \leq n \iff A' \subset AS^{\leq n}$$

Fact 5.2. Let $x, y \in X$, $a \in G$, $S, T \subset G$ and $n \in \mathbb{Z}_{\geq 0}$.

- (1) $d_S \in \mathcal{M}_{as}(X)$, d_S is nondegenerate $\therefore \widehat{d_S} \in \mathcal{M}(X)$
 - (i) $d_S(x, y) \leq n \iff y \in xS^{\leq n}$
 - (ii) $d_S(x, y) = 0 \iff x = y$
 - (iii) $B_{d_S}(x, n) := \{y \in X \mid d_S(x, y) \leq n\} = xS^{\leq n}$
- (2) (i) $d_S(x, y) = d_{S^{-1}}(y, x)$ (ii) $S = S^{-1} \implies d_S \in \mathcal{M}(X)$
- (3) (i) $d_{S^a}(xa, ya) = d_S(x, y)$ (ii) $d_S(xa, ya) = d_S(x, y)$ if S is conjugation-invariant.
- (4) (i) $d_S = d_{S \cup \{e\}}$ (ii) $d_S(x, \cdot)^{-1}(1) = xS - x$
- (5) $S \subset T \implies d_S^X \geq d_T^X$.

Example 5.1.

- (1) In the case $(X, \eta) = (G, R)$, where R denotes the action of G on itself by the right translation, for any $S \subset G$ the metric d_S is just the word metric defined in Definition 3.2.
- (2) Suppose $X = (X, *)$ is a set with a binary operation. Consider the group $G = \text{Aut}(X)$ with the multiplication $f \cdot g = g \circ f$ ($f, g \in G$).
 - (i) The group G has the canonical right action on X : $x \cdot f = f(x)$ ($x \in X, f \in G$).
 - (ii) Any subset $S \subset G$ induces the word metric d_S on X .

When $X = (X, *)$ is a quandle, it has the symmetries :

$$\sigma_a \in \text{Aut}(X) \quad (a \in X) : \quad \sigma_a(x) = x * a \quad (x \in X)$$

The group G includes the following subgroups :

$$G > \text{Inn}(X) := \langle \sigma_a \mid a \in X \rangle > \text{Dis}(X) := \langle \sigma_a \sigma_b^{-1} \mid a, b \in X \rangle.$$

For example, we can take S as the generating sets of these subgroups.

Fact 5.3. Each $S \in \mathcal{S}(G)$ induces the word metrics $d_S \equiv d_S^G \in \mathcal{M}_{as}(G)$ and $d_S \equiv d_S^X \in \mathcal{M}_{as}(X)$.

- (1) For $x \in X$ we have the map $\eta_x : (G, d_S) \twoheadrightarrow (xG, d_S|_x)$.
 - (i) η_x is a 1-Lipschitz map.
 - (ii) If η_x is injective, then η_x is an isometric bijection.
- (2) $d_S^X \leq \nu_H(S, T) d_T^X$ ($S, T \in \mathcal{S}(G)$) (Here, we set $\infty \cdot 0 = 0$.)

Recall the bijections $\mathcal{S}'(G) \approx \mathcal{S}''(G) \approx \widetilde{\mathcal{S}}(G)$ in Notation 3.3 [II](4).

Proposition 5.1. Consider the map $\varphi : (\mathcal{S}''(G)^*, \widehat{\varrho}) \approx (\widetilde{\mathcal{S}}(G)^*, \widetilde{\varrho} = \widehat{\varrho}) \longrightarrow (F(X^2 - \Delta(X), \widetilde{\mathbb{R}}_{\geq 0}), \widehat{\mu})$.

$$\begin{matrix} S & [S] & \log d_S^X| \end{matrix}$$

Here, $d_S^X| := d_S^X|_{X^2 - \Delta(X)}$. Suppose X is not a singleton.

- (1) φ is 1-Lipschitz.
- (2) If $\eta_x : G \rightarrow X$ is injective for some $x \in X$, then
 - (i) $\lambda(d_T^X|, d_S^X|) = \nu_H(S, T)$ ($S, T \in \mathcal{S}(G)^*$).
 - (ii) φ is an isometric injection. (φ is an anti-isometric injection with respect to ϱ and μ).

Proof. Let $r := \lambda(d_T^X, d_S^X) \in \widetilde{\mathbb{R}}_{\geq 1}$.

(1) $r \leq \nu_H(S, T)$ by Fact 5.3 (2).

(2) (i) $d_S^X \leq r d_T^X \quad \therefore d_S^G \leq r d_T^G$ by Fact 5.3 (1)(ii). $\therefore \nu_S \leq r \nu_T$
 $\therefore \nu_H(S, T) = \sup \nu_S(T) \leq r \sup \nu_T(T) = r$ □

5.2. Symmetry of (X, G, ϱ) .

Notation 5.2.

- (1) Suppose (X, G, ϱ) and (Y, H, ζ) are right group actions on sets X and Y . $X \times G \xrightarrow{f \times \varphi} Y \times H$
 An equivariant map $(f, \varphi) : (X, G, \varrho) \longrightarrow (Y, H, \zeta)$ is $\begin{array}{ccc} \varrho & & \zeta \\ \downarrow & f & \downarrow \\ X & \xrightarrow{f} & Y \end{array}$
 a pair of a map $f : X \rightarrow Y$ and a group homomorphism $\varphi : G \rightarrow H$
 such that $f(xa) = f(x)\varphi(a)$ ($\forall (x, a) \in X \times G$).

The composition of equivariant maps and the identity maps are defined as follows.

- (i) $(g, \psi)(f, \varphi) := (gf, \psi\varphi)$ for $(X, G, \varrho) \xrightarrow{(f, \varphi)} (Y, H, \zeta) \xrightarrow{(g, \psi)} (Z, I, \eta)$.
 (ii) $\text{id}_{(X, G, \varrho)} := (\text{id}_X, \text{id}_G)$
 (2) In the category of right group actions on sets and equivariant maps between them.
 (i) $\text{End}(X, G, \varrho) \supset \text{Aut}(X, G, \varrho)$ denote the monoid of endomorphisms and the group of automorphisms of (X, G, ϱ) .
 (ii) An equivariant map $(f, \varphi) : (X, G, \varrho) \longrightarrow (Y, H, \zeta)$ is an isomorphism if and only if both f and φ are bijective. In this case $(f, \varphi)^{-1} = (f^{-1}, \varphi^{-1})$.

Suppose (X, G, ϱ) is a right group action on a set X , M is a set with a binary operation and $(f, \varphi) : M \ni u \longmapsto (f_u, \varphi_u) \in \text{End}(X, G, \varrho)$ is an anti-homomorphism.

Notation 5.3. Let $u, v \in M$, $x \in X$, $a, b \in G$ and $S \subset G$.

- (1) $\varphi : M \longrightarrow \text{End}(G)$ is an anti-homomorphism. $\circ (f_{uv}, \varphi_{uv}) = (f_v, \varphi_v)(f_u, \varphi_u) = (f_v f_u, \varphi_v \varphi_u)$
 (2) $x^u := f_u(x)$, $a^u := \varphi_u(a)$
 (i) $x^{uv} = (x^u)^v$ (ii) $(xa)^u = x^u a^u$
 (iii) $a^{uv} = (a^u)^v$ (iv) $(ab)^u = a^u b^u$ (v) $(a^b)^u = (a^u)^{b^u}$
 (3) (i) $S^u := \varphi_u(S) = \{a^u \mid a \in S\}$ (ii) $S \subset G$ is φ -invariant. $\iff S^u \subset S$ ($\forall u \in M$)

Fact 5.4. Let $A \subset X$, $S, T \subset G$, $u, v \in M$ and $n \in \mathbb{Z}_{\geq 0}$.

- (1) (i) $(A^u)^v = A^{uv}$ (ii) $(AS)^u = A^u S^u$
 (2) (i) $(S^u)^v = S^{uv}$ (ii) $(ST)^u = S^u T^u$ (iii) $(S^T)^u = (S^u)^{T^u}$
 (3) (i) $(S^n)^u = (S^u)^n$ (ii) $(S^{\leq n})^u = (S^u)^{\leq n}$ (iii) $(S^\infty)^u = (S^u)^\infty$

Fact 5.5. Let $S \in \mathcal{S}(G)$.

- (1) $d_{S^u}^X(x^u, y^u) \leq d_S^X(x, y)$ ($\forall x, y \in X$, $u \in M$)
 (2) If S is φ -invariant, then $d_S^X(x^u, y^u) \leq d_S^X(x, y)$ ($\forall x, y \in X$, $u \in M$).
 (3) Suppose M is a monoid, $e \in M$ is the unit element and (f, φ) is an anti-homomorphism of monoids (i.e., $(f_e, \varphi_e) = \text{id}$ ($x^e = x, a^e = a$)). Let $U(M)$ denote the group of units of M . For $u \in U(M)$ the following holds.
 (i) $d_{S^u}^X(x^u, y^u) = d_S^X(x, y)$ ($\forall x, y \in X$)

- (ii) If S is φ -invariant, then $S^u = S$ and $d_S^X(x^u, y^u) = d_S^X(x, y)$ ($\forall x, y \in X$).

Example 5.2.

- (1) Suppose M is a $*$ -subset of $\text{End}(X, G, \varrho)$ (i.e., M is closed under the multiplication). We reverse the multiplication of M so that the inclusion $M \subset \text{End}(X, G, \varrho)$ is an anti-homomorphism.
- (i) M acts on X and G by $x^{(f, \varphi)} = f(x)$ and $a^{(f, \varphi)} = \varphi(a)$ ($x \in X, a \in G, (f, \varphi) \in M$).
- (ii) If $S \subset G$ is M -invariant (i.e., $S^{(f, \varphi)} = \varphi(S) \subset S$ ($\forall (f, \varphi) \in M$)), then
- $$d_S^X(f(x), f(y)) \leq d_S^X(x, y) \quad (\forall x, y \in X, (f, \varphi) \in M).$$
- (2) When $M \subset \text{Aut}(X, G, \varrho)$ is a subgroup under the reverse multiplication,
- (i) $d_{\varphi(S)}^X(f(x), f(y)) = d_S^X(x, y)$ ($\forall x, y \in X, (f, \varphi) \in M$)
- (ii) If $S \subset G$ is M -invariant, then $\varphi(S) = S$ and
- $$d_S^X(f(x), f(y)) = d_S^X(x, y) \quad (\forall x, y \in X, (f, \varphi) \in M).$$

Example 5.3. Each action (X, G, ϱ) admits a canonical anti-homomorphism

$$(f, \varphi) : G \longrightarrow \text{Aut}(X, G, \varrho) : (f_a, \varphi_a) = (\varrho_a, a^{-1}(\cdot)a).$$

However, this symmetry yields no new information, as seen below.

- (1) Let $x \in X, a, g \in G$ and $S \subset G$.
- (i) $x^a = \varrho_a(x) = xa, g^a = \varphi_a(g) = a^{-1}ga = g^a$ (in the sense of Section 3)
- (ii) $S^a = \varphi_a(S) = a^{-1}Sa = S^a$ (in the sense of Section 3)
- (iii) S is φ -invariant. $\iff a^{-1}Sa \subset S$ ($\forall a \in G$) $\iff S$ is conjugation invariant.
- (2) Fact 5.5 (3) implies the following statements, which exactly restate Fact 5.2 (3).
- (i) $d_{S^a}^X(xa, ya) = d_S^X(x, y)$ ($\forall x, y \in X, \forall a \in G$).
- (ii) If S is conjugation invariant, then $d_S^X(xa, ya) = d_S^X(x, y)$ ($\forall x, y \in X, \forall a \in G$).
- (3) If we consider the action of a group G on itself by the right translation

$$(G, G, R) : R : G \curvearrowright G : R(x, a) = xa$$

and the anti group homomorphism

$$(f, \varphi) : G \longrightarrow \text{Aut}(G, G, R) : (f_a, \varphi_a) = (R_a, a^{-1}(\cdot)a),$$

then we recover the statements in Fact 3.2.

6. WORD LENGTH ON $*$ -SETS

6.1. $*$ -sets.

In this section we extend the arguments for groups in the previous sections to the case of sets with binary operations.

Definition 6.1. A $*$ -set means a set X with binary operations $*_i : X \times X \rightarrow X$ ($i = 1, \dots, m$).

Let $X = (X, (*_i)_{i \in [m]})$ be a nonempty $*$ -set.

Notation 6.1.

- (1) A $*$ -subset of X means a subset $Y \subset X$ such that $y_1 *_i y_2 \in Y$ ($\forall i \in [m]$) for any $y_1, y_2 \in Y$.
- (i) In the case of unital $*$ -sets (i.e., $*$ -sets with an identity element), we can consider unital $*$ -subsets.
- (2) $\mathcal{S}_*(X) :=$ the set of $*$ -subsets of X

- (i) $\mathcal{S}_*(X)$ is closed under intersection (i.e., $Y_\lambda \in \mathcal{S}_*(X) \ (\lambda \in A) \implies \bigcap_{\lambda \in A} Y_\lambda \in \mathcal{S}_*(X)$).
- (3) Let $x, y, x_1, \dots, x_n \in X$.
 - (i) The symbol $x * y$ denotes any of $x *_i y \ (i \in [m])$.
 - (ii) The symbol $x_1 * \dots * x_n$ denotes any product of x_1, \dots, x_n in this order under the (mixed) operations $*_i \ (i \in [m])$ and suitable $(\)$'s.
 - $\mathcal{S}_*(X) \ni Y \ni x_1, \dots, x_n \ (n \in \mathbb{Z}_{\geq 1}) \implies x_1 * \dots * x_n \in Y$
- (4) $A * B := \{a * b \mid a \in A, b \in B\} \equiv \{a *_i b \mid a \in A, b \in B, i \in [m]\} \quad (A, B \subset X)$
 $A_1 * \dots * A_n := \{a_1 * \dots * a_n \mid a_i \in A_i \ (i \in [n])\} \quad (n \in \mathbb{Z}_{\geq 1}, A_1, \dots, A_n \subset X)$

Notation 6.2. Let $S \subset X$.

- (1) The $*$ -subset of X generated by S means the smallest $*$ -subset of X including S . It exists and is obtained formally by $\langle S \rangle := \bigcap \{Y \in \mathcal{S}_*(X) \mid S \subset Y\}$.
- (2) The $*$ -subset $\langle S \rangle$ has the following explicite description : $\langle S \rangle = S^\infty$
 - (i) $S^n := \underbrace{S * \dots * S}_n \equiv \{x_1 * \dots * x_n \mid x_i \in S \ (i \in [n])\} \quad (n \in \mathbb{Z}_{\geq 1})$
 - Inductive definition :
 $S^1 = S \quad S^k \ (k \in [n]) \rightsquigarrow S^{n+1} = \bigcup \{S^k * S^\ell \mid k, \ell \in \mathbb{Z}_{\geq 1}, k + \ell = n + 1\}$
 - $S^2 = S * S, \quad S^3 = (S^1 * S^2) \cup (S^2 * S^1) = (S * (S * S)) \cup ((S * S) * S), \text{ etc.}$
 - (ii) $S^{\leq n} := \bigcup_{k \in [n]} S^k \quad (n \in \mathbb{Z}_{\geq 1}) \quad S^\infty := \bigcup_{n \in \mathbb{Z}_{\geq 1}} S^n$
 - (iii) In the unital case (i.e., X has the identity element e), we set $S^0 := \{e\}$ and $S^\infty := \bigcup_{n \in \mathbb{Z}_{\geq 0}} S^n$.

Fact 6.1.

- (1) (i) $S^k * S^\ell \subset S^{k+\ell} \quad (k, \ell \in \mathbb{Z}_{\geq 1})$ ◦ This implies that $S^\infty \in \mathcal{S}_*(X)$ and $\langle S \rangle = S^\infty$.
- (ii) $S^{\leq k} * S^{\leq \ell} \subset S^{\leq k+\ell} \quad (k, \ell \in \mathbb{Z}_{\geq 1})$
- (2) (i) $(S^k)^\ell \subset S^{k\ell}$ (ii) $(S^{\leq k})^{\leq \ell} \subset S^{\leq k\ell} \quad (k, \ell \in \mathbb{Z}_{\geq 1})$

6.2. Word length on a $*$ -set.

Definition 6.2. (Word length) Let $S \subset X$.

- (1) $\nu_S : X \longrightarrow \tilde{\mathbb{Z}}_{\geq 1} : \nu_S(x) = \begin{cases} \min\{n \in \mathbb{Z}_{\geq 1} \mid x \in S^n\} & (x \in S^\infty) \\ \infty & (x \in G - S^\infty) \end{cases}$
- (i) In the unital case $\mathbb{Z}_{\geq 1}$ is replaced by $\mathbb{Z}_{\geq 0}$.

Fact 6.2.

- (1) (i) $\nu_S(x) \leq n \iff x \in S^{\leq n} \quad (x \in X, n \in \mathbb{Z}_{\geq 1})$
- (ii) $\nu_S^{-1}(1) = S \quad (\nu_S^{-1}(1) = S - \{e\} \text{ in the unital case})$
- (2) (i) $\nu_S(x * y) \leq \nu_S(x) + \nu_S(y) \quad (x, y \in X)$
- (ii) $\nu_S(x_1 * \dots * x_n) \leq \nu_S(x_1) + \dots + \nu_S(x_n) \quad (x_i \in X \ (i \in [n])) \quad (n \in \mathbb{Z}_{\geq 1})$

Definition 6.3. (the induced metric on $\mathcal{S}(X)^\times = \mathcal{S}(X) - \{\emptyset\}$)

- (1) $\nu_H : \mathcal{S}(X)^\times \times \mathcal{S}(X)^\times \longrightarrow \tilde{\mathbb{Z}}_{\geq 1} : \nu_H(S, S') = \sup \nu_S(S')$
- (2) $\varrho := \log \nu_H : \mathcal{S}(X)^\times \times \mathcal{S}(X)^\times \longrightarrow \log \tilde{\mathbb{Z}}_{\geq 1} \subset \mathbb{R}_{\geq 0}$

Fact 6.3.

- (1) $\nu_H(S, S') \leq n \iff S' \subset S^{\leq n} \quad (S, S' \in \mathcal{S}(X)^\times, n \in \mathbb{Z}_{\geq 1})$

- (2) $\nu_H \in \mathcal{M}_{m,as}(\mathcal{S}(X)^\times)$, ν_H is nondegenerate.
- (i) $\nu_H(S, S') = 1 \iff S' \subset S$ (i)' $\nu_H(S, S') = \nu_H(S', S) = 1 \iff S = S'$
- (ii) $\nu_H(S, S'') \leq \nu_H(S, S')\nu_H(S', S'')$ ($S, S', S'' \in \mathcal{S}(X)^\times$)
- (3) $\varrho \in \mathcal{M}_{as}(\mathcal{S}(X)^\times)$, ϱ is nondegenerate. $\therefore \widehat{\varrho} \in \mathcal{M}(\mathcal{S}(X)^\times)$

Example 6.1. Some examples of subspaces $\mathcal{X} \subset \mathcal{S}(X)^\times$ with $\widehat{\varrho}|_{\mathcal{X}} < \infty$

Let $\mathcal{S}_g(X) := \{S \in \mathcal{S}(X) \mid X = S^\infty\} \subset \mathcal{S}(X)^\times$.

- (1) $\mathcal{S}_{fg}(X) := \{S \in \mathcal{S}_g(X) \mid |S| < \infty\}$
- (2) $\mathcal{S}_g(X; \text{finite rel } S_0) := \{S \in \mathcal{S}_g(X) \mid |S - S_0| < \infty, |S_0 - S| < \infty\}$ ($S_0 \in \mathcal{S}_g(X)$)
- (i) If $S \in \mathcal{S}_g(X)^\times$, $S' \in \mathcal{S}(X)^\times$ and $|S' - S| < \infty$, then $\nu_H(S, S') < \infty$

For word metrics on the $*$ -set X we have the following candidates.

Notation 6.3. (Word metrics) Let $S \subset X$.

$$d_S : X \times X \longrightarrow \widetilde{\mathbb{Z}}_{\geq 0} : d_S(x, y) = \begin{cases} \min\{n \in \mathbb{Z}_{\geq 0} \mid y \in [x, S, n]\} & (y \in \bigcup_{n \in \mathbb{Z}_{\geq 0}} [x, S, n]) \\ \infty & (\text{otherwise}) \end{cases}$$

Here, we can take the family $[x, S, n]$ ($x \in X, n \in \mathbb{Z}_{\geq 0}$) as follows :

- (1) $[x, S, n] := x * \underbrace{S * \cdots * S}_n$ ($n \in \mathbb{Z}_{\geq 1}$) $[x, S, 0] := \{x\}$ $\circ [x, S, n] \supset x * (S^n)$
- (2) $[x, S, n]' := ((\cdots ((x * S) * S) * \cdots * S) * S$ ($n \in \mathbb{Z}_{\geq 1}$) $[x, S, 0]' := \{x\}$

We denote by d'_S the metric d_S defined in the case (2).

Fact 6.4. $d_S, d'_S \in \mathcal{M}_{as}(X)$ and they are nondegenerate. $\therefore \widehat{d_S}, \widehat{d'_S} \in \mathcal{M}(X)$.

- (1) (i) $d_S(x, y) = 0 \iff x = y$ (ii) $d_S(x, z) \leq d_S(x, y) + d_S(y, z)$ ($\forall x, y, z \in X$)
- (2) (i) $d'_S(x, y) = 0 \iff x = y$ (ii) $d_S(x, z) \leq d_S(x, y) + d_S(y, z)$ ($\forall x, y, z \in X$)

6.3. Symmetry of a $*$ -set.

Let X be a $*$ -set.

Notation 6.4.

- (1) A map $f : X \rightarrow X$ is said to be a $*$ -endomorphism if $f(x *_i y) = f(x) *_i f(y)$ ($x, y \in X, i \in [m]$).
- \circ In this case, (i) $S \in \mathcal{S}_*(X) \implies f(S) \in \mathcal{S}_*(Y)$ (ii) $R \in \mathcal{S}_*(Y) \implies f^{-1}(R) \in \mathcal{S}_*(X)$
- (2) $\text{End}(X)$ denotes the monoid of $*$ -endomorphisms of X

Suppose (R, \star) is a nonempty set with a binary operation and $\varphi : R \longrightarrow \text{End}(X)$ is an anti-homomorphism.

Notation 6.5.

- (1) $x^a := \varphi(a)(x) \in X$ ($x \in X, a \in R$) (2) $S^A := \{x^a \mid x \in S, a \in A\}$ ($S \subset X, A \subset R$)
- (2) $S \subset X$ is φ -invariant $\stackrel{\text{def}}{\iff} S^a \subset S$ ($\forall a \in R$) $\iff S^R \subset S$

Fact 6.5. Let $x, y \in X$, $S \subset X$, $a, b \in R$ and $A, B \subset R$.

- (1) (i) $(x * y)^a = x^a * y^a$ (ii) $(x^a)^b = x^{a \star b}$ (iii) $(S^A)^B = S^{A \star B}$ (iv) $(S \cup T)^A = S^A \cup T^A$
- (2) (i) $(Y * Z)^a = Y^a * Z^a$ ($Y, Z \subset X$) (ii) $(\bigcup_{\lambda \in \Lambda} Y_\lambda)^a = \bigcup_{\lambda \in \Lambda} Y_\lambda^a$ ($Y_\lambda \subset X$ ($\lambda \in \Lambda$))

- (3) (i) $(x_1 * \cdots * x_n)^a = x_1^a * \cdots * x_n^a \quad (x_i \in X \ (i \in [n]))$
 (The products are taken in the same form in the left side and right side.)
 (ii) $(S^n)^a = (S^a)^n \quad (n \in \mathbb{Z}_{\geq 1}) \quad (S^\infty)^a = (S^a)^\infty \quad \langle S \rangle^a = \langle S^a \rangle$
 (4) (i) S^R is φ -invariant. (ii) $(S \cup S^R)^R = S^R \subset S \cup S^R \quad \therefore S \cup S^R$ is φ -invariant.
 (ii) S is φ -invariant. $\implies (S^n)^a \subset S^n \ (n \in \mathbb{Z}_{\geq 1}), (S^\infty)^a \subset S^\infty$

Fact 6.6. Suppose $S \in \mathcal{S}(X)^\times$ is φ -invariant.

- (1) $\nu_S(x^a) \leq \nu_S(x) \quad (\forall x \in X, \forall a \in R)$
 (2) $\nu_H(S, T^R) \leq \nu_H(S, T \cup T^R) = \nu_H(S, T) \quad (\forall T \in \mathcal{S}(X)^\times)$

Example 6.2. Further examples of subspaces $\mathcal{X} \subset \mathcal{S}(X)^\times$ with $\widehat{\varrho}|_{\mathcal{X}} < \infty$

- (1) $\mathcal{S}_{\varphi\text{-}fg}(X) := \{S \in \mathcal{S}_g(X) \mid S = F^R \text{ or } F \cup F^R \ (\exists F \in \mathcal{S}_f(X)^\times)\}$
 ◦ For any $S, T \in \mathcal{S}_{\varphi\text{-}fg}(X)$
 (i) $S \in \mathcal{S}_g(X)$, S is φ -invariant. (ii) $T = F^R$ or $F \cup F^R$ for some $F \in \mathcal{S}_f(X)^\times$.
 (iii) $\nu_H(S, F^R) \leq \nu_H(S, F \cup F^R) = \nu_H(S, F) < \infty \quad \therefore \nu_H(S, T) < \infty$
 (2) $\mathcal{S}_{\varphi\text{-}g}(X; \text{finite rel } F_0) := \{S \in \mathcal{S}_g(X) \mid S = F^R \text{ or } F \cup F^R \ (\exists F \in \mathcal{S}(X, \text{finite rel } F_0)^\times)\}$
 ◦ $\mathcal{S}(X; \text{finite rel } F_0)^\times := \{F \in \mathcal{S}(X)^\times \mid |F - F_0| < \infty, |F_0 - F| < \infty\} \quad (F_0 \in \mathcal{S}(X)^\times)$
 ◦ For any $S, T \in \mathcal{S}_{\varphi\text{-}g}(X; \text{finite rel } F_0)$
 (i) $S \in \mathcal{S}_g(X)$, S is φ -invariant. (ii) $S = F \cup F^R$ for some $F \in \mathcal{S}(X, \text{finite rel } F_0)^\times$.
 (iii) $T = G \cup G^R$ for some $G \in \mathcal{S}(X, \text{finite rel } F_0)^\times$.
 (iv) $G - S \subset G - F \subset (G - F_0) \cup (F_0 - F) \in \mathcal{S}_f(X)$
 $\therefore \nu_H(S, G) < \infty \quad (\text{cf. Example 6.1 (2)(i)})$
 (v) $\nu_H(S, G^R) \leq \nu_H(S, G \cup G^R) = \nu_H(S, G) < \infty \quad \therefore \nu_H(S, T) < \infty$

REFERENCES

- [1] D. Burago, S. Ivanov and L. Polterovich, *Conjugation-invariant norms on groups of geometric origin*, Advanced Studies in Pure Math., 52 (2008), Groups of diffeomorphisms, 221-250.
 [2] K. Fujiwara, *Quasi-homomorphisms on mapping class groups*, Handbook of Teichmüller theory. Vol. II, 241-269, IRMA Lect. Math. Theor. Phys., 13, Eur. Math. Soc., Zürich, 2009.
 [3] K. Fukui, T. Rybicki, T. Yagasaki, *The uniform perfectness of diffeomorphism groups of open manifolds*, arXiv math: 1905.07664.
 [4] K. Fukui and T. Yagasaki, *Boundedness of bundle diffeomorphism groups over a circle*, to appear in Jour. of Math. Soc. Japan, arXiv:2209.07848.
 [5] K. Fukui and T. Yagasaki, *Boundedness of diffeomorphism groups of manifold pairs – Circle case*, arXiv:2501.11363 [math.GT].
 [6] J. Kędra, A. Libman, and B. Martin, *Strong and uniform boundedness of groups*, J. Topol. Anal., 15 (3) (2023) 707-739.
 [7] J. Kędra, A. Libman, and B. Martin, *Uniform boundedness for algebraic groups and Lie groups*, arXiv:2202.13885 [math.GR].
 [8] M. Kawasaki, M. Kimura, H. Kodama, Y. Matsuda, T. Matsushita and R. Orita, *Relative simplicity of the universal coverings of transformation groups and Tsuboi's metric*, arXiv:2412.00839 [math.GR].
 [9] T. Tsuboi, *On the uniform perfectness of diffeomorphism groups*, Advanced Studies in Pure Math. 52, (2008), Groups of diffeomorphisms, 505-524.
 [10] T. Tsuboi, *On the uniform simplicity of diffeomorphism groups*, In Differential Geometry, World scientific Publ., Hackensack, N.J., (2009), 43-55.

- [11] T. Tsuboi, *On the uniform perfectness of the groups of diffeomorphism groups of even-dimensional manifolds*, Comm. Math. Helv., 87 (2012) 141-185.

PROFESSOR EMERITUS, KYOTO INSTITUTE OF TECHNOLOGY, KYOTO, JAPAN
Email address: yagasaki@kit.ac.jp